

## GENERAL INFORMATION

Suggested dosages refer to pounds of 7.5% Sevin Dust to use per acre of crop treated. Where a dosage range is indicated use the lower rate on small or young plants and the higher rate on large or mature plants. Always use sufficient dust to obtain good coverage of the crop plants being treated. Apply when insects or their damage appear and repeat applications as necessary, or at 7 to 10 day intervals unless shorter intervals are specified in DIRECTIONS FOR USE.

## DIRECTIONS FOR USE

**COTTON:** For early-season control of Thrips, Flea Beetles, Cotton Fleahopper, Cotton Leafworm, and Striped Blister Beetle apply 7 to 15 pounds.

For mid- and late-season control of Boll Weevil, Bollworm, Fall Armyworm, Cotton Leaf Perforator, Leaf Rollers, Leafhoppers, and Tarnished Plant Bug apply 15 to 25 pounds. Make applications every 5 to 7 days as long as control is necessary.

For control of Pink Bollworm apply 20 to 30 pounds on a 5 to 7 day schedule.

For control of Stink Bugs and Salt-Marsh Caterpillar apply 25 pounds.

Aphid populations will be suppressed by repeated applications of 7.5% Sevin Dust.

**SMALL FRUITS:** Blackberries, Boysenberries, Dewberries, Loganberries and Raspberries - Use 25 pounds for Japanese Beetles, Leaf Rollers, Leafhoppers and Raspberry Aphid. Do not apply later than 7 days before harvesting berries.

Grapes - Use 15 to 25 pounds for Grape Leaf Folder, Grape Leafhopper and Grape Leaf Skeletonizer. Apply just before first Brood Leaf Folder Larvae emerge from rolls and as needed for Leafhoppers. Use 25 pounds for Grape Berry Moth, Japanese Beetle, June Beetles and Red-Banded Leaf Roller.

Strawberries - Use 15 to 25 pounds for Meadow Spittlebug, Strawberry Leaf Roller and Strawberry Weevil. Do not apply later than 1 day before harvesting berries.

**VEGETABLES:** Asparagus - Use 15 to 25 pounds for Asparagus Beetle on seedlings or spears. Do not apply later than 1 day before harvesting asparagus. Use 30 to 50 pounds for Asparagus Beetle and Apache Cicada on ferns or brush growth in the post-harvest period. Do not apply more often than once every three days.

Beans (Green beans, lima beans, snap beans, cowpeas and black-eyed peas) - Use 7 pounds for Mexican Bean Beetle, 15 pounds for Bean Leaf Beetle, Cucumber Beetles, Flea Beetles, Japanese Beetle, Leafhoppers, Velvetbean Caterpillar and Westernbean Cutworm.

Use 15 to 20 pounds for Armyworms, Corn Earworm, Stink Bugs and Tarnished Plant Bug.

Corn - Use 15 to 25 pounds for Corn Earworm, Corn Rootworm Adults, European Corn Borer, Fall Armyworm, Flea Beetles, Japanese Beetle, Sap Beetles and Leafhoppers. For larvae in whorl, Budworm damage, and foliage feeders, apply to entire plant. Repeat as necessary.

For insects attacking silks and ears, apply at 2 to 3-day intervals starting when first silks appear and continuing until silks begin to dry. Three or more applications may be required depending on severity of infestation. Timing and good coverage are absolutely essential for effective control.

Cucumber, Melons, Pumpkin and Squash - Use 7 to 13 pounds for Pickleworm and Melonworm, 15 pounds for Cucumber Beetles, Flea Beetles, Leafhoppers and Squash Bugs. Avoid excessive applications. Some leaf injury may occur if treatments are made when tender foliage is wet or in the presence of prolonged high humidity. 7.5% Sevin Dust injures watermelons in Florida.

Garden Peas - Use 15 pounds for Leafhoppers. Use 15 pounds for Colorado Potato Beetle. For Colorado Potato Beetle control, direct application to weeds in fields of garden peas. This application is intended to help prevent insect contamination in processed peas. Use 15 to 20 pounds for Armyworms and Grasshoppers.

Potato, Tomato, Equolant and Pepper - Use 7 to 15 pounds for Colorado Potato Beetle, Flea Beetles and Leafhoppers. Use 15 to 25 pounds for European Corn Borer, Fall Armyworm, Lace Bugs, Tomato Fruitworm, Tomato Hornworm, Tarnished Plant Bugs and Stink Bugs.

ORZ - Use 15 to 25 pounds for Corn Earworm and Stink Bugs. Repeat at 5 to 7-day intervals or as necessary.

Carrots, Chinese Cabbage, Collards, Dandelion, Endive (Escarole), Garden Beets, Horse-radish, Kale, Lettuce, Mustard Greens, Parsley, Parsnips, Radishes, Rutabagas, Salsify, Spinach, Swiss Chard and Turnips - Use 7 to 15 pounds for Flea Beetles, Harlequin Bug and Leafhoppers. To control Six-Spotted Leafhopper (aster yellows vector), apply 15 to 20 pounds in a 5 to 7-day schedule.

Use 15 to 25 pounds for Armyworms, Cabbage Caterpillars, Corn Earworm, Tarnished Plant Bug and Stink Bugs. For optimum control of Corn Earworm after lettuce heads begin to form, use 25 pounds in a 5 to 7-day schedule.

Where Cabbage Looper is the principal problem, use an alternative pesticide that is specifically recommended. For control of small Cabbage Loopers after edible portions begin to form, use 15 to 25 pounds in a 5 to 7-day schedule.

Do not apply to Chinese Cabbage, collards, dandelion, endive (escarole), garden beets, kale, lettuce, mustard greens, parsley, salsify, spinach, Swiss chard or turnips later than 14 days before harvest, or to horseradish, parsnips, radishes, and rutabagas later than 3 days before harvest.

Cabbage, Broccoli, Brussels Sprouts, Cauliflower and Kohlrabi - Use 7 to 15 pounds for Flea Beetles and Harlequin Bug.

Use 15 to 25 pounds for Armyworms, Cabbage Caterpillars and Corn earworm. Where Cabbage Looper is the principal problem, use an alternative pesticide that is specifically recommended. For control of small Cabbage Loopers after edible portions begin to form, use 7.5% Sevin Dust in a 5 to 7-day schedule. Do not apply to these crops later than 3 days before harvest.

**TOBACCO:** In Plant Beds (For Tobacco Flea Beetle) - Apply 1 to 2 pounds per 100 square yards when insects appear. Avoid excessive application; plant injury may result. Repeat as needed at 10 to 14-day intervals and just before transplanting.

In Fields, Broadcast applications for Hornworms and Tobacco Flea Beetles - Apply 10 to 15 pounds per acre depending on size of the plants. Good coverage is essential. Use the highest dosage only on large plants and repeat as needed. Heavier applications may cause plant injury.

If late applications are necessary, treat immediately after priming rather than before priming. Treated fields may be entered immediately after dust has settled. Some phytotoxicity may occur on tender foliage in the presence of rain or high humidity of several days duration following application.

**FORAGE AND FIELD CROPS:** Alfalfa, Clovers, Cowpeas, Peanuts and Soybeans - Use 7 to 15 pounds for Blister Beetles and Mexican Bean Beetle.

Use 15 pounds for Alfalfa Caterpillar, Bean Leaf Beetle, Cucumber Beetles, Green Cloverworm, Japanese Beetle, Leafhoppers, Three-cornered Alfalfa Hopper, Thrips and Velvetbean Caterpillar.

Use 15 to 20 pounds for Armyworms, Corn Earworm, Stink Bugs and Webworms. To avoid possible injury on tender foliage, do not apply when foliage is wet or when rain or excessive humidity is expected during the next two days.

Forage Grasses and Pasture - Use 15 to 20 pounds for Armyworms and Thrips. For optimum Thrips control in grasses grown for seed, direct dust into boot.

Rice - In the Mississippi Delta and Texas - Use 15 to 20 pounds for Armyworms and Stink Bugs. Warning! Do not apply before heading if propanil (such as Staff F-34 or Rogue) has been or will be applied; plant injury may result. Do not apply later than 14 days before harvesting rice.

Sorghums - Including Milo and Grain Sorghum - Use 15 to 25 pounds for Armyworms, Corn Earworm, Stink Bugs and Webworms. Use 20 pounds for Sorghum Midge. For optimum insect control on grain sorghum, direct dust into the forming heads. Treatment for Sorghum Midge control should be made 3 to 4 days after heads have emerged from boot. Do not apply later than 21 days before harvesting sorghums.

**GRASSHOPPERS:** Apply 7 to 20 pounds as often as necessary to control grasshoppers on the above crops. A 7 to 15 pounds rate is suggested for nymphs on small plants or sparse vegetation in wasteland, rangeland, ditchbanks, and borders. A 15 to 20 pound dosage should be used when grasshoppers are mature or material is applied to crops requiring greater coverage.

Net Weight:  
50 Pounds

# 7.5%

7.5% Sevin Dust is  
of certain small fruit  
and field crops.

READ ENTIRE

ACTIVE INGREDIENT  
Carbaryl (1-naphthyl  
INERT INGREDIENTS

KEEP OUT OF REACH

HARMFUL IF INHALED  
Repeated Breathing  
Prolonged or Repeated  
Before Eating. Take  
Long Sleeved Work

AVOID CONTAMINATION  
TROUGH

NOTE FOR PHYSICIANS  
cholinesterase inhibitor

Sold By



Harold

ACCEPTED

100