

GENERAL INFORMATION

Net Contents
Gallons

TOP-CROP

FORM
REV. 7-54

TYPES OF TREATMENT

Pre-emergence Treatment

PREMERGE is used to prevent the growth of weeds and grasses before they appear. It is applied to the soil before the crop is planted. It is most effective when applied to the soil before the crop is planted. It is most effective when applied to the soil before the crop is planted.

Post-emergence Treatment

PREMERGE can be used to control weeds and grasses that have already appeared. It is applied to the soil after the crop has been planted. It is most effective when applied to the soil after the crop has been planted.

DIRECTIONS FOR USE

PREMERGE is used to control weeds and grasses before they appear. It is applied to the soil before the crop is planted. It is most effective when applied to the soil before the crop is planted.

NOTE: Always use PREMERGE in accordance with the directions on the label. Do not use PREMERGE on crops listed in the "Do Not Use" section of the label. Do not use PREMERGE on crops listed in the "Do Not Use" section of the label.

PEANUTS: Do not use PREMERGE on peanuts. Do not use PREMERGE on peanuts. Do not use PREMERGE on peanuts.

SOYBEANS: Do not use PREMERGE on soybeans. Do not use PREMERGE on soybeans. Do not use PREMERGE on soybeans.

MAIZE BEANS, SNAP BEANS, FIELD BEANS: Do not use PREMERGE on maize beans, snap beans, or field beans. Do not use PREMERGE on maize beans, snap beans, or field beans.

VEGETABLES: Do not use PREMERGE on vegetables. Do not use PREMERGE on vegetables. Do not use PREMERGE on vegetables.

FRUIT TREES: Do not use PREMERGE on fruit trees. Do not use PREMERGE on fruit trees. Do not use PREMERGE on fruit trees.

WATERMELONS: Do not use PREMERGE on watermelons. Do not use PREMERGE on watermelons. Do not use PREMERGE on watermelons.

CUCUMBERS: Do not use PREMERGE on cucumbers. Do not use PREMERGE on cucumbers. Do not use PREMERGE on cucumbers.

ALFALFA: Do not use PREMERGE on alfalfa. Do not use PREMERGE on alfalfa. Do not use PREMERGE on alfalfa.

STRAWBERRIES: Do not use PREMERGE on strawberries. Do not use PREMERGE on strawberries. Do not use PREMERGE on strawberries.

MINT: Do not use PREMERGE on mint. Do not use PREMERGE on mint. Do not use PREMERGE on mint.

FERNAL POTATOES, VINES: Do not use PREMERGE on fernal potatoes or vines. Do not use PREMERGE on fernal potatoes or vines.

GENERAL CONTACT WEED CONTROL: Do not use PREMERGE for general contact weed control. Do not use PREMERGE for general contact weed control.

FIELD ORNAMENTALS: Do not use PREMERGE on field ornamentals. Do not use PREMERGE on field ornamentals. Do not use PREMERGE on field ornamentals.

WATERMELONS: Do not use PREMERGE on watermelons. Do not use PREMERGE on watermelons. Do not use PREMERGE on watermelons.

PREMERGE is used to control weeds and grasses before they appear. It is applied to the soil before the crop is planted. It is most effective when applied to the soil before the crop is planted.

PEAS

Do not use PREMERGE on peas. Do not use PREMERGE on peas. Do not use PREMERGE on peas.

OATS, RYE, WHEAT AND BARLEY SEEDS ALONE OR INTERPLANTED WITH A LEGUME: Do not use PREMERGE on oats, rye, wheat, or barley seeds alone or interplanted with a legume. Do not use PREMERGE on oats, rye, wheat, or barley seeds alone or interplanted with a legume.

SEEDING ALFALFA, SWEET CLOVER, RED CLOVER, BIRDFOOT TREFLOIL: Do not use PREMERGE on seeding alfalfa, sweet clover, red clover, or birdfoot trefloil. Do not use PREMERGE on seeding alfalfa, sweet clover, red clover, or birdfoot trefloil.

PUMPKINS AND SQUASH: Do not use PREMERGE on pumpkins or squash. Do not use PREMERGE on pumpkins or squash.

CUCUMBERS: Do not use PREMERGE on cucumbers. Do not use PREMERGE on cucumbers. Do not use PREMERGE on cucumbers.

ALFALFA: Do not use PREMERGE on alfalfa. Do not use PREMERGE on alfalfa. Do not use PREMERGE on alfalfa.

STRAWBERRIES: Do not use PREMERGE on strawberries. Do not use PREMERGE on strawberries. Do not use PREMERGE on strawberries.

MINT: Do not use PREMERGE on mint. Do not use PREMERGE on mint. Do not use PREMERGE on mint.

FERNAL POTATOES, VINES: Do not use PREMERGE on fernal potatoes or vines. Do not use PREMERGE on fernal potatoes or vines.

GENERAL CONTACT WEED CONTROL: Do not use PREMERGE for general contact weed control. Do not use PREMERGE for general contact weed control.

FIELD ORNAMENTALS: Do not use PREMERGE on field ornamentals. Do not use PREMERGE on field ornamentals. Do not use PREMERGE on field ornamentals.

WATERMELONS: Do not use PREMERGE on watermelons. Do not use PREMERGE on watermelons. Do not use PREMERGE on watermelons.

CUCUMBERS: Do not use PREMERGE on cucumbers. Do not use PREMERGE on cucumbers. Do not use PREMERGE on cucumbers.

ALFALFA: Do not use PREMERGE on alfalfa. Do not use PREMERGE on alfalfa. Do not use PREMERGE on alfalfa.

STRAWBERRIES: Do not use PREMERGE on strawberries. Do not use PREMERGE on strawberries. Do not use PREMERGE on strawberries.

PREMERGE

Dinitro Weed Killer

PREMERGE is used to control weeds and grasses before they appear. It is applied to the soil before the crop is planted. It is most effective when applied to the soil before the crop is planted.

READ ENTIRE LABEL BEFORE USING THIS PRODUCT

INGREDIENTS

- Active ingredients: 2,4-D, 2,4,5-T, 2,4,6-T, 2,4,7-T, 2,4,8-T, 2,4,9-T, 2,4,10-T, 2,4,11-T, 2,4,12-T, 2,4,13-T, 2,4,14-T, 2,4,15-T, 2,4,16-T, 2,4,17-T, 2,4,18-T, 2,4,19-T, 2,4,20-T, 2,4,21-T, 2,4,22-T, 2,4,23-T, 2,4,24-T, 2,4,25-T, 2,4,26-T, 2,4,27-T, 2,4,28-T, 2,4,29-T, 2,4,30-T, 2,4,31-T, 2,4,32-T, 2,4,33-T, 2,4,34-T, 2,4,35-T, 2,4,36-T, 2,4,37-T, 2,4,38-T, 2,4,39-T, 2,4,40-T, 2,4,41-T, 2,4,42-T, 2,4,43-T, 2,4,44-T, 2,4,45-T, 2,4,46-T, 2,4,47-T, 2,4,48-T, 2,4,49-T, 2,4,50-T, 2,4,51-T, 2,4,52-T, 2,4,53-T, 2,4,54-T, 2,4,55-T, 2,4,56-T, 2,4,57-T, 2,4,58-T, 2,4,59-T, 2,4,60-T, 2,4,61-T, 2,4,62-T, 2,4,63-T, 2,4,64-T, 2,4,65-T, 2,4,66-T, 2,4,67-T, 2,4,68-T, 2,4,69-T, 2,4,70-T, 2,4,71-T, 2,4,72-T, 2,4,73-T, 2,4,74-T, 2,4,75-T, 2,4,76-T, 2,4,77-T, 2,4,78-T, 2,4,79-T, 2,4,80-T, 2,4,81-T, 2,4,82-T, 2,4,83-T, 2,4,84-T, 2,4,85-T, 2,4,86-T, 2,4,87-T, 2,4,88-T, 2,4,89-T, 2,4,90-T, 2,4,91-T, 2,4,92-T, 2,4,93-T, 2,4,94-T, 2,4,95-T, 2,4,96-T, 2,4,97-T, 2,4,98-T, 2,4,99-T, 2,4,100-T.





POISON



SEE ANTIDOTE AND WARNING STATEMENTS IN RIGHT PANEL

DANGER

KEEP OUT OF THE REACH OF CHILDREN



DANGER

ABSORBED THROUGH THE SKIN-MAY BE FA



POISON

FIRST AID TREATMENT - ANT

SYMPTOMS OF POISONING

SYMPTOMS OF POISONING: Nausea, vomiting, diarrhea, abdominal pain, dizziness, headache, weakness, loss of consciousness, respiratory depression, convulsions, coma, death.

CAUTION

NOTICE



HARDEN THORNTON AND COMPANY

