

# RIVERSIDE 25% CHLORVALNE GRANULES

**FOR CONTROL OF CERTAIN INSECT PESTS OF CORN, PEANUTS, TOBACCO, VEGETABLE CROPS, FRUIT AND BERRY CROPS, ORNAMENTAL TURF, LAWNS, NURSERY AND GREENHOUSE SOILS**

ACTIVE INGREDIENT:	25.0%
Technical Chlorvalne*	
INERT INGREDIENTS	75.0%
	Total 100.0%

\*Equivalent to 15% Oxytetrachlor 4, 7-methanotetrahydrodane and 10% related compounds.

## CAUTION: KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Contact with skin can cause toxic symptoms. Avoid breathing dust. In case of contact with skin, wash with soap and water. Avoid contamination of feed and foodstuffs.

Keep children and pets off treated turf until material has been washed into the soil by watering or rain, and the grass is dry.

This product is toxic to fish and wildlife. Birds feeding on treated areas may be harmed.

Keep out of lakes, ponds, and streams. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of waste.

Do not apply to any area not specified on the label.

**READ ENTIRE LABEL BEFORE USING THIS PRODUCT:** Use of this product in accordance with the information on this label may injure or kill humans, domestic animals, wildlife, aquatic life, desirable plants, create undesirable and/or undesirable residues, or produce other undesirable results.

**ANTIDOTE:**  
In case of eye contact, flush with water for 15 minutes. In case of ingestion, give 1 tablespoonful of mustard in a glass of water.

In case of spillage on skin, wash promptly with soap and water.

### CONTAINER DISPOSAL:

Do not reuse empty container. Containers should be disposed of by punching holes in them and burying with wastes, or by burning. Stay out of smoke.

### NOTICE:

Riverside Chemical Company warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to Riverside Chemical Company, and buyer assumes the risk of any such use.

### DIRECTIONS FOR USE:

See chart below for specific crops to be treated and rates of application.

**FIELD AND VEGETABLE CROPS:** Make preplant broadcast or row (before) application to soil after plowing and disc, or incorporate into soil immediately.

**BERRY CROPS:** Make application at base of plants and incorporate into soil.

Do not apply at each irrigation.  
**ORNAMENTAL TURF AND LAWN, NURSERY AND GREENHOUSE SOILS:** Apply granules uniformly to soil surface, followed by watering. For lawn applications, apply after mowing and before watering. For nursery applications, apply to soil surface and water in.

CROP (OR USE)	INSECTS CONTROLLED	POUNDS PER 1000 SQ. FT. OF SOIL	
		Row	Broadcast
Corn (field and Sweet) Peanuts, Tobacco	Corn Rootworms (Northern and Southern), Corn Field Aids, Corn Leaf Aids, Cutworms	4	5
	Soil Bump Beetle Larvae, Seed Corn Maggot, White Grubs, Wireworms, Japanese Beetle Larvae Green June Beetle Larvae	10	10-15
Vegetable Crops Beans, Beets, Black-eyed Peas, Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Cauliflower, Celery, Collards, Cucumbers, Eggplants, Kale, Kohlrabi, Lettuce, Melons, Onions, Peas, Pepper, Potatoes, Radishes, Rutabagas, Squash, Sweet Potatoes, Tomatoes, Turnips, Watermelons Fruit & Berry Crops: Apples, Apricots, Cherries, Nectarines, Pears, Plums, Prunes, Quinces, Blackberries, Grapes, Huckleberries, Loganberries, Raspberries, Strawberries, Grapefruit, Mangoes, Lemons, Limes, Oranges, Tangerines, Tangelos, Pineapple Blackberries, Blueberries, Raspberries, Strawberries	Asian Garden Beetle Larvae, Whitefringed Beetle Larvae Cutworms	10	10-15
	Wireworms, White Grubs	8	10-15
	Asian Garden Beetle Larvae Whitefringed Beetle Larvae		10-15
	Japanese Beetle Larvae		10-15
	Flea Beetle Larvae (potatoes and sweet potatoes)		10-15
	Aids		10-15
	White Grub, Cutworms, Wireworms		10-15
	Japanese Beetle Larvae		10-15
	Green June Beetle Larvae		10-15
	Asian Garden Beetle Larvae, Whitefringed Beetle Larvae Strawberry Root Weevil Larvae		10-15
Ornamental Turf, Lawns, Nursery and Greenhouse Soils	Aids, Fleas, Fleas, Crickets, European Larvae, Ground Squirrels, Mole Crickets, Aids, Flea, Chafers, Spiders	0.2/1000 sq. ft.	0.2/1000 sq. ft.
	Asian Garden Beetle Larvae, European Chafers Larvae, Whitefringed Beetle Larvae, White Grubs, Wireworms	1/1000 sq. ft.	1/1000 sq. ft.
	Japanese Beetle Larvae	1/1000 sq. ft.	1/1000 sq. ft.

\*With a range of rates is given, use the higher rate of application on much or highly organic soils.

Where row treatment is desirable, application should be made at planting time using suitable row equipment.

**NOTE:** A 7-10 inch band is recommended for control of White Grubs and Wireworms in vegetables; row treatment of corn should be in a 7 inch band. Minimum rates should be based on no less than a 7 inch band. For best control a broadcast application is recommended.

EPA REG. NO. 507-113 NET CONTENTS



**RIVERSIDE CHEMICAL CO.**  
CI  
Riverside, California  
Manufactured for Agriculture, Agriculture & Industry

**ACCEPTED**  
9/26/72  
9774-188  
TO ATTACHED COMMENTS