

RIVERSIDE 10% CHLORDANE GRANULES

FOR CONTROL OF CERTAIN INSECT PESTS OF CORN, PEANUTS, TOBACCO, VEGETABLE CROPS, FRUIT AND BERRY CROPS, ORNAMENTAL TURF, LAWNS, NURSERY AND GREENHOUSE SOILS

ACTIVE INGREDIENT:	Technical Chlordane*	10.0%
INERT INGREDIENTS		90.0%
	Total	100.0%

*Equivalent to 6% Octachloro-4, 7-methanotetrahydroindane and 4% related compounds.

CAUTION: KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Contact with skin can cause toxic symptoms. Avoid breathing dust. In case of contact with skin, wash with soap and water. Avoid contamination of feed and foodstuffs.

Keep children and pets off treated turf until material has been washed into the soil by watering or rain, and the grass is dry.

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed.

Keep out of lakes, ponds, and streams. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes.

Do not apply to any area not specified on the label.

READ ENTIRE LABEL BEFORE USING THIS PRODUCT: Use of this product in a manner not in accordance with the information on this label may injure or kill humans, domestic animals, wildlife, aquatic life, desirable plants, create excessive and unlawful chemical residues, or produce other undesirable results.

ANTIDOTE:

INTERNAL — Give an emetic of 1 tablespoonful of mustard in a glass of water. Call a physician.

EXTERNAL — In case of spillage on skin wash promptly with soap and water

CONTAINER DISPOSAL

Do not reuse empty container. Containers should be disposed of by punching holes in them and burying with wastes, or by burning. Stay out of smoke.

NOTICE

Riverside Chemical Company warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to Riverside Chemical Company, and buyer assumes the risk of any such use.

DIRECTIONS FOR USE

See right panel for specific crops to be treated and rates of application.

FIELD AND VEGETABLE CROPS: Make preplant broadcast or row (where applicable) application to soil after plowing and disc or incorporate into soil immediately.

BERRY CROPS: Make application at base of plants and incorporate into soil.

Do not apply when fruit is present.

ORNAMENTAL TURF AND LAWN, NURSERY AND GREENHOUSE SOILS: Apply granules uniformly, using a lawn fertilizer spreader or other suitable granular application equipment. Incorporate into bare soil immediately.

TREE FRUITS: Make soil application only when no fruits are present.

CROP (OR USE)	INSECTS CONTROLLED	POUNDS OF RIVERSIDE 10% CHLORDANE GRANULES PER ACRE		
		Row	Broadcast	
Corn (field and sweet) Peanuts, Tobacco	Corn Rootworms (Northern and Southern), Corn Field Ants, Corn Root Aphids	10	20	
	Cutworms	25	25-40	
	Seed Corn Beetle Larvae, Seed Corn Maggot, White Grubs, Wireworms	25	25-40	
	Japanese Beetle Larvae		40	
	Green June Beetle Larvae		30-40	
	Asiatic Garden Beetle Larvae, Whitefringed Beetle Larvae		30-40	
	Cutworms		25-40	
	Wireworms, White Grubs	20	25-40	
	Asiatic Garden Beetle Larvae		25-50	
	Whitefringed Beetle Larvae		25-50	
Vegetable Crops Beans, Beets, Blackeyed Peas, Broccoli, Brussels Sprouts, Cabbage, Cantaloupes, Cauliflower, Celery, Collards, Cucumbers, Eggplants, Kale, Kohlrabi, Lettuce, Melons, Okra, Onions, Peas, Peppers, Pimientos, Potatoes, Radishes, Rutabagas, Squash, Sweet Potatoes, Tomatoes, Turnips, Watermelons	Japanese Beetle Larvae		50-100	
	Flea Beetle Larvae (potatoes and sweet potatoes)		25-50	
	Ants		15-20	
	White Grubs, Cutworms, Wireworms		25-40	
	Japanese Beetle Larvae		100	
	Green June Beetle Larvae		30-50	
	Asiatic Garden Beetle Larvae, Whitefringed Beetle Larvae		25-50	
	Strawberry Root Weevil Larvae		50	
	Fruit & Berry Crops: Apples, Apricots, Cherries, Nectarines, Peaches, Plums, Prunes, Quinces, Blackberries, Grapes, Huckleberries, Loganberries, Raspberries, Strawberries, Grapefruit, Kumquats, Lemons, Limes, Oranges, Tangerines, Tangelos, Pineapple	Blackberries, Blueberries, Raspberries, Strawberries		50
		Ornamental Turf, Lawns, Nursery and Greenhouse Soils	Ants, Ticks, Chiggers	6
Crickets, European Earwigs, Grasshoppers, Mosquitoes (Adult), Rose Chafer, Spiders			9-18	25-50
Asiatic Garden Beetle Larvae, European Chafer Larvae, White Fringed Beetle Larvae, White Grubs, Wireworms			18-23	50-100
Japanese Beetle Larvae			23	100

*Where a range of rates is given, use the higher rate on muck or highly organic soils.

Where row treatment is desirable, application should be made at planting time using suitable row equipment.

NOTE: A 7-10 inch band is recommended for control of White Grubs and Wireworms in vegetables; row treatment of corn should be in a 7 inch band. Minimum rates should be based on no less than a 7 inch band. For best control a broadcast application is recommended.

EPA REG. NO. 9779-198

NET CONTENTS: _____ POUNDS



MANUFACTURED BY
RIVERSIDE CHEMICAL CO.
DIVISION OF COOK INDUSTRIES, INC., MEMPHIS, TENN. 38116

Riverside serves agriculture. Agriculture serves everyone.

