

RIVERSIDE WEEDGARD

NONSELECTIVE WEED AND GRASS KILLER.

CONTAINS SODIUM METABORATE, SODIUM CHLORATE, AND DUPONT BROMACIL.

USEFUL AS DRY OR SPRAY APPLICATION FOR GENERAL CONTROL OF GRASSES AND BROADLEAVED WEEDS ON INDUSTRIAL SITES OR SIMILAR NON-CROPPED LAND.

ACTIVE INGREDIENTS:

Sodium Metaborate Tetrahydrate (Na ₂ B ₄ O ₇ ·4H ₂ O)	88.5%
Sodium Chlorate (NaClO ₃)	30.0%
Bromacil (5-bromo-3-sec-butyl-6-methyluracil)	1.5%
INERT INGREDIENTS:	2.0%

U.S. Patents 2,700,604 & 3,227,541. Other Patents Pending

CAUTION: KEEP OUT OF REACH OF CHILDREN

Avoid contamination of feed and foodstuffs. Do not leave container where children or animals may take internally. Harmful if swallowed. May cause irritation of eyes, nose, throat and skin. Avoid breathing spray mist. Avoid contact with skin and clothing. Do not store in direct contact with ground or concrete floors.

READ ENTIRE LABEL BEFORE USING THIS PRODUCT: Use of this product in a manner not in accordance with the information on this label may injure or kill humans, domestic animals, wildlife, aquatic life, desirable plants, create excessive and unlawful chemical residues, or produce other undesirable results.

GENERAL INFORMATION

RIVERSIDE WEEDGARD G is designed for kill of grasses and broadleaved weeds. It may be used for quick knock-down of existing vegetation as well as for long lasting control.

RIVERSIDE WEEDGARD G should be used on non-cropped land such as industrial sites, fence rows, storage yards, roadsides and railroads.

RIVERSIDE WEEDGARD G may be applied at any time during the growing season. Unwanted vegetation is easiest to kill when it is young and growing vigorously.

RIVERSIDE WEEDGARD G is unique in that it may be applied dry as is to the soil surface or it may be dissolved in water and applied to growing unwanted vegetation as a spray.

DRY APPLICATIONS may be made with a shaker can or with a hand operated spreader designed for this use. Power spreaders can be used for large areas. Effects are slow to appear and will not be apparent until the product is activated by moisture.

SPRAY APPLICATIONS should be made with equipment having mechanical agitation. Up to 3 pounds of RIVERSIDE WEEDGARD G may be dissolved in a gallon of water. Add RIVERSIDE WEEDGARD G gradually to the spray tank during filling with agitator operating. Make sure RIVERSIDE WEEDGARD G granules are all dissolved before spraying. Spray so as to thoroughly wet all surfaces of the vegetation to be controlled.

For small areas, or for spot treatment, application can be made with a knapsack type sprayer or sprinkling can.

DIRECTIONS FOR USE

GRASS AND BROADLEAVED WEED CONTROL

ANNUAL WEEDS AND GRASSES: For control of annuals such as sunflower, Kochia, Russian thistle, pigweed, crabgrass, downy brome, foxtail, ryegrass, wild oats, ragweed, lambquarters, puncturevine, and turkey mullein apply 1 pound per 100 square feet for a dry application; for spray application dissolve RIVERSIDE WEEDGARD G at the rate of 1 pound in a gallon of water and apply this solution at the rate of 1 gallon per 100 square feet.

PERENNIAL WEEDS AND GRASSES: For control of perennials such as smooth brome, Bahiagrass, bluegrass, redtop, purpletop, quackgrass, broom-sedge, aster, dandelion, dog fennel, goldenrod, plantain, wild carrot, Johnson grass, Bermuda grass, Dallisgrass, Canada thistle, field bindweed, apply 2-3 pounds per 100 square feet for a dry application; for spray application dissolve RIVERSIDE WEEDGARD G at the rate of 2-3 pounds per gallon of water and apply this solution at the rate of 1 gallon per 100 square feet.

IMPORTANT

This material is not selective in action and application and may render the entire treated area for one or more years. Care should be taken to the particular area intended to be destroyed or injured. This includes lawns, trees, shrubs, crops and other plants which may be underlain by roots of adjacent plants. Do not drain such areas. Do not contaminate domestic water supplies by fish and wildlife. Wash clothing and animals off treated areas until rainfall has soil. Mechanical equipment should be thoroughly cleaned.

DISPOSAL OF CONTAINER

Burn bag immediately when empty. Stay out of area.

NOTICE

Seller warrants that the product conforms reasonably fit for the purposes stated on the label with directions under normal conditions of use. No warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product, or under abnormal conditions, or for use for any purpose not intended or foreseeable to seller, and buyer assumes the responsibility.


EPA REG. NO. 9779-209

MANUFACTURED FOR



MANUFACTURED BY
RIVERSIDE
A SUBSIDIARY OF COOK

Riverside serves agriculture



9779-209

80411692

FOR LAKESHORE, DRIVEWAYS AND SIDEWALKS,
 where lasting residual effect is desired.

ACTIVE INGREDIENTS:

- Sodium chlorate (NaClO₃).....40.0%
- Sodium metaborate (Na₂B₂O₄ • H₂O)*..... 57.0%
- Monuron (3-(p-chlorophenyl)-1, 1-dimethylurea** 1.0%


INERT INGREDIENTS..... 2.0%

*Equivalent to 5.14% boron expressed as elemental boron.

**Equivalent to 1.25% Telvar.

"Telvar," a trademark of E. I. duPont de Nemours & Company, Inc.

U.S. Patent Nos. 2,847,293 and 2,655,445. Other Patents Pending



GENERAL RECOMMENDATIONS:

AMOUNTS TO USE: For annuals or shallow-rooted plants—Apply 1 to 2 pounds of PRO-2 per 100 square feet.

FOR PERENNIALS AND DEEP-ROOTED PLANTS—Apply 2 to 3 pounds per 100 square feet. For first application where Johnson, Bermuda and Vasey grasses are a serious problem apply 3 to 4 pounds per 100 square feet.

EXAMPLES OF USES: PRO-2 may be used wherever noncropland weed control is desired, and in any areas where vegetation is not wanted (see CAUTION below). Examples are as follows:

Lakeshore: Along the edges of lakes to kill unwanted grass, weeds, and brush.

Sidewalks, driveways and patios: Kill off unwanted vegetation.

Others: Airfields, runways and landing lights; outdoor theaters; tennis courts; school playgrounds; driveways and parking areas.

APPLICATION

NOTE: It is essential that PRO-2 be spread uniformly and at the rates given above.

Sprinkler can application preferred to avoid drift. Clean thoroughly after use. Agitate when mixing PRO-2 with water. Apply to give thorough coverage.

DRY—Use undiluted and apply uniformly by hand sifter, fertilizer spreader or by hand.

IMPORTANT—The killing action of PRO-2 is primarily through the roots of plants, therefore, moisture is required after application. Too little moisture will not carry the chemical throughout the root zone and follow up applications may be necessary.

WHEN TO APPLY

For Annual Weeds and Grasses, it is best to apply in the spring when weeds are small.

For Perennial Weeds and Grasses, in areas of light winter rain (less than 20 inches) it is generally best to treat during fall or winter. In areas of more than 20 inches, spring application is effective.

General Note—Avoid application in hot dry weather as moisture is required to activate the chemical. By preventing weed growth in the year, fire hazards from dead and dry vegetation are avoided later on.

Cautions in Application—Do not apply on or near turf, valuable plants, or in areas into which their roots may extend. Do not use on slopes where "run-off" will move the chemical into the area of useful plants. The chemical should be used only where the soil is not objectionable. Avoid contamination of irrigation or domestic water.

CAUTIONS: Harmful if swallowed. May cause irritation of eyes, throat and skin. Avoid handling with cuts or abrasions on hands. Wear gloves. Avoid breathing dust. Avoid contact with skin, eyes, clothing. Wash clothing and equipment after application. Do not use in containers where children and animals may take the chemical. Avoid contamination of food and foodstuff.

Local Conditions—may affect the use of this chemical. Consult your experiment station or extension service weed specialist for specific recommendations as to dosage and time of application and control of other weeds and grasses.

WHEN TO APPLY

and Grasses, it is best to apply in the spring when

eds and Grasses, in areas of light winter rainfall
(es) it is generally best to treat during fall or early
f more than 20 inches, spring application is most

aid application in hot dry weather as moisture is
to the chemical. By preventing weed growth early
hazards from dead and dry vegetation are avoided

caution—Do not apply on or near turf, valuable trees
r in areas into which their roots may extend. Do not
ere "run-off" will move the chemical into the root
ants. The chemical should be used only where un-
not objectionable. Avoid contamination of irrigation

ful if swallowed. May cause irritation of eyes, nose,
Avoid handling with cuts or abrasions on hands or
d breathing dust. Avoid contact with skin, eyes and
othing and equipment after application. Do not leave
children and animals may take the chemical in-
ntemination of food and foodstuff.

—may affect the use of this chemical. Consult ex-
er extension service weed specialist for specific
es to dosage and time of application and for con-
s and grasses.

CAUTION

See Caution Statements on Back Panel

KEEP OUT OF REACH OF CHILDREN

**DO NOT STORE WITH FOODSTUFFS
STORE IN A COOL, DRY PLACE**

Avoid Direct Contact with Ground or Concrete Floors

Distributed By

BIRCH INDUSTRIES, INC.

6521 Central Avenue N.E.
MINNEAPOLIS, MINN. 55432

USDA Reg. No. 9798-2

DEATH !! TO

LAKESHORE WEEDS

BRUSH AND GRASS

5 LBS. TREATS UP TO 500 SQ. FT.

Net Weight 5 Pounds

PRO-2

WEED & GRASS KILLER

Contains "TELVAR" a trademark of E.I. DuPont
de Nemours & Co., Inc.