

PM 13 9779-210

F. 1/6/33



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

MAR 24 1994

Glenda Haage
RIVERSIDE/TERRA CORP.
600 Fourth Street
Sioux City, IA 51101

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Subject: Label Amendment Submission of 01/05/94 in Response to PR Notice 93-7
EPA Reg. No. 9779-210
RIVERSIDE DIAZINON 4AG

Dear Registrant:

The labeling cited above and submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is accepted subject to the comments reflected on the enclosed sheet. A copy of your proposed labeling stamped "ACCEPTED WITH COMMENTS" is enclosed.

WHAT THIS ACCEPTANCE MEANS:

Based on your certification, the Agency has accepted the labeling changes that are necessary to comply with the Worker Protection Standard (WPS) labeling requirements of 40 CFR part 156, subpart K, described in PR Notices 93-7 and 93-11. Any other labeling changes submitted in connection with this amendment application but not directly related to compliance with the WPS have not been reviewed or accepted by the Agency. If you wish to make such changes, you must submit a separate amendment application proposing them. If your product is currently suspended, the acceptance of this labeling amendment does not affect the suspension in any way.

WHAT YOU NEED TO DO NEXT:

By the next label printing make all the specified changes to your labeling. Send to EPA one (1) copy of the final printed labeling:

- BEFORE selling or distributing any product bearing the final printed labeling
- AND
- WITHIN one year from date of this acceptance.



Recycled/Recyclable
Printed with Soy/Canola Ink on paper that
contains at least 50% recycled fiber


Submit the final printed labeling via the U.S. Postal Service to:

Document Processing Desk (FIN-LABEL)
Office of Pesticide Programs (7505C)
U.S. Environmental Protection Agency
401 M Street, SW
Washington, D.C. 20460-0001

Hand or courier deliveries of final printed labeling may be made to:

Document Processing Desk (FIN-LABEL)
Office of Pesticide Programs
Room 266A, Crystal Mall 2
1921 Jefferson Davis Highway
Arlington, VA 22202

Sincerely,



Jim Tompkins, Deputy Chief
Registration Support Branch
Registration Division (7505W)

Attachment

Glenda Haage
RIVERSIDE/TERRA CORP

COMMENTS FOR: EPA Reg. No. 9779-210
RIVERSIDE DIAZINON 4AG

The following specific comments pertain to your WPS labeling submission concerning the product cited above:

Move the statement "Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application." from its current position in the Agricultural Use Box to the "Directions for Use"

The following statement on your proposed label is not required for this product: "Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. You may delete it or leave it on the label."

Delete the crossed out statements on your proposed label. They are redundant statements or phrases.

Correct the typographical errors circled on your proposed label.

40933

R*

RESTRICTED USE PESTICIDE
DUE TO AVIAN AND AQUATIC TOXICITY
FOR RETAIL SALE TO AND USE BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT SUPERVISION AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION.

THIS PRODUCT MUST NOT BE USED ON GOLF COURSES AND SOD FARMS

DIAZINON 4AG

For control of certain insects on fruits, nuts, vegetables, field crops, lawns, and ornamentals

ACTIVE INGREDIENTS

| | |
|---|-------|
| Diazinon (O,O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate)* | 48.9% |
| INERT INGREDIENTS** | 51.1% |
| Total 100.0% | |

*Contains 4 pounds of diazinon per gallon.
 **Contains xylene range aromatic solvents.

EPA REG. NO. 9779-210

EPA EST. NO. 9779-

STOP - READ LABEL BEFORE USING!

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT
ORGANOPHOSPHATE

ACCEPTED
with COMMENTS
In EPA Letter Dated

MAR 24 1981

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

9779-210

IF SWALLOWED: Call a physician or Poison Control Center immediately. DO NOT INDUCE VOMITING. Vomiting may cause aspiration pneumonia. Do not give anything by mouth if person is unconscious or convulsing.

IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

IF ON SKIN: Wash promptly with soap and water. Rinse thoroughly. Get medical attention.

IF IN EYES: Rinse eyes with plenty of water. Call a physician if irritation persists.

NOTE TO PHYSICIAN: This product is an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine. This product contains a xylene range aromatic solvent which may present an aspiration hazard. Gastric lavage may be indicated if product was taken internally.

Manufactured For
RIVERSIDE/TERRA CORPORATION
 P.O. Box 8000, Sioux City, Iowa 51102
 Riverside Serves Agriculture. Agriculture Serves Everyone

Net Contents
____ GALS

5 of 33

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Harmful if swallowed, inhaled or absorbed through skin. Causes moderate eye injury. ~~Avoid contact with skin, eyes, or clothing. Avoid breathing vapor or spray mist. Wash thoroughly with soap and water after handling and before eating or smoking. Remove contaminated clothing and wash before reuse.~~ Avoid contamination of food and feed. Food utensils such as tablespoons and measuring cups should not be used for food purposes after use in measuring pesticides. Keep out of reach of domestic animals. Do not use on humans, household pets or livestock. Do not contaminate ornamental fish ponds. ~~Do not permit children or pets to go onto treated areas until spray has dried.~~

Resistant Protective Equipment

Some materials that are chemical-resistant to this product are listed below. If you want more options follow the instructions for category E on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear long-sleeved shirt and long pants, chemical-resistant gloves such as Barrier Laminate, Butyl Rubber, Nitrile Rubber, or Viton, shoes plus socks.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

This product is highly toxic to birds, fish and other wildlife. Birds, especially waterfowl, feeding or drinking on treated areas may be killed. Because of the migratory habits of Atlantic coast waterfowl, do not apply this product to lawns in Nassau County, New York, between November 1 and May 20. Do not exceed maximum permitted label rates. Rates above those recommended significantly increase potential hazards to birds, especially waterfowl. Avoid overlapping of sprays. If waterfowl; i.e., ducks or geese, can be expected in the treated area after treatment (except spot treatment), apply at least 1/4 inch of water immediately after spraying this product; however, stop watering before puddling occurs. Keep out of lakes, streams, ponds, tidal marshes and estuaries. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, shrimp, crab, and other aquatic life are important resources. Do not contaminate water by cleaning of equipment or disposal of equipment wash waters.

This pesticide is highly toxic to bees exposed to direct treatment or to residues on blooming crops or weeds. Do not apply this pesticide or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

Physical or Chemical Hazards

Do not use, pour, spill, or store near heat or open flame.

BEST AVAILABLE COPY

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR INSECT CONTROL, CROP INJURY OR ILLEGAL RESIDUES.

~~of accidental exposure, as described under Statement of Practical Treatment in the Precautionary Statements on this label. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Written warnings must include the following information: WARNING. Area treated with DIAZINON 4AG on (date of application). Do not enter without appropriate personal protective equipment for 24 hours after application. In case of accidental exposure, flush eyes or skin with plenty of water. Call physician if irritation persists. Remove and wash contaminated clothing before reuse.~~

Note: The California Department of Agriculture has set a 5-day reentry period for diazinon on grapes, peaches, and nectarines. Five days should elapse between the time of application and the time a worker may enter the field to engage in any activity requiring substantial contact with treated foliage. When a mixture of two or more organophosphate pesticides are applied in combination, the interval should be prolonged by adding to the largest applicable interval an additional 50% of that interval.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

Pesticide

Pesticide wastes are toxic, improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If pesticide, spray mixture or rinsate cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container

Triple rinse (or equivalent). Then puncture and dispose of in a sanitary landfill, or incinerate, or burn, if allowed by state and local authorities. Stay out of smoke from burning container.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup and disposal of wastes. In the event of a major spill, fire or other emergency, call 1-800-424-8802 (CHEM TREC) day or night.

GENERAL INFORMATION

DIAZINON 4AG is an emulsifiable solution which, when diluted with water according to the directions, is intended for the control of insect pests of agricultural crops and other plants. Do not use on food crops grown in greenhouses. Do not use in barns or dwellings. Do not use on livestock.

In the directions for use which follow, best control is obtained when application is made at first sign of pests. Application should be repeated only as directed to maintain control.

To avoid spray drift, do not apply under windy conditions. Avoid spray overlap, since crop or plant injury may result.

Work Safety Rules

REPEATED EXPOSURES TO CHOLINESTERASE INHIBITORS SUCH AS ARE CONTAINED IN THIS PRODUCT MAY, WITHOUT WARNING, CAUSE PROLONGED SUSCEPTIBILITY TO VERY SMALL DOSES OF ANY CHOLINESTERASE INHIBITOR.

When handling DIAZINON 4AG, do not rub eyes or mouth with hands. If you feel sick in any way, STOP work and get help right away. See Practical Treatment section of this label. ~~When handling this product, wear the personal protective equipment listed in the following section.~~

Personal Protective Equipment

~~Long sleeved shirt, long legged pants, shoes and socks, and chemical resistant gloves.~~

~~Do not apply this product in a way that will contact unprotected workers, either directly or through drift. Only protected handlers may be in the area during application.~~

BEST AVAILABLE COPY

CHEMIGATION
APPLICATION THROUGH AN IRRIGATION SYSTEM (CHEMIGATION) - CRANBERRIES ONLY

DIAZINON 4AG may be applied alone or in combination with other pesticides registered for application through sprinkler irrigation systems to cranberries. To insure compatibility, pour the products into a small container of water in the correct proportions. After thorough mixing, let stand for five minutes. If the combination remains mixed, or can be remixed readily, the mixture is compatible.

Do not apply this product through any type of irrigation system to other crops. For chemigation, apply this product to cranberries only through sprinkler irrigation systems. Do not apply this product to cranberries through any other type of irrigation system.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

Do not connect the irrigation system used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments, if the need arises.

Chemigation System Connected to Public Water Systems

1. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
2. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.

Sprinkler Chemigation

1. The system must contain a functional check valve, a vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
2. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
3. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
5. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended for treatment.

Operating Instructions

1. Determine the acreage to be treated by the sprinkler system.
2. Measure the appropriate amount of DIAZINON 4AG needed to treat the acreage to be treated. Refer to the specific directions for use for cranberries for the application rate and the amount of water per acre.

BEST AVAILABLE COPY

3. Add the premeasured DIAZINON 4AG to the chemigation system supply tank and dilute with water if needed. Proper agitation is needed if DIAZINON 4AG is tank mixed with other pesticides registered for sprinkler chemigation application to cranberries.
4. Start the irrigation system and bring the system up to full pressure, making certain that all sprinkler heads are functioning properly.
5. Engage the chemigation injection or venturi system to add the diazinon to the irrigation lines.
6. Continue to run irrigation system for a few minutes after supply tank is empty to allow material in the irrigation lines to reach the sprinkler heads furthest from the chemigation injection point.

FRUIT AND NUT CUPS

DIAZINON 4AG may be applied using ground or aerial application equipment as specified in the following table. This table indicates the *minimum* amount of water that can be used for the various methods of application. The rate of DIAZINON 4AG for aerial spray or concentrated spray applied to fruit trees by ground equipment is based on a dilute full cover spray applied with conventional ground equipment at a rate of 300-400 gals. of water per acre. The specific amount of product to be used appears under the separate directions for use for each crop. If the Rate calls for 1 pt. of DIAZINON 4AG in 100 gals. of water, this amounts to 3-4 pts. of product per acre when applied at the usual dilute ground spray volumes of 300-400 gals. of water per acre. In no case should the amount of product used per acre exceed the maximum amount per acre per application that is specified in the Timing/Special Directions for individual crops.

| CROP | AERIAL APPLICATION MINIMUM GALLONS/ACRE | GROUND APPLICATION MINIMUM GALLONS/ACRE | |
|-----------------|--|--|-------------|
| | | DILUTE | CONCENTRATE |
| Almonds* | 20 | 100 | 20 |
| Apricots | 20 | 100 | 20 |
| Blackberries** | 20 | 100 | 20 |
| Boysenberries** | 20 | 100 | 20 |
| Cherries | 20 | 100 | 20 |
| Cranberries | 20 | 15 | -- |
| Dewberries** | 20 | 100 | 20 |
| Grapes | 20 | 100 | 20 |
| Loganberries** | 20 | 100 | 20 |
| Nectarines | 20 | 100 | 20 |
| Peaches | 20 | 100 | 20 |
| Plums | 20 | 100 | 20 |
| Prunes | 20 | 100 | 20 |
| Raspberries** | 20 | 100 | 20 |
| Strawberries | 20 | 100 | 20 |
| Walnuts* | 20 | 100 | 20 |

* CA only.

** CA, OR, or WA only.

Observe stated time intervals between last application and harvest, as well as intervals between applications and total amount of product that can be applied per season.

| CROP | PHI* |
|---------------|------|
| Almonds | ** |
| Apricots | 21 |
| Blackberries | 7 |
| Boysenberries | 7 |
| Cherries | 21 |
| Cranberries | 7 |
| Dewberries | 7 |
| Grapes | 28 |
| Loganberries | 7 |
| Nectarines | 21 |
| Peaches | 21 |
| Plums, Prunes | 21 |
| Raspberries | 7 |
| Strawberries | 5 |
| Walnuts | 45 |

* PHI - Pre-Harvest Interval or the number of days between the last application and harvest; may vary based on use pattern selected. See specific directions for use on grapes.

** Dormant spray only.

Note: To protect bees, do not apply this product to fruit trees when trees or substantial numbers of weeds in the orchard are in bloom.

| CROP | PEST | RATE | TIMING/SPECIAL DIRECTIONS |
|-------------------|---|---|---|
| Almonds (CA only) | San Jose Scale, Partstoria Scale | 1-1½ pts. + 2-3 gals. dormant oil or 1-1½ gals. superior type oil in 100 gals. of water | Apply as a dormant spray. Do not apply more than 6 pts. of DIAZINON 4AG or 9 gals. of oil per acre. |
| Apricots | Aphids, Brown Mites, (Clover Mites), Two-spotted Spider Mites, Olive Scale Crawlers, San Jose Scale, Crawlers | 1 pt. in 100 gals. of water | Apply as infestations occur. Repeat application after 14 days if necessary. |
| | Olive Scale Crawlers | ½ pt. + 1½ gals. of light medium horticultural oil per 100 gals. of water | Apply when scale crawlers are present. |

| CROP | PEST | RATE | TIMING/SPECIAL DIRECTIONS |
|--------------------------------|-------------------|---------------------------------|--|
| Apricots <i>(continued)</i> | Apricot Mealybugs | 1 pt. per 100 gals. of water | Apply as a cover spray from petal fall to June. Allow 14 days between applications. |

Notes: (1) Do not apply more than 4 pts. of product per acre per application and no more than 12 pts. of product per acre per season. (2) To avoid illegal residues, allow a minimum of 21 days between the last application and harvest.

| | | | |
|--|---|-------------------------------|--|
| Caneberries (CA, OR and WA only) (Blackberries, Boysenberries, Dewberries, Loganberries, Raspberries) | Leafhoppers, Thrips, Aphids, Two-spotted Spider Mites, Raspberry Fruitworms, Raspberry Sawflies | 1 qt. per acre | Apply as insects occur. Repeat application after 14 days if necessary. Do not make more than 5 applications per season. Do not apply within 7 days of harvest or illegal residues may occur. Apply in a minimum of 100 gals. of water per acre (200 gals. per acre maximum). For Raspberry Fruitworms make one application when blossom buds separate. Make a second application immediately before blossoms open. |
| | Dryberry Mites | 1-2 qts. per acre | Apply as a dormant spray in the early Spring when canes are still on the ground and buds are closed. Make a second application when buds are well developed but still closed. Apply in a minimum of 100 gals. of water per acre. Do not apply within 7 days of harvest or illegal residues may occur. |
| | Raspberry Crown Borers (Raspberry Root Borers) | 2 qts. per acre | Apply a single application in a minimum of 100 gals. of water per acre as a drench to the crown and lower canes in the Spring before buds break. |
| Cherries | Eyespotted Bud Moths, Fruit Tree Leafrollers | 1 pts. in 100 gals. | Apply in cover sprays as infestations occur beginning at petal fall. Repeat application after 7 days if necessary. |
| | Leafhoppers | 1 pt. in 100 gals. of water | Make 3-4 applications beginning at petal fall and continuing through the growing season as infestations occur. Allow 7 days between applications. |
| | Cherry Fruitflies | ½-1 pt. in 100 gals. of water | Make 3-4 applications beginning when adult flies begin to emerge and repeating at 10-day intervals. |
| | Black Cherry Aphids | 1 pt. in 100 gals. of water | Apply as a cover spray when aphids appear. Repeat application after 10 days if necessary. |

| CROP | PEST | RATE | TIMING/SPECIAL DIRECTIONS |
|--------------------------------|----------------------------|----------------------------------|---|
| Cherries <i>(continued)</i> | San Jose Scale Crawlers | 1 pt. in 100 gals. of water | Apply during the pre-pink stage or when crawlers first appear. |
| | Cherry Rust Mites | ½-1 pt. in 100 gals. of water | Apply to trees as a single supplemental spray after harvest. |

Notes: (1) Do not make more than 3 applications per season. (2) Do not apply more than 4 pts. of product per acre per application and no more than 12 pts. of product per acre per season. (3) Allow a minimum of 21 days between the last application and harvest.

| | | | |
|-------------|---|---|---|
| Cranberries | Blackheaded Fireworms <i>(Rhopobota naevana)</i> | 2 qts. in a min. of 15 gals. of water per acre for ground application and up to 400 gals. of water per acre for chemigation | Apply when pest appears. Repeat application after 14 days if necessary. A maximum of 6 applications per season is permitted. Do not apply within 7 days of harvest or illegal residues may occur. |
| | Cranberry Fruitworms | 3 qts. in a min. of 15 gals. of water per acre for ground application and up to 400 gals. of water per acre for chemigation | Apply when pest appears. Repeat application after 14 days if necessary. A maximum of 4 applications per season is permitted. Do not apply within 7 days of harvest or illegal residues may occur. |

Note: This product may be applied to cranberries through sprinkler irrigation. Refer to the Chemigation section for specific chemigation directions for use.

| | | | |
|--------|--|--|--|
| Grapes | Pacific Spider Mites, Leafhoppers, Grape Berry Moths, Grape Leafrollers, Omnivorous Leafrollers, Aphids, Spider Mites | 1-2 pts. per acre in a minimum of 100 gals. of water | Apply as pest occurs as a thorough cover spray. Repeat application after 7 days if necessary. |
| | <i>Drosophila spp.</i> | 6 fl. oz. per acre in a minimum of 100 gals. of water (300 gals. of water maximum) | Apply as pest occurs as a thorough cover spray. Repeat application after 7 days if necessary. |

Notes: (1) Do not apply more than 2 pts. of product per acre per application and no more than 10 pts. per acre per season. (2) To avoid illegal residues, allow 28 days between the last application and harvest.

| CROP | PEST | RATE | TIMING/SPECIAL DIRECTIONS |
|------------|--|-----------------------------|--|
| Nectarines | Oriental Fruit Moths | 1 pt. in 100 gals. of water | Apply at 100% petal fall. Make 2 additional applications at 10-15 day intervals. Time the last application to peak adult flight. |
| | Aphids, Brown Mites (Clover Mites), Two-spotted Spider Mites, Olive Scale Crawlers, San Jose Scale Crawlers, Peach Twig Borers | 1 pt. in 100 gals. of water | Apply as infestations occur. Repeat application after 14 days if necessary. |

Notes: (1) Do not apply more than 4 pts. of product per acre per application and no more than 12 pts. per acre per season. (2) Allow a minimum of 21 days between the last application and harvest.

| | | | |
|---------|---|---------------------------------|--|
| Peaches | Aphids, Brown Mites (Clover Mites), Two-spotted Spider Mites, Olive Scale Crawlers, San Jose Scale Crawlers | 1 pt. in 100 gals. of water | Apply as infestations occur. Repeat application after 14 days if necessary. |
| | Oriental Fruit Moths | 1 pt. in 100 gals. of water | Apply at 100% petal fall. Make 2 additional applications at 10-15 day intervals. Time the last application to peak adult flight. |
| | Peach Twig Borers | 1 pt. in 100 gals. of water | Apply as insects occur in May and June. Allow 7 days between applications. |
| | Leafhoppers | 1 pt. in 100 gals. of water | Make 3-4 applications beginning at petal fall and continuing through the growing season. Allow at least 7 days between applications. |
| | White Peach Scale | 1½-2 pts. in 100 gals. of water | Apply post-harvest timed to coincide with peak crawler and immature scale activity. Under conditions of heavy infestation, 2 applications (7 days apart) may be needed. Do not apply before fruit is harvested. |

Notes: (1) Do not apply more than 4 pts. of product per acre per application and no more than 12 pts. of product per acre per season. (2) Allow a minimum of 21 days between the last application and harvest.

| | | | |
|-------|---|-------------------------------|-------------------------------|
| Plums | Brown Mites (Clover Mites), European Red Mites | 1 pt. in 100 gals. of water | Apply as needed every 7 days. |
| | Leafcurl Plum Aphids, Mealy Plum Aphids, Thistle Aphids | ½-1 pt. in 100 gals. of water | |

Notes: (1) Do not apply more than 4 pts. of product per acre per application and no more than 12 pts. of product per acre per season. (2) Allow a minimum of 21 days between the last application and harvest.

| CROP | PEST | RATE | TIMING/SPECIAL DIRECTIONS |
|--------|---|-------------------------------|------------------------------|
| Prunes | Brown Mites (Clover Mites), European Red Mites | 1 pt. in 100 gals. of water | Apply as needed every 7 days |
| | Leafcurl Plum Aphids, Mealy Plum Aphids, Thistle Aphids | ½-1 pt. in 100 gals. of water | |

Notes: (1) Do not apply more than 4 pts. of product per acre per application and no more than 12 pts. of product per acre per season. (2) Allow a minimum of 21 days between the last application and harvest.

| | | | |
|-------------------|--|--------------------------------------|--|
| Strawberries | Aphids, Two-spotted Spider Mites | 1 pt. in 100 gals. of water per acre | Apply as pests occur. Repeat application if necessary every 7 days, but do not make more than 3 applications per season. Do not apply within 5 days of harvest or illegal residues may occur. |
| | Cyclamen Mites | 1 qt. in 100 gals. of water per acre | Direct spray to the plant crowns. Plants should be sufficiently agitated to assure thorough coverage of the foliage. Repeat application after 7 days if necessary. Do not make more than 3 foliar applications per season. Do not apply within 5 days of harvest or illegal residues may occur. Application may be made to plants after harvest. |
| | Mole Crickets | 1 qt. per acre | Broadcast in sufficient water to obtain even coverage when insects are present. Apply 1-2 days before transplanting and immediately incorporate into the top 1-2 inches of soil. |
| | Strawberry Leafrollers | ¾-1 pt. in 100 gals. of water | Apply 200 gals. of spray per acre when blossoms show color. Do not apply within 5 days of harvest or illegal residues may occur. |
| Walnuts (CA only) | Aphids, Spider Mites | 1 qt. per acre | Apply as a foliar spray. Repeat application in 14 days if necessary. Do not make more than 3 applications per season. Do not apply within 45 days of harvest or after husks open or illegal residues may occur. Make ground applications in a minimum of 100 gals. of water per acre. |
| | Codling Moths, Scale Crawlers, Walnut Caterpillars | 3 qts. per acre | |

VEGETABLE CROPS

For foliar or soil applications, DIAZINON 4AG should always be applied as a spray in sufficient water to assure thorough coverage of the foliage or soil. DIAZINON 4AG may be applied using ground or aerial application equipment as specified in the following table. This table indicates the amount of water that can be used for the various methods of application. The specific amount of product to be used appears under the separate directions for use for each crop.

| CROP | GROUND APPLICATION | | AERIAL APPLICATION MINIMUM GALS/ACRE |
|--|----------------------|-------------------------------------|---|
| | MINIMUM GALS/ACRE | OR MINIMUM GALS/1,000 SQ. FT. | |
| Beans (Lima, Pole, and Snap) (succulent only) | 10 | 2 | 5 |
| Beets, Red (Table) | 10 | 2 | 5 |
| Broccoli | 10 | 2 | 5 |
| Brussels Sprouts | 10 | 2 | 5 |
| Cabbage | 10 | 2 | 5 |
| Cantaloupes | 5 | 1 | 5 |
| Carrots | 10 | 2 | 5 |
| Casabas | 5 | 1 | 5 |
| Cauliflower | 10 | 2 | 5 |
| Chinese Broccoli | 10 | 2 | 5 |
| Chinese Cabbage | 10 | 2 | 5 |
| Chinese Mustard | 10 | 2 | 5 |
| Chinese Radish (CA, FL only) | 10 | 2 | 5 |
| Collards | 10 | 2 | 5 |
| Crenshaws | 5 | 1 | 5 |
| Cucumbers | 5 | 1 | 5 |
| Endive (Escarole) | 10 | 2 | 5 |
| Ginseng | 10 | 2 | 5 |
| Honeydew Melons | 5 | 1 | 5 |
| Kale | 10 | 2 | 5 |
| Lettuce, Head | 10 | 2 | 5 |
| Lettuce, Leaf | 10 | 2 | 5 |
| Muskmelons | 5 | 1 | 5 |
| Mustard | 10 | 2 | 5 |
| Onions (Bulb and Green) | 10 | 2 | 5 |
| Parsley | 10 | 2 | 5 |
| Parsnips | 10 | 2 | 5 |
| Peas (succulent only) | 10 | 2 | 5 |

| CROP | GROUND APPLICATION | | AERIAL APPLICATION MINIMUM GALS/ACRE |
|-------------------|----------------------|-------------------------------------|---|
| | MINIMUM GALS/ACRE | OR MINIMUM GALS/1,000 SQ. FT. | |
| Peppers | 10 | 2 | 5 |
| Persian Melons | 5 | 1 | 5 |
| *Potatoes (Irish) | 10 | 2 | 5 |
| Radishes | 10 | 2 | 5 |
| Spinach | 10 | 2 | 5 |
| Squash, Summer | 5 | 1 | 5 |
| Squash, Winter | 5 | 1 | 5 |
| Sweet Corn | 20 | 4 | 5 |
| Sweet Potatoes | 10 | 2 | 5 |
| Swiss Chard | 10 | 2 | 5 |
| Tomatoes | 10 | 2 | 5 |
| Turnips | 10 | 2 | 5 |
| Watermelons | 5 | 1 | 5 |

* Do not use on commercially grown potatoes which will be hand-harvested.

Observe stated time intervals between last application and harvest as well as intervals between and maximum numbers of applications per season. Do not apply to food crops grown in greenhouses.

Pre-Harvest Intervals (PHI*) for Foliar Applications

| CROP | PHI* (DAYS) |
|------------------------------|-------------|
| Beans (Lima, Pole, and Snap) | 7 |
| Beets, Red (Table) | 14 |
| Broccoli | 7 |
| Brussels Sprouts | 7 |
| Cabbage | 21 |
| Cantaloupes | 3 |
| Carrots | 14 |
| Casabas | 3 |
| Cauliflower | 7 |
| Chinese Broccoli | 10 |
| Chinese Cabbage | 10 |
| Chinese Mustard | 10 |

| CROP | PHI* (DAYS) |
|-------------------------|-------------|
| Chinese Radish | 10 |
| Collards | 10 |
| Crenshaws | 3 |
| Cucumbers | 7 |
| Endive (Escarole) | 14 |
| Ginseng | 30 |
| Honeydew Melons | 3 |
| Kale | 10 |
| Lettuce | 14 |
| Muskmelons | 3 |
| Mustard | 10 |
| Onions (Bulb and Green) | 14 |
| Parsnips | 14 |
| Peas | 7 |
| Peppers | 5 |
| Persian Melons | 3 |
| Potatoes (Irish) | 35 |
| Radishes | 14 |
| Spinach | 14 |
| Squash, Summer | 7 |
| Squash, Winter | 3 |
| Sweet Corn | 7 |
| Swiss Chard | 14 |
| Tomatoes | 1 |
| Turnips, Roots and Tops | 14 |
| Watermelons | 3 |

*PHI - Pre-Harvest Interval or the number of days between the last application and harvest.

Notes: (1) Soil Incorporation: Following application of this product to control vegetable soil insects, immediately incorporate the product into the soil to the recommended depth (see Timing/Special Directions) using a rotary hoe, cultivator, disk, harrow, or other suitable means. For control of surface cutworms, incorporate 2-3 inches, for subterranean cutworms incorporate 3-6 inches. (2) To Protect Bees: Foliar applications of this product should not be made to peas or beans if these crops or weeds in the treatment area are in bloom, or to corn during the pollen shed period.

| CROP | PEST | RATE | TIMING/SPECIAL DIRECTIONS |
|--|---|--------------------|---|
| Beans (Lima, Pole, and Snap) (succulent only) | Cutworms | 2-4 qts. per acre | Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.) |
| | Wireworms | 3-4 qts. per acre | Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil. |
| | Black Bean Aphids, Mexican Bean Beetles, Dipterous Leafminers | 1-1½ pts. per acre | Apply as insects occur. Repeat application, as necessary, every 5 days. Do not make more than 3 applications per season. Do not apply within 7 days of harvest or illegal residues may occur. Bean forage or bean hay may be fed to dairy and beef cattle and sheep. Wait 4 days following treatment if forage is to be cut for hay. |
| | Cucumber Beetles | ¾ pt. per acre | Make cucumber beetle application in 100 gals. of water per acre. |
| Beans, Lima (succulent only) | Leafhoppers, Spider Mites | 1 pt. per acre | Apply as pests occur. Repeat application, as necessary, every 5 days. Do not make more than 3 applications per season. Notes: (1) DIRZINON 4-AG will not control organophosphate-resistant leafminers, (2) To protect bees, do not apply foliarty if the crop or weeds in the treatment area are in bloom. Do not apply within 7 days of harvest or illegal residues may occur. Bean forage or bean hay may be fed to dairy and beef cattle and sheep. Wait 4 days following treatment if forage is to be cut for hay. Note: To protect bees, do not apply foliarty if the crop or weeds in the treatment area are in bloom. |
| Beets, Red (Table) | Cutworms | 2-4 qts. per acre | Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.) |
| | Mole Crickets | 1 qt. per acre | Apply to plant beds or fields 1-2 days before sowing of seed or transplanting and immediately incorporate into the top 1-2 inches of soil. |
| | Wireworms | 3-4 qts. per acre | Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil. |

| CROP | PEST | RATE | TIMING/SPECIAL DIRECTIONS |
|--|----------------------|-------------------------------|--|
| Beets, Red (Table) <i>(continued)</i> | Dipterous Leafminers | ½-1 pt. per acre | <p>Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season.</p> <p>Do not apply within 14 days of harvest, or illegal residues may occur.</p> <p>Note: DIAZINON 4AG will not control organo-phosphate-resistant leafminers.</p> |
| Broccoli | Root Maggots | 2-3 qts. per acre | <p>Apply to plant beds or fields just before sowing of seed or transplanting and immediately incorporate into the top 3-4 inches of soil.</p> |
| | | 4-8 oz. per 50 gals. of water | <p>In transplant water as a drench application when 200-300 gals. of water are used per acre. Apply ½-1 cup (4 to 8 oz.) per plant by hand or by tractor mounted sprayer equipped with drop nozzles to direct spray to the base of the plant.</p> <p>Note: Transplant water treatments may result in stand reduction due to plant stress at time of transplanting.</p> |
| | Cutworms | 2-4 qts. per acre | <p>Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)</p> |
| | Mole Crickets | 1 qt. per acre | <p>Apply to plant beds or fields 1-2 days before sowing of seed or transplanting and immediately incorporate into the top 1-2 inches of soil.</p> |
| | Wireworms | 3-4 qts. per acre | <p>Broadcast just before planting and immediately incorporate into the top 1-2 inches of soil.</p> |
| Brussels Sprouts | Root Maggots | 2-3 qts. per acre | <p>Apply to plant beds or fields just before sowing of seed or transplanting and immediately incorporate into the top 3-4 inches of soil.</p> |

| CROP | PEST | RATE | TIMING/SPECIAL DIRECTIONS |
|---------------------------------|--|------------------------------|---|
| Brussels Sprouts (continued) | | 4-8 oz. in 50 gals. of water | In transplant water as a drench application when 200-300 gals. of water are used per acre. Apply ½-1 cup (4 to 8 oz.) per plant by hand or by tractor-mounted sprayer equipped with drop nozzles to direct spray to the base of the plant. Note: Transplant water treatments may result in stand reduction due to plant stress at time of transplanting. |
| | Cutworms | 2-4 qts. per acre | Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.) |
| | Mole Crickets | 1 qt. per acre | Apply to plant beds or fields 1-2 days before sowing of seed or transplanting and immediately incorporate into the top 1-2 inches of soil. |
| | Wireworms | 3-4 qts. per acre | Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil. |
| | Aphids, Diamondback Moths, Imported Cabbageworms | ½-1 pt. per acre | Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 7 days of harvest or illegal residues may occur. |
| Cabbage | Root Maggots | 2-3 qts. per acre | Apply to plant beds or fields just before sowing of seed or transplanting and immediately incorporate into the top 3-4 inches of soil. |
| | | 4-8 oz. in 50 gals. of water | In transplant water as a drench application when 200-300 gals. of water are used per acre. Apply ½-1 cup (4 to 8 oz.) per plant by hand or by tractor-mounted sprayer equipped with drop nozzles to direct spray to the base of the plant. Note: Transplant water treatments may result in stand reduction due to plant stress at time of transplanting. |
| | Cutworms | 2-4 qts. per acre | Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.) |
| | Mole Crickets | 1 qt. per acre | Apply to plant beds or fields 1-2 days before sowing of seed or transplanting and immediately incorporate into the top 1-2 inches of soil. |
| | Wireworms | 3-4 qts. per acre | Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil. |

2/9/33

| CROP | PEST | RATE | TIMING/SPECIAL DIRECTIONS |
|-------------------------------|--|------------------------------|--|
| Cabbage <i>(continued)</i> | Aphids, Diamondback Moths, Imported Cabbageworms, Dipterous Leafminers | ½-1 pt. per acre | <p>Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season.</p> <p>Do not apply within 21 days of harvest or illegal residues may occur.</p> <p>Note: DIAZINON 4AG will not control organo-phosphate-resistant leafminers.</p> |
| Carrots | Cutworms | 2-4 qts. per acre | Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.) |
| | Mole Crickets | 1 qt. per acre | Apply to plant beds or fields 1-2 days before sowing of seed or transplanting and immediately incorporate into the top 1-2 inches of soil. |
| | Wireworms | 3-4 qts. per acre | Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil. |
| | Aphids | 1 pt. per acre | <p>Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season.</p> <p>Do not apply within 14 days of harvest or illegal residues may occur.</p> |
| Cauliflower | Root Maggots | 2-3 qts. per acre | Apply to plant beds or fields just before sowing of seed or transplanting and immediately incorporate into the top 3-4 inches of soil. |
| | | 4-8 oz. in 50 gals. of water | <p>In transplant water as a drench application when 200-300 gals. of water are used per acre. Apply ½-1 cup (4 to 8 oz.) per plant by hand or by tractor-mounted sprayer equipped with drop nozzles to direct spray to the base of the plant.</p> <p>Note: Transplant water treatments may result in stand reduction due to plant stress at time of transplanting.</p> |
| | Cutworms | 2-4 qts. per acre | Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.) |
| | Mole Crickets | 1 qt. per acre | Apply to plant beds or fields 1-2 days before sowing of seed or transplanting and immediately incorporate into the top 1-2 inches of soil. |
| | Wireworms | 3-4 qts. per acre | Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil. |

220/53

| CROP | PEST | RATE | TIMING/SPECIAL DIRECTIONS |
|--|--|------------------|--|
| Cauliflower <i>(continued)</i> | Aphids, Diamondback Moths, Imported Cabbageworms, Dipterous Leafminers | ½-1 pt. per acre | <p>Apply as insects occur. Repeat application, as necessary, up to 5 times, with a minimum of 7 days between applications.</p> <p>Do not apply within 5 days of harvest or illegal residues may occur.</p> <p>Note: DIAZINON 4AG will not control organo-phosphate-resistant leafminers.</p> |
| Chinese Broccoli | Aphids, Diamondback Moths, Imported Cabbageworms, Dipterous Leafminers | ½-1 pt. per acre | <p>Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season.</p> <p>Do not apply within 10 days of harvest or illegal residues may occur.</p> <p>Note: DIAZINON 4AG will not control organo-phosphate-resistant leafminers.</p> |
| Chinese Cabbage (Bok Choy and Napa) | Aphids, Diamondback Moths, Imported Cabbageworms, Dipterous Leafminers | ½-1 pt. per acre | <p>Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season.</p> <p>Do not apply within 10 days of harvest or illegal residues may occur.</p> <p>Note: DIAZINON 4AG will not control organo-phosphate-resistant leafminers.</p> |
| Chinese Mustard (Gai Choy) | Aphids, Diamondback Moths, Imported Cabbageworms, Dipterous Leafminers | ½-1 pt. per acre | <p>Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season.</p> <p>Do not apply within 10 days of harvest or illegal residues may occur.</p> <p>Note: DIAZINON 4AG will not control organo-phosphate-resistant leafminers.</p> |
| Chinese Radish (Japanese Radish or Daikon) (FL and CA only) | Aphids, Diamondback Moths, Imported Cabbageworms, Dipterous Leafminers | 1 pt. per acre | <p>Mix in a minimum of 100 gals. of water per acre and apply as a foliar spray as pests occur. Repeat application as necessary, every 14 days. Do not make more than 3 applications per season.</p> <p>Do not apply within 10 days of harvest or illegal residues may occur.</p> <p>Note: DIAZINON 4AG will not control organo-phosphate-resistant leafminers.</p> |

| CROP | PEST | RATE | TIMING/SPECIAL DIRECTIONS |
|-------------------|---|-------------------|---|
| Collards | Cutworms | 2-4 qts. per acre | Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.) |
| | Mole Crickets | 1 qt. per acre | Apply to plant beds or fields 1-2 days before sowing of seed or transplanting and immediately incorporate into the top 1-2 inches of soil. |
| | Wireworms | 3-4 qts. per acre | Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil. |
| | Aphids, Diamondback Moths, Imported Cabbageworms, Harlequin Bugs, Corn Earworms | ½-1 pt. per acre | Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 10 days of harvest or illegal residues may occur. |
| Cucumbers | Cutworms | 2-4 qts. per acre | Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.) |
| | Wireworms | 3-4 qts. per acre | Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil. |
| | Aphids, Serpentine Leafminers, Thrips, Striped Cucumber Beetles, Melonworms | 1 pt. per acre | Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 7 days of harvest or illegal residues may occur. Note: DIAZINON 4AG will not control organo-phosphate-resistant leafminers. |
| Endive (Escarole) | Cutworms | 2-4 qts. per acre | Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.) |
| | Mole Crickets | 1 qt. per acre | Apply to plant beds or fields 1-2 days before sowing of seed or transplanting and immediately incorporate into the top 1-2 inches of soil. |
| | Wireworms | 3-4 qts. per acre | Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil. |
| | Aphids, Dipterous Leafminers | ½-1 pt. per acre | Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 14 days of harvest or illegal residues may occur. Note: DIAZINON 4AG will not control organo-phosphate-resistant leafminers. |

| CROP | PEST | RATE | TIMING/SPECIAL DIRECTIONS |
|---------|---|-------------------|--|
| Ginseng | Leafhoppers, Aphids, Lygus Bugs, Flea Beetles, Jumping Plant Lice | ½ 1 pt. per acre | Spray when insects first appear. For heavy insect infestations, use 1 pt. per acre. Apply by ground equipment using 10 to 100 gals. of water per acre. Do not apply more than one application per growing season. Do not apply during flowering of three or four year old crops. Do not apply within 30 days of harvest. Do not graze treated areas or feed treated forage to livestock. |
| Kale | Cutworms | 2-4 qts. per acre | Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.) |
| | Mole Crickets | 1 qt. per acre | Apply to plant beds or fields 1-2 days before sowing of seed or transplanting and immediately incorporate into the top 1-2 inches of soil. |
| | Wireworms | 3-4 qts. per acre | Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil. |
| | Aphids, Diamondback Moths, Imported Cabbageworms, Harlequin Bugs, Corn Earworms | ½-1 pt. per acre | Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 14 days of harvest or illegal residues may occur. |
| Lettuce | Cutworms | 2-4 qts. per acre | Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.) |
| | Mole Crickets | 1 qt. per acre | Apply to plant beds or fields 1-2 days before sowing of seed or transplanting and immediately incorporate into the top 1-2 inches of soil. |
| | Wireworms | 3-4 qts. per acre | Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil. |
| | Aphids, Dipterous Leafminers | ½-1 pt. per acre | Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 14 days of harvest. Note: DIAZINON 4AG will not control organo-phosphate-resistant leafminers. |

| CROP | PEST | RATE | TIMING/SPECIAL DIRECTIONS |
|--|--|--------------------|---|
| Melons (Cantaloupes, Casabas, Crenshaws, Honeydews, Muskmelons, Persians, and Hybrids of these) and Watermelons | Cutworms | 2-4 qts. per acre | Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.) |
| | Wireworms | 3-4 qts. per acre | Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil. |
| | Aphids, Dipterous Leafminers, Thrips, Striped Cucumber Beetles, Melonworms, Leafhoppers, Spider Mites | ½-1½ pts. per acre | Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 3 days of harvest or illegal residues may occur. Note: DIAZINON 4AG will not control organo-phosphate-resistant leafminers. |
| Mustard | Cutworms | 2-4 qts. per acre | Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.) |
| | Mole Crickets | 1 qt. per acre | Apply to plant beds or fields 1-2 days before sowing of seed or transplanting and immediately incorporate into the top 1-2 inches of soil. |
| | Wireworms | 3-4 qts. per acre | Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil. |
| | Aphids, Dipterous Leafminers, Diamondback Moths, Imported Cabbageworms | ½-1 pt. per acre | Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 10 days of harvest or illegal residues may occur. Note: DIAZINON 4AG will not control organo-phosphate-resistant leafminers. |
| Onions (Bull and Green) | Onion Maggots | 2-4 qts. per acre | Broadcast just before planting and mix into the top 3-4 inches of soil Note: DIAZINON 4A will not control organo-phosphate-resistant leafminers. |
| | Onion Thrips | 1 pts. per acre. | Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 3 applications per season. Do not apply within 14 days of harvest or illegal residues may occur. |
| | Wireworms | 3-4 qts. per acre | Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil. |

| CROP | PEST | RATE | TIMING/SPECIAL DIRECTIONS |
|-----------------------|--|-------------------|---|
| Parsley | Cutworms | 2-4 qts. per acre | Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note). |
| | Wireworms | 3-4 qts. per acre | Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil. |
| Parsnips | Aphids, Flea Beetles, Dipterous Leafminers | ½-1 pt. per acre | <p>Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season.</p> <p>Do not apply within 14 days of harvest or illegal residues may occur.</p> <p>Note: DIAZINON 4AG will not control organo-phosphate-resistant leafminers.</p> |
| Peas (succulent only) | Cutworms | 2-4 qts. per acre | Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note). |
| | Wireworms | 3-4 qts. per acre | Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil. |
| | Aphids, Dipterous Leafminers | ½-1 pt. per acre | <p>Apply as insects occur. Repeat application, as necessary, every 5 days. Do not make more than 3 applications per season.</p> <p>To avoid illegal residues, allow a minimum of 7 days between the last application and harvest.</p> <p>Vines and hay may be fed to dairy and beef cattle and sheep. To avoid illegal residues, wait 7 days following treatment if forage is to be cut for hay.</p> <p>Notes: (1) DIAZINON 4AG will not control organo-phosphate-resistant leafminers. (2) To protect bees, do not apply foliarty if the crop or weeds in the treatment area are in bloom.</p> |
| Peppers | Cutworms | 2-4 qts. per acre | Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note). |
| | Mole Crickets | 1 qt. per acre | Apply to plant beds or fields 1-2 days before sowing of seed or transplanting and immediately incorporate into the top 1-2 inches of soil. |
| | Wireworms | 3-4 qts. per acre | Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil. |

| CROP | PEST | RATE | TIMING/SPECIAL DIRECTIONS |
|---|---|-------------------|--|
| Peppers <i>(continued)</i> | Aphids, Serpentine Leafminers | ½-1 pt. per acre | <p>Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season.</p> <p>Do not apply within 5 days of harvest or illegal residues may occur.</p> <p>Note: DIAZINON 4AG will not control organo-phosphate-resistant leafminers.</p> |
| Potatoes | Cutworms (Surface and Subterranean) | 2-4 qts. per acre | Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note). |
| | Wireworms | 3-4 qts. per acre | Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil. |
| | Southern Potato Wireworms | 2-4 qts. per acre | For potatoes grown on mineral soils in the Southeast, apply before planting and immediately incorporate into the top 3-6 inches of soil depending on location of wireworms. |
| | Aphids, Flea Beetles, Colorado Potato Beetles, Dipterous Leafminers | ½-¾ pt. per acre | <p>Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season.</p> <p>Do not apply within 35 days of harvest or illegal residues may occur.</p> <p>Note: DIAZINON 4AG will not control organo-phosphate-resistant leafminers.</p> |
| | Banded Cucumber Beetles, Leafhoppers, Southern Armyworms | ¾-1 pt. per acre | <p>Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season.</p> <p>Do not apply within 35 days of harvest or illegal residues may occur.</p> |
| Note: Do not use on commercially grown potatoes which will be hand-harvested. | | | |
| Radishes | Cutworms | 2-4 qts. per acre | Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note). |
| | Mole Crickets | 1 qt. per acre | Apply to plant beds or fields 1-2 days before sowing of seed or transplanting and immediately incorporate into the top 1-2 inches of soil. |
| | Wireworms | 3-4 qts. per acre | Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil. |

| CROP | PEST | RATE | TIMING/SPECIAL DIRECTIONS |
|-------------------------|--|---------------------|---|
| Redishes (continued) | Aphids, Flea Beetles, Dipterous Leafminers | ½-1 pt. per acre | Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 3 applications per season. Do not apply within 14 days of harvest or illegal residues may occur. Note: DIAZINON 4AG will not control organo-phosphate-resistant leafminers. |
| Spinach | Cutworms | 2-4 qts. per acre | Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note). |
| | Wireworms | 3-4 qts. per acre | Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil. |
| | Aphids, Dipterous Leafminers | ½-1 pt. per acre | Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 14 days of harvest. Note: DIAZINON 4AG will not control organo-phosphate-resistant leafminers. |
| Squash, Summer | Cutworms | 2-4 qts. per acre | Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note). |
| | Wireworms | 3-4 qts. per acre | Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil. |
| | Aphids, Serpentine Leafminers, Thrips, Striped Cucumber Beetles, Melonworms, Leafhoppers, Spider Mites | ½-1 ½ pts. per acre | Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 7 days of harvest or illegal residues may occur. Note: DIAZINON 4AG will not control organo-phosphate-resistant leafminers. |
| Squash, Winter | Cutworms | 2-4 qts. per acre | Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.) |

| CROP | PEST | RATE | TIMING/SPECIAL DIRECTIONS |
|---------------------------------------|--|--------------------|---|
| Squash, Winter <i>(continued!)</i> | Wireworms | 3-4 qts. per acre | Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil. |
| | | ½-1½ pts. per acre | <p>Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season.</p> <p>Do not apply within 3 days of harvest or illegal residues may occur.</p> <p>Note: DIAZINON 4AG will not control organo-phosphate-resistant leafminers.</p> |
| Sweet Corn | Cutworms | 2-4 qts. per acre | Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.) |
| | Seed Corn Maggots | 2-4 qts. per acre | Broadcast just before planting and immediately incorporate into the top 2 inches of soil. |
| | Wireworms | 3-4 qts. per acre | Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil. |
| | Corn Earworms | 1-1½ qts. per acre | Apply when silks first appear. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. |
| | Corn Rootworm (Adults) | ½-1 pt. per acre | Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. |
| | Corn Leaf Aphids | 1-2 pts. per acre | Apply in sufficient water to provide complete coverage to the upper portions of the plants. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. |
| | Grasshoppers, Spider Mites, Flea Beetles | 1 pt. per acre | Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. |
| | Sap Beetles | 2-2½ pts. per acre | Consult local agricultural authorities for proper timing of sprays. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. |

Notes: (1) To protect bees, do not apply foliarty to corn during the pollen shed period. (2) Allow 7 days after the last application before grazing or cutting for forage. Forage may be fed to beef and dairy cattle and sheep. (3) Corn may be picked 7 days after the last application.

| | | | |
|----------------|-------------------------|-------------------|---|
| Sweet Potatoes | Wireworms, Flea Beetles | 3-4 qts. per acre | Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil. |
|----------------|-------------------------|-------------------|---|

| CROP | PEST | RATE | TIMING/SPECIAL DIRECTIONS |
|-------------------------|---|--------------------|---|
| Swiss Chard | Cutworms | 2-4 qts. per acre | Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.) |
| | Wireworms | 3-4 qts. per acre | Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil. |
| | Aphids, Imported Cabbageworms, Diamondback Moths, Harlequin Bugs | ½-1 pt. per acre | Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 14 days of harvest or illegal residues may occur. |
| Tomatoes | Cutworms | 2-4 qts. per acre | Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.) |
| | Mole Crickets | 1 qt. per acre | Apply to plant beds or fields 1-2 days before sowing of seed or transplanting and immediately incorporate into the top 1-2 inches of soil. |
| | Wireworms | 3-4 qts. per acre | Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil. |
| | Aphids, Dipterous Leafminers | ½ pt. per acre | Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. |
| | Banded Cucumber Beetles, Fall Armyworms, Southern Armyworms, Beet Armyworms | ¾-1 pt. per acre | Do not apply within 1 day of harvest or illegal residues may occur. Note: DIAZINON 4AG will not control organo-phosphate-resistant leafminers. |
| | Vinegar Flies (<i>Drosophila spp.</i>) | ½-1½ pts. per acre | |
| Turnips, Roots and Tops | Cutworms | 2-4 qts. per acre | Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.) |
| | Mole Crickets | 1 qt. per acre | Apply to plant beds or fields 1-2 days before sowing of seed or transplanting and immediately incorporate into the top 1-2 inches of soil. |

| CROP | PEST | RATE | TIMING/SPECIAL DIRECTIONS |
|-------------------------------|---|-------------------|--|
| Turnips <i>(continued)</i> | Wireworms | 3-4 qts. per acre | Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil. |
| | Aphids, Flea Beetles, Dipterous Leafminers | ½-1 pt. per acre | Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 14 day of harvest or illegal residues may occur. Note: DIAZINON 4AG will not control organo-phosphate-resistant leafminers. |

FIELD CROPS

Soil and foliar applications of DIAZINON 4AG should always be applied as a spray in sufficient water to ensure thorough coverage. For foliar applications, observe stated time intervals between last application and harvest, as well as intervals between applications and total number of applications that can be made per season.

Pre-Harvest Interval (PHI*) For Foliar Applications

| CROP | PHI* (Days) |
|-------------|-------------|
| Hops | 14 |
| Sugar Beets | 14 |

*PHI - Pre-Harvest interval or the number of days between the last application and harvest.

| CROP | PEST | RATE | TIMING/SPECIAL DIRECTIONS |
|-------------|--|-------------------|---|
| Hops | Aphids, Spider Mites | 1 qt. per acre | Apply as pests occur. Repeat application, as necessary, every 14 days. Do not make more than 4 applications per season. Do not apply within 14 days of harvest. |
| Sugar Beets | Wireworms | 3-4 qts. per acre | Broadcast just before planting. Incorporate into soil 4-6 inches. Do not apply more than 4 qts. of this product to the soil per season. |
| | Beet Leafhoppers, Dipterous Leafminers, Aphids | ¾-1 pt. per acre | Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 14 days of harvest. Note: DIAZINON 4AG will not control organo-phosphate-resistant leafminers. |

| CROP | PEST | RATE | TIMING/SPECIAL DIRECTIONS |
|----------------------------|--------------|----------------|--|
| Sugar Beets (continued) | Grasshoppers | 1 pt. per acre | Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 14 days of harvest |

Note: Tops may be fed to beef and dairy animals.

GRASSLAND INSECTS

Ditch Banks, Roadsides, Noncrop Areas, Barrier Strips

Grasshoppers: Spray ¾-1 pt. per acre in water (minimum of 1 gal. of water per acre) for aerial applications or a minimum of 5 gals. of water per acre for ground applications or in oil (a minimum of 1 gal. of oil per acre) when insects first appear, preferably in the nymphal stage. Thorough coverage of the foliage is essential.

Do not graze or use treated grass for forage or feed.

Insect Control on Ornamentals

To control certain insects on Arborvitae, Azalea, Birch, Boxwood, Camelia, Carnation, Chrysanthemum, Douglas Fir, Elm, Gladioli, Hawthorn, Holly, Juniper, Lilac, Locust, Maple, Oak, Pine, Ornamental Plum, Poplar, Rhododendron, Rose, Spruce and Willow, apply the recommended rates indicated below. Try to spray underside of leaves and penetrate dense foliage.

| Insects | Amount of D-z-n Diazinon AG500 to Use | |
|---|--|-----------------------------|
| | Rate Per 3 Gals. Water | Rate Per 100 Gals. Water |
| Aphids, Bagworms, Carnation Bud Mites, Carnation Shoot Mites, Clover Mites, Cyclamen Mites, Dipterous Leafminers, European Pine Shoot Moths, European Red Mites, Flea Beetles, Holly Bud Moths, Leafhoppers, Obscure Root Weevils, Omnivorous Leaf-tiers, Privet Mites, Scale Crawlers (Cottony-cushion, Lecanium, Pine Needle, San Jose, Soft), Thrips, Two-spotted Spider Mites, Juniper Webworms, Whiteflies | ½ fl. oz.* | 1 pt.** |
| Apple-and-Thorn Skeletonizers, Cotoneaster Webworms, Fall Webworms, Hemlock Chermes, Oak Loopers, Obliquebanded Leafrollers, Pear Slugs, Tent Caterpillars | 1 ½ fl. oz.* | 3 pts.** |

* 1 fl. oz. - 2 tablespoons

** 16 fl. oz. - 1 pint

Precaution: Do not use on Ferns, Poinsettia, Hibiscus, Papaya, Piles and Gardenia, since injury to the plants may occur.

Mimosa Webworms: Mix ½ fl. oz. in 3 gals. of water (or 1 pt. in 100 gals. of water) and apply as a thorough foliar spray when adults first appear. Make a second and third application at 4-5 week intervals.

LAWN PEST CONTROL (excluding Golf Courses and Sod Farms)

Precautions: Birds, especially waterfowl, feeding or drinking on treated areas may be killed. Apply this product only as specified on this label. Because of the migratory habits of certain waterfowl, do not apply in Nassau County, New York.

between November 1 and May 20. Do not exceed maximum permitted label rates, since rates above those recommended significantly increase potential hazards to birds, especially waterfowl. Avoid overlapping sprays. If waterfowl; i.e., ducks or geese, can be expected to enter the treated area after treatment (except spot treatments), water lawn with at least 1/4 inch of water immediately after application; however, stop watering before puddling occurs.

To control pests listed below, apply recommended rates.

| Pest | Rate/1000 sq. ft. | | Remarks |
|--|----------------------------|---------|--|
| | D-2-n Diazinon AG500 | Water | |
| Lawn Chinch Bugs | 2-3 fl. oz. | 3 gals. | Use higher rate for longer residual control and in lawns of dense growth such as St. Augustine lawns. |
| Ants, Armyworms, Bermudagrass Mites, Clover Mites, Crickets, Cutworms, Digger Wasps, Earwigs, Frit Flies, Lawn Billbugs, Leafhoppers, Sod Webworms (Lawn Moths), Sowbugs (Pillbugs), Springtails (<i>Collembola</i>) | 2-3 fl.oz. | 3 gals. | Spot spray ant hills and wasp ground nest openings. For digger wasps, apply at dusk when wasps are not active. For frit flies, mow grass and water well before treatment, but delay application until grass is dry. For billbug control, apply when activity is first observed or when chewed or brown grass is present, indicating damage from this insect. |
| Brown Dog Ticks, Chiggers, Fleas | 3 fl. oz. | 3 gals. | Note: Do not apply to animals. For brown dog ticks, spray grass and under shrubbery, particularly near house. |
| Hyperodes Weevils | 3 fl. oz. | 3 gals. | Treat problem areas in mid-April and again in mid-May. |
| Millipedes | 3 fl. oz. | 3 gals. | |
| Rhodesgrass Scales | 3 fl. oz. | 3 gals. | Apply when crawlers first emerge. |
| White Grubs (such as Japanese Beetle Larvae, European Chafer, Southern Chafer) | 3 fl. oz. | 3 gals. | Apply any time between late July and early October. |

Nuisance Pests in Outside Areas

Ants, Boxelder Bugs, Brown Dog Ticks, Chiggers, Clover Mites, Cockroaches, Crickets, Earwigs, Fleas, Flies, Millipedes, Sowbugs (Pillbugs), Spiders, Springtails (*Collembola*)

Mix 3 fl. oz. in 3 gals. of water and thoroughly spray window and door screens, sills, foundations, porch and patio walls, entranceways, walks, garbage cans, tree trunks, and into cracks and other places where insects can hide. To prevent entry into houses, spray a five-foot band of soil around the house next to the foundation and spray the foundation of the house to a height of 2-3 feet. Repeat application as necessary. Do not apply to animals. Do not treat plants (other than tree trunks) at this nuisance pest rate.

NOTICE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to Seller, and Buyer assumes the risk of any such use.