

# RIVERSIDE DIAZINON 4EP

**AN EMULSIFIABLE CONCENTRATE CONTAINING 4 POUNDS DIAZINON PER GALLON.  
FOR USE BY PEST CONTROL OPERATORS ONLY.  
USEFUL FOR CONTROL OF COCKROACHES, CERTAIN OTHER HOUSEHOLD PESTS,  
AND TURF AND ORNAMENTAL PESTS.**

For specific insects and uses refer to label directions.  
DIAZINON is a trademark of CIBA-GEIGY Corporation.

ACTIVE INGREDIENTS		
Diuron (3,4-dichlorophenyl 6-methyl-4-pyridinyl phosphorothioate)	47.6%	
Water-soluble solvent	30.3%	
<b>INERT INGREDIENTS</b>	<b>22.1%</b>	
<b>DIAZINON</b>	<b>100.0%</b>	<b>TOTAL</b>
Each gallon contains 4 lbs. Diazinon per gallon.		

## WARNING: KEEP OUT OF REACH OF CHILDREN

May be harmful if swallowed. May be absorbed through skin. Do not breathe vapors. Avoid contact with eyes, nose, mouth, and skin. Wash thoroughly after handling. Avoid contamination of food and feed. Keep out of reach of children. Do not use in or near human habitation, bird pits, or livestock. Do not permit children to play in or near the sprayed area until spray has completely dried.

Toxic to fish and other aquatic life. Birds and other wildlife may be killed if they come in contact with treated areas. Do not apply where birds or other wildlife are present. Do not apply when weather conditions favor drift. Avoid contact with water by washing of equipment and disposal of wastes. Apply this product only as identified on this label. Do not use if bees are exposed to direct treatment or residues. For more information, contact your Cooperative Agricultural Extension Office.

## DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME

**READ ENTIRE LABEL BEFORE USING THIS PRODUCT.** Use of this product is restricted to those uses specifically mentioned on the label. May cause skin irritation. Avoid contact with eyes, nose, mouth, and skin. Do not breathe vapors. Avoid contact with water by washing of equipment and disposal of wastes.

## GENERAL INFORMATION

Each gallon of DIAZINON 4EP contains 4 pounds of Diazinon per gallon. Each gallon contains 4.76% Diuron, 30.3% water-soluble solvent, and 22.1% inert ingredients.

Each gallon of DIAZINON 4EP contains 4 pounds of Diazinon per gallon. Each gallon contains 4.76% Diuron, 30.3% water-soluble solvent, and 22.1% inert ingredients.

**NOTE:** This product is not intended for use on lawns, ornamentals, or other areas where it may be absorbed through the skin. Do not breathe vapors. Avoid contact with eyes, nose, mouth, and skin. Wash thoroughly after handling. Avoid contamination of food and feed. Keep out of reach of children. Do not use in or near human habitation, bird pits, or livestock. Do not permit children to play in or near the sprayed area until spray has completely dried.

## DIRECTIONS FOR USE—FOOD HANDLING ESTABLISHMENTS AND HOUSEHOLDS

Dilute RIVERSIDE DIAZINON 4EP with water to make a 1/2% or 1% Diazinon spray and apply as directed below.

Gals. of Finished Spray	Amount RIVERSIDE DIAZINON 4EP to use for 1/2% spray	1% spray
1	1.05 fl. oz.*	2.5 fl. oz.
2	2.5 fl. oz.	5.0 fl. oz.
3	3.75 fl. oz.	7.5 fl. oz.
10	12.5 fl. oz.	25.0 fl. oz. (12.6 pts.)
25	32.0 fl. oz. (2.0 pts.)	64.0 fl. oz. (4.0 pts.)

## DIRECTIONS FOR USE — FOOD HANDLING ESTABLISHMENTS (Places other than private residences in which exposed food is held, processed, prepared or served.)

**FOOD AREAS** (Application limited to crack and crevice treatment only)

In food areas, cockroaches, ants, silverfish, and other household pests may be controlled by spraying in cracks and crevices. Do not spray on food or food preparation surfaces. Do not spray in areas where food is exposed and if food is in contact with walls, do not spray on walls.

Apply a solution of RIVERSIDE DIAZINON 4EP as a 1/2% solution in cracks and crevices. Do not spray on food or food preparation surfaces. Do not spray in areas where food is exposed and if food is in contact with walls, do not spray on walls. Do not spray on food preparation surfaces. Do not spray in areas where food is exposed and if food is in contact with walls, do not spray on walls.

**SEWERAGE AREAS** (Facilities where prepared foods are served, such as dining rooms, but excluding areas where foods may be prepared or held.)

Apply a solution of RIVERSIDE DIAZINON 4EP as a 1/2% solution in cracks and crevices. Do not spray on food or food preparation surfaces. Do not spray in areas where food is exposed and if food is in contact with walls, do not spray on walls.

**NON-FOOD AREAS** (Garage rooms, incinerators, floor drains, to sewers, entries and vestibules, utility, boiler rooms, machine rooms, boiler rooms, garages, mop closets, and storage after cleaning or mopping.)

Apply a solution of RIVERSIDE DIAZINON 4EP as a 1/2% solution in cracks and crevices. Do not spray on food or food preparation surfaces. Do not spray in areas where food is exposed and if food is in contact with walls, do not spray on walls.

## DIRECTIONS FOR USE — HOUSEHOLD INSECTS

**COCKROACHES (WATERBUGS), ANTS AND SILVERFISH**—Make spot applications in cracks and crevices. Do not spray on food or food preparation surfaces. Do not spray in areas where food is exposed and if food is in contact with walls, do not spray on walls.

For heavy cockroach infestations, make the first application at a strength of 1% Diazinon, followed by monthly applications of 1/2% Diazinon.

**CARPET BEETLES**—Apply a solution of RIVERSIDE DIAZINON 4EP as a 1/2% solution in cracks and crevices. Do not spray on food or food preparation surfaces. Do not spray in areas where food is exposed and if food is in contact with walls, do not spray on walls.

**SPIDERS AND SCORPIONS**—Apply a solution of RIVERSIDE DIAZINON 4EP as a 1/2% solution in cracks and crevices. Do not spray on food or food preparation surfaces. Do not spray in areas where food is exposed and if food is in contact with walls, do not spray on walls.

**PANTRY PESTS**—Exposed stages (Saw-Toothed Grain Beetle, Confused Flour Beetle, Rice Weevil, Cigarette Beetle, Drugstore Beetle, Indian Meal Moth). Apply a solution of RIVERSIDE DIAZINON 4EP as a 1/2% solution in cracks and crevices. Do not spray on food or food preparation surfaces. Do not spray in areas where food is exposed and if food is in contact with walls, do not spray on walls.

**BROWN DOG TICKS IN BUILDINGS**—Make spot applications in cracks and crevices. Do not spray on food or food preparation surfaces. Do not spray in areas where food is exposed and if food is in contact with walls, do not spray on walls.

## DIRECTIONS FOR USE—ORNAMENTAL INSECTS

To control certain insects on Aspidistra, Asplenium, Buxus, Camellia, Carnation, Chrysanthemum, Douglas Fir, Elm, Euonymus, Holly, Juniper, Liquidambar, Lycium, Oak, Pine, Plum, Pyrus, Rhododendron, Rose, Spruce, and Walden apply the recommended rates indicated below.

Do not use on certain Ferns, Poinsettia, Hibiscus, Papaya, Freesia, and Gardenia since injury to the plants may occur.

**MIMOSA WEBWORMS**—Mix 1/2 fl. oz. of 1/2% solution in 3 pts. water or 1 pt. in 100 gal. water and apply as a thorough foliar spray when adults first appear. Make a second and third application at 4 week intervals.

## RATE OF APPLICATION

Insects	Amount RIVERSIDE DIAZINON 4EP to use per 3 gal. water	per 100 gal. water
Aspidistra		
Asplenium		
Buxus		
Camellia		
Chrysanthemum		
Douglas Fir		
Elm		
Euonymus		
Holly		
Juniper		
Liquidambar		
Lycium		
Oak		
Pine		
Plum		
Pyrus		
Rhododendron		
Rose		
Spruce		
Walden		
Aspidistra	1/2 fl. oz.*	1 pt.
Asplenium		
Buxus		
Camellia		
Chrysanthemum		
Douglas Fir		
Elm		
Euonymus		
Holly		
Juniper		
Liquidambar		
Lycium		
Oak		
Pine		
Plum		
Pyrus		
Rhododendron		
Rose		
Spruce		
Walden		

## DIRECTIONS FOR USE—LAWN INSECTS

To control certain insects on lawns, apply a solution of RIVERSIDE DIAZINON 4EP as a 1/2% solution in cracks and crevices. Do not spray on food or food preparation surfaces. Do not spray in areas where food is exposed and if food is in contact with walls, do not spray on walls.

**BOXELDER BUGS**—Apply a solution of RIVERSIDE DIAZINON 4EP as a 1/2% solution in cracks and crevices. Do not spray on food or food preparation surfaces. Do not spray in areas where food is exposed and if food is in contact with walls, do not spray on walls.

Insects	Amount RIVERSIDE DIAZINON 4EP to use per 3 gal. water	per 100 gal. water
Aspidistra		
Asplenium		
Buxus		
Camellia		
Chrysanthemum		
Douglas Fir		
Elm		
Euonymus		
Holly		
Juniper		
Liquidambar		
Lycium		
Oak		
Pine		
Plum		
Pyrus		
Rhododendron		
Rose		
Spruce		
Walden		
Aspidistra	1/2 fl. oz.*	1 pt.
Asplenium		
Buxus		
Camellia		
Chrysanthemum		
Douglas Fir		
Elm		
Euonymus		
Holly		
Juniper		
Liquidambar		
Lycium		
Oak		
Pine		
Plum		
Pyrus		
Rhododendron		
Rose		
Spruce		
Walden		

Apply a solution of RIVERSIDE DIAZINON 4EP as a 1/2% solution in cracks and crevices. Do not spray on food or food preparation surfaces. Do not spray in areas where food is exposed and if food is in contact with walls, do not spray on walls.

## DISPOSAL OF CONTAINER

Do not reuse empty container. Destroy by perforating or crushing and disposing in a sanitary manner.

## NOTICE

This product is not intended for use on lawns, ornamentals, or other areas where it may be absorbed through the skin. Do not breathe vapors. Avoid contact with eyes, nose, mouth, and skin. Wash thoroughly after handling. Avoid contamination of food and feed. Keep out of reach of children. Do not use in or near human habitation, bird pits, or livestock. Do not permit children to play in or near the sprayed area until spray has completely dried.

EPA REG. NO. 9779-12 NET CONTENTS 1 GALLON

EPA Est. 9779-AL-12, AR-12, GA-1, LA-1, MS-12, TN-1, TX-1

