

AUG 2 1974

RIVERSIDE PECAN INSECTICIDE

AN EMULSIFIABLE CONCENTRATE CONTAINING 4 POUNDS OF EPN AND 2.5 POUNDS OF PARATHION PER GALLON.
USEFUL FOR THE CONTROL OF CERTAIN INSECT PESTS OF PECANS.

USE ONLY WHEN WEARING THE FOLLOWING PROTECTIVE EQUIPMENT AND CLOTHING
1. Wear water proof pants, coat, hat, rubber boots, and gloves. 2. Wear safety goggles. 3. Wear a respirator with a filter. 4. Wear safety shoes. 5. Wear safety glasses. 6. Wear safety glasses and a respirator with a filter. 7. Wear safety glasses and a respirator with a filter. 8. Wear safety glasses and a respirator with a filter.

WORK SAFETY RULES
1. Keep children and pets away from treated areas. 2. Do not eat, drink, or smoke while working with this product. 3. Wash hands thoroughly after use. 4. Do not use near open flames. 5. Do not use in confined spaces. 6. Do not use in windy conditions. 7. Do not use in rain. 8. Do not use in temperatures below 50°F. 9. Do not use in temperatures above 90°F. 10. Do not use in areas where bees are present.

POISON SIGNS (Symptoms)
1. Nausea and vomiting. 2. Headache and dizziness. 3. Blurred vision. 4. Difficulty breathing. 5. Weakness and fatigue. 6. Irritation of the skin. 7. Irritation of the eyes. 8. Irritation of the nose and throat. 9. Irritation of the lungs. 10. Irritation of the stomach and intestines. 11. Irritation of the kidneys. 12. Irritation of the liver. 13. Irritation of the heart. 14. Irritation of the nervous system. 15. Irritation of the reproductive system.

FIRST AID TREATMENT
IF SWALLOWED: Induce vomiting by drinking water or milk. Do not give anything if the person is unconscious. IF BREATHING HAS STOPPED: Start artificial respiration immediately. IF IN CONTACT WITH EYES: Flush with water for at least 15 minutes. IF IN CONTACT WITH SKIN: Wash with soap and water.

IN CASE OF CONTACT
Immediately flush eyes with water for at least 15 minutes. Wash contaminated clothing and shoes. Do not use if you are pregnant or nursing.

NOTE TO PHYSICIAN
Antidote: Atropine. Symptoms: Nausea, vomiting, headache, dizziness, blurred vision, difficulty breathing, weakness, fatigue, irritation of the skin, eyes, nose, throat, lungs, stomach, intestines, kidneys, liver, heart, nervous system, reproductive system.

POST TREATED AREA
Do not eat, drink, or smoke in treated areas. Do not use treated areas for food storage. Do not use treated areas for animal husbandry.

ACTIVE INGREDIENTS	
O-ethyl O-p-nitrophenyl phenylphosphonothioate	4.4%
Parathion (O,O-diethyl O-p-nitrophenyl phosphorothioate)	25.9%
Other inert ingredients	15.5%
INERT INGREDIENTS	15.5%
TOTAL	100.0%

STOP - READ LABEL BEFORE USING!

POISON
SEE SIDE PANELS FOR ANTIDOTE AND PRECAUTIONS




DANGER
KEEP OUT OF THE REACH OF CHILDREN

NOT FOR HOME USE



DO NOT SWALLOW
CAN KILL IF SWALLOWED



DO NOT TOUCH
CAN KILL BY SKIN CONTACT



DO NOT BREATHE
CAN KILL IF BREATHED

POISONOUS TO FISH AND WILDLIFE
Do not use near water bodies. Do not use in areas where birds and other wildlife are present.

HIGHLY TOXIC TO BEES
Do not use when bees are present. Do not use in areas where bees are present.

GENERAL INFORMATION
RIVERSIDE PECAN INSECTICIDE is a concentrate of Parathion and O-ethyl O-p-nitrophenyl phenylphosphonothioate. It is used for the control of certain insect pests of pecans. It is highly toxic to humans and animals. It is also highly toxic to bees and other beneficial insects.

TIMING OF SPRAY APPLICATIONS
Timing of spray applications is essential. Make applications according to the directions for use. Do not use in windy conditions. Do not use in rain. Do not use in temperatures below 50°F. Do not use in temperatures above 90°F.

DIRECTIONS FOR USE
Apply to pecan trees during the growing season. Apply to the leaves and twigs. Apply to the soil around the base of the tree. Do not use on other plants. Do not use on animals. Do not use on humans.

DO NOT USE POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME
HEAD ENTIRE LABEL BEFORE USING THIS PRODUCT

FALL WEEDING
Use for the control of certain insect pests of pecans. Use for the control of certain insect pests of pecans.

REGISTERED TRADEMARK
RIVERSIDE PECAN INSECTICIDE is a registered trademark of the manufacturer. All other trademarks are the property of their respective owners.

INSECTICIDE

GENERAL INFORMATION
 RIVERSIDE PECAN INSECTICIDE is a combination of Dursin, diethyl para-
 and ERB...
 pecans is an important part of the pecan production program. The insect pests
 that attack pecans are numerous and their damage is often severe. The use of
 insecticides is necessary to protect the crop and to insure a high quality
 product. Riverside Pecan Insecticide is a highly effective and safe insecticide
 for the control of these pests. It is a mixture of organic phosphate compounds
 and is easy to apply. It is effective against a wide range of insect pests
 and is safe for the user and the environment. It is a valuable addition to
 any pecan grower's insecticide program.

MODE OF APPLICATION
 Riverside Pecan Insecticide is applied as a spray to the leaves and
 twigs of the pecan tree. It is most effective when applied during the
 growing season. The spray should be applied to the undersides of the
 leaves and to the twigs. It is important to cover the entire tree
 thoroughly. The spray should be applied every 7 to 10 days during the
 growing season. It is important to follow the directions on the label
 for the most effective results.

MODE OF SPRAY APPLICATIONS
 To obtain adequate control of insect pests, the quantity
 of spray applications is essential. Make applications according to the
 directions for use. The spray should be applied to the undersides of the
 leaves and to the twigs. It is important to cover the entire tree
 thoroughly. The spray should be applied every 7 to 10 days during the
 growing season. It is important to follow the directions on the label
 for the most effective results.

**DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR
 IN FLAME**
 Read the entire label before using this product.
 Do not use, pour, spill, or store near heat or in flame.
 Do not use in home plantings.

DIRECTIONS FOR USE
 Apply Riverside Pecan Insecticide to the undersides of the leaves
 and to the twigs of the pecan tree. It is most effective when applied
 during the growing season. The spray should be applied every 7 to 10
 days during the growing season. It is important to follow the directions
 on the label for the most effective results.

DISPOSAL OF CONTAINER AND WASTES
 Do not reuse this container for ANY purpose. When empty, it should
 be disposed of properly. Do not dump it into the water, soil, or any other
 place. It should be disposed of in a way that will not harm the
 environment. Do not use in home plantings.

NOTICE
 Riverside Chemical Co. warrants that the product conforms to its
 label. It does not warrant that the product will be effective under all
 conditions. It is the user's responsibility to follow the directions on
 the label. Do not use in home plantings.

REGISTERED TRADEMARKS
 RIVERSIDE CHEMICAL CO. is a registered trademark of Riverside Chemical Co.
 Do not use in home plantings.

MITES—Apply RIVERSIDE PECAN INSECTICIDE when mites or their damage
 are noticed.

PECAN LEAF CASEBEARERS—Application of RIVERSIDE PECAN INSECTI-
 CIDE in the first pre-pollination spray will control pecan leaf casebearers. Its
 inclusion in cover sprays from late June until Mid-August will reduce leaf
 casebearer damage during the following season.

PECAN NUT CASEBEARERS—Apply RIVERSIDE PECAN INSECTICIDE
 when nuts begin to turn brown. Repeat in 7 days and in subsequent cover
 sprays if needed.

PECAN WEEVILS—The first application of RIVERSIDE PECAN INSECTICIDE
 should be made when the pecan shells begin to harden on the earliest varie-
 ties. For better control of nut drop results from adult feeding. Make appli-
 cations every 7 days until Mid-September and even later if adult pecan weevils
 are still found in the grove.

SPITTLE BUGS—If past experience indicates spittlebugs will be a problem,
 apply RIVERSIDE PECAN INSECTICIDE to the second pre-pollination spray.
 Otherwise apply when the bugs are noticed.

TWIG GIRDLERS—Apply RIVERSIDE PECAN INSECTICIDE when twig gird-
 lers are found making 3 to 5 applications at 2 to 3 week intervals.


WALNUT CATERPILLAR—Apply RIVERSIDE PECAN INSECTICIDE when
 caterpillars are found and are still small.

Do not apply later than 21 days prior to harvest. Do not apply after husks
 open. Do not feed treated husks to livestock. DO NOT GRAZE LIVESTOCK
 ON TREATED GROVES. DO NOT USE IN HOME PLANTINGS.

DISPOSAL OF CONTAINER AND WASTES
 Do not reuse this container for ANY purpose. When empty, it should
 be disposed of properly. Do not dump it into the water, soil, or any other
 place. It should be disposed of in a way that will not harm the
 environment. Do not use in home plantings.

NOTICE
 Riverside Chemical Co. warrants that the product conforms to its
 label. It does not warrant that the product will be effective under all
 conditions. It is the user's responsibility to follow the directions on
 the label. Do not use in home plantings.

REGISTERED TRADEMARKS
 RIVERSIDE CHEMICAL CO. is a registered trademark of Riverside Chemical Co.
 Do not use in home plantings.



ORGANIC PHOSPHATE COMPOUNDS, MIXTURE, LIQUID