
KEEP OUT OF REACH OF CHILDREN - 12 point

DANGER -red - 18 point

PELIGRO - red

POISON

VENENO

Skull & Crossbones - red
Can Cause Death

PRECAUCION
AL USUARIO

Si usted no lee ingles,
no use este producto
hasta que la etiqueta
le haya sido explicada
ampliamente.

STATEMENT OF PRACTICAL TREATMENT

FIRST AID TREATMENT + Call a doctor (physician), clinic, or hospital immediately. Explain that the victim has been exposed to methyl parathion and describe his condition.

IF BREATHING HAS STOPPED, start artificial respiration immediately and maintain until doctor sees victim.

IF SWALLOWED, call a physician or poison control center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger, or if available, by administering syrup of ipecac. Do not induce vomiting or give anything by mouth to an unconscious person.

IN CASE OF CONTACT, immediately flush eyes or skin with plenty of water for at least 15 minutes while removing contaminated clothing and shoes. See doctor immediately.

See left side panel for additional PRECAUTIONARY STATEMENTS.

EPA Reg. No. 9779-218

EPA Est. No. 9779-AL-1

Manufactured For

RIVERSIDE/TERRA CORPORATION

Terra Centre, 600 Fourth Street, Sioux City, Iowa 51101

Riverside Serves Agriculture. Agriculture Serves Everyone.

NET CONTENTS

_____ GALLONS