


FRONT PANEL

RED-I-CAT
Concentrate

Contains PROLIN, New Improved Warfarin Rodenticide (Warfarin plus anti-bacterial agent).



Kills RATS & MICE . . . For Home Bait Mixing

Active ingredients:

| | |
|--|------|
| Warfarin 3-(a-acetonylbenzyl)-4-hydroxycoumarin | 0.5% |
| Sulfaquinoxaline (N'-(2-quinoxaliny) sulfanilamide | 0.5% |

Inert ingredients:

| | |
|--------|--------|
| Total: | 99.0% |
| | 100.0% |

Prolin is a Trade Mark of Wisconsin Alumni Research Foundation.

CAUTION: Keep out of reach of children.
(See back panel for additional cautions)

BACK PANEL

DIRECTIONS

RATS: Mix thoroughly one part of Red-I-Cat Concentrate with 19 parts of a suitable bait material which is available, such as corn meal, rolled oats, dog food, laying mash, stock food, etc. A small amount of a bland oil, such as mineral, corn, peanut, or any other edible oil, may be used up to 5% when the bait is so dry that it might have a tendency to dust.

When perishable baits, such as ground meat, fish, vegetables or similar material are used, bait placements should be made in the late afternoon and replaced daily with fresh bait.

Baits should be placed in amounts of 4 oz. up to 1 lb. in locations frequented by rats or where rats have been known to feed. Baits must be protected from dogs, cats, poultry, livestock, etc., by bait boxes, cages, or other suitable means.

Baits should be examined regularly and replenished before completely eaten. If bait molds, decomposes or becomes sour, it should be replaced. An uninterrupted supply of bait should be maintained for at least 10 days.

MICE: For controlling mice, follow the same general directions as for rats. Table-spoonful amounts of bait should be placed at 8 to 12 foot intervals.

Mice are more difficult to control than rats and complete control may take a longer period of baiting. An uninterrupted supply of bait should be maintained for at least 15 days.

IMPORTANT

Baiting should be continued as long as evidence of feeding is observed. Except in extremely heavy infestations, this should be from 5 days to 2 weeks. Where a continuous source of infestation is present from nearby dumps or fields, permanent bait stations should be used and bait placed in them as often as necessary.

CAUTION

Red-I-Cat Concentrate contains as its active ingredient, an anti-coagulant chemical, therefore baits prepared with Red-I-Cat Concentrate must be protected from or placed where humans, livestock, poultry or pets will not eat them. In case baits are eaten, give a tablespoonful of salt in a glass of warm water and