




MALACIDE

SWINE DUST

4



Contains 4% Malathion

DIRECTIONS FOR USE

HOGS: For control of Lice treat animal thoroughly. Repeat in 10 days and thereafter as needed. Avoid contamination of feed, food containers and watering troughs.

POULTRY: Direct Application for control of Northern Fowl Mite, Poultry Lice, and Chicken Red Mite. Dust individual birds with a can shaker or rotary hand duster. Repeat in 4 to 8 weeks if necessary. For Chicken Red Mite control this is a supplement to the premise treatment. Premise Treatment for control of Northern Fowl Mite, Poultry Lice, Chicken Red Mite, and Flies. Apply 1 lb. per 50-60 sq. ft. of litter and floor space and to nest, roost, and adjacent areas. Apply with a rotary hand duster, puff duster or by sprinkling from a can. Dust Bath Boxes for control of Northern Fowl Mite and Poultry Lice. Use 1 lb. per box (18"x12"x3") per 30 hens. Remove boxes when dust has been used. Dust Bath Boxes or Natural Soil Wallows for Control of Stick-tight Flea. Use 15 lbs. per box (2'x3'x1') per 100 birds or apply 7½ lbs. in natural soil wallows per 50 birds. Brooder House for control of Stick-tight Flea. Use 1 lbs. per 100 chickens. Distribute the dust evenly in each corner of the brooding house.

NOTICE: Avoid contamination of feed and feeding troughs and water fountains with sprays or dusts.

WARNING

KEEP OUT OF REACH OF CHILDREN

ACTIVE INGREDIENTS:

Malathion (O, O-dimethyl Dithiophosphate of Diethylmercaptosuccinate) 4.0%
 INERT INGREDIENT 96.0%
 TOTAL 100.0%

U.S.D.A. REG. NO. 240-137

Seller's guarantee shall be limited to the terms of the label, and subject thereto the buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.

CATTLE (Beef and Non-Milking), **HORSES:** For Lice control treat animal thoroughly. Repeat in 10 days and thereafter as needed. For Tick control repeat application at 2 week intervals if needed. For Horn Fly control apply 4 tablespoons on back and neck of the animals and repeat applications at 10-14 day intervals. Do not treat animals under one month of age. Do not apply to lactating dairy animals or non-lactating dairy animals within 2 weeks of freshening. Avoid contamination of feed, food containers and watering troughs.

DAIRY CATTLE: For Horn Fly control apply 4 tablespoons on back and neck of the animals and repeat applications at 10-14 day intervals. Do not apply to dairy cattle later than 5 hours before milking or during milking. Do not contaminate water troughs.

Pet Quarters (Dog & Cats): For Flea control remove debris, then apply 1-2 lbs. per 1,000 square feet of surface. Repeat when necessary.

CAUTION: Avoid breathing of dust or spray mist. Avoid contact with skin; wash thoroughly after using. Avoid contamination of feed and foodstuffs. Keep out of reach of children.

Not for use or storage in or around the home.

Manufactured by

DALY-HERRING COMPANY

Ahoskie, N. C. 27910 Greensboro, N. C. 27408 Dillon, S. C. 29536
 KINSTON, N. C. 28501





MALACIDE 4

SWINE DUST

Contains 4% Malathion

DIRECTIONS FOR USE

HOGS: For control of Lice treat animal thoroughly. Repeat in 10 days and thereafter as needed. Avoid contamination of feed, food containers and watering troughs.

POULTRY: Direct Application for control of Northern Fowl Mite, Poultry Lice, and Chicken Red Mite. Dust individual birds with a can shaker or rotary hand duster. Repeat in 4 to 8 weeks if necessary. For Chicken Red Mite control this is a supplement to the premise treatment. **Premise Treatment for control of Northern Fowl Mite, Poultry Lice, Chicken Red Mite, and Flies:** Apply 1 lb. per 50-60 sq. ft. of litter and floor space and to nest, roost, and adjacent areas. Apply with a rotary hand duster, puff duster or by sprinkling from a can. **Dust Bath Boxes for control of Northern Fowl Mite and Poultry Lice:** Use 1 lb. per box (18"x12"x3") per 30 hens. Remove boxes when dust has been used. **Dust Bath Boxes or Natural Soil Wallows for Control of Stick-tight Flea:** Use 15 lbs. per box (2'x3'x1') per 100 birds or apply 7 1/2 lbs. in natural soil wallows per 50 birds. **Brooder House for control of Stick-tight Flea:** Use 1 lb. per 100 chickens. Distribute the dust evenly in each corner of the brooding house.

NOTICE: Avoid contamination of feed and feeding troughs and water fountains with sprays or dusts.

WARNING
KEEP OUT OF REACH OF CHILDREN

ACTIVE INGREDIENTS:
Malathion (O, O-dimethyl Dithiophosphate of Diethylmercaptosuccinate) 4.0%
INERT INGREDIENT: 96.0%
TOTAL 100.0%

U.S.D.A. REG. NO. 240-137

Seller's guarantee shall be limited to the terms of the label, and subject thereto the buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.

CATTLE
Lice control repeat needed on back cations, mals, lactating animals, tami, troughs.

DAIRY
tablespoons repeat apply to milking water troughs.

Pet Quads move ca feet of s

CAUTION
Avoid using. Keep out.

Not for



Manufactured by
DALY-HERRING COMPANY

Ahoskie, N. C. 27910 Greensboro, N. C. 27408 Dillon, S. C. 29536
KINSTON, N. C. 28501

BEST DOCUMENT AVAILABLE

MALACIDE

SWINE DUST

Contains 4% Malathion

DIRECTIONS FOR USE

For control of Lice treat animal thoroughly at in 10 days and thereafter as needed. Contamination of feed, food containers and troughs.

Direct Application for control of Fowl Mite, Poultry Lice, and Chicken Dust individual birds with a can shaker and duster. Repeat in 4 to 8 weeks if needed. For Chicken Red Mite control this is a treatment to the premise treatment. For control of Northern Fowl Mite, Poultry Lice, and Chicken Red Mite, and Flies, Apply 1 lb. per 60 sq. ft. of litter and floor space and roost, and adjacent areas. Apply with a hand duster, puff duster or by sprinkling. Dust Bath Boxes for control of Northern Fowl Mite and Poultry Lice, Use 1 lb. per box (2'x3'x1') per 30 hens. Remove boxes when used. Dust Bath Boxes or Natural Soil Wallows for Control of Stick-tight Flea. Use 1 lb. per box (2'x3'x1') per 100 birds or apply in natural soil wallows per 50 birds. For control of Stick-tight Flea. Use 1 lb. per 100 chickens. Distribute the dust evenly in the brooding house.

Avoid contamination of feed and feeding and water fountains with sprays or dusts.

ACTIVE INGREDIENTS:

Malathion (O, O-dimethyl Dithiophosphate of Diethylmercaptosuccinate) 4.0%

INERT INGREDIENT: 96.0%

TOTAL 100.0%

U.S.D.A. REG. NO. 240-137

Seller's guarantee shall be limited to the terms of the label, and subject thereto the buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.

Manufactured by

DALY-HERRING COMPANY

Ahoscie, N. C. 27910 Greensboro, N. C. 27408 Dillon, S. C. 29536
KINSTON, N. C. 28501

4



CATTLE (Beef and Non-Milking), **HORSES**: For Lice control treat animal thoroughly. Repeat in 10 days and thereafter as needed. For Tick control repeat application at 2 week intervals if needed. For Horn Fly control apply 4 tablespoons on back and neck of the animals and repeat applications at 10-14 day intervals. Do not treat animals under one month of age. Do not apply to lactating dairy animals or non-lactating dairy animals within 2 weeks of freshening. Avoid contamination of feed, food containers and watering troughs.

DAIRY CATTLE: For Horn Fly control apply 4 tablespoons on back and neck of the animals and repeat applications at 10-14 day intervals. Do not apply to dairy cattle later than 5 hours before milking or during milking. Do not contaminate water troughs.

Pet Quarters (Dog & Cats): For Flea control remove debris, then apply 1-2 lbs. per 1,000 square feet of surface. Repeat when necessary.

CAUTION: Avoid breathing of dust or spray mist. Avoid contact with skin; wash thoroughly after using. Avoid contamination of feed and foodstuffs. Keep out of reach of children.

Not for use or storage in or around the home.



15

DIRECTIONS FOR USE:

General: Mice infestations occur mostly in spring and fall. Check your home and seal all possible areas of entry, such as pipes, conduit and door entries.

Mice Habits: Because mice are nervous feeders and follow no particular feeding pattern, it is important that smaller portions of Okay Mice Kill be placed at intervals of 8 to 12 feet.

Amount: Place tablespoon of Okay Mice Kill in shallow containers at intervals of 8 to 12 feet where mice are known to run.

Kitchen area: Set under stoves, refrigerators and lower part of kitchen cabinets (drain pipe under kitchen sink). Set under SHELTERED areas along the wall.

Basement area: Set Okay Mice Kill on top of heating ducts and on top of plate on foundation wall. Bait should be set along walls, provided it is placed behind sheltered areas, out of reach of children and pets.

When Okay Mice Kill is placed in garages and enclosed patios, it should be placed in enclosed, locked metal bait stations.

OKAY

MICE KILL

ACTIVE INGREDIENTS:

N¹-(2-Quinoxaliny1) Sulfanilamide
(sulfaguinoxaline)----- 0.025%
3-(alpha-acetonyl-benzyl)-4-
hydroxycoumarin (warfarin) - 0.025%

INERT INGREDIENTS ----- 99.950%

Total ----- 100.000%

Contains Prolin* a better, more effective Warfarin rodenticide-(warfarin plus sulfaguinoxaline) is manufactured under license from the Wisconsin Alumni Research Foundation, U. S. Patent No. 2,427,578, other patent applications pending. The antibacterial agent, sulfaguinoxaline, increases the effectiveness of this bait and assures more thorough control of rats and mice.

CAUTIONS: KEEP OUT OF REACH OF CHILDREN AND PETS, WHEN USING-WHEN STORING.

SEE SIDE PANEL FOR ADDITIONAL CAUTIONS.

LEVENSCHON CHEMICAL COMPANY
1407 Harney St., Omaha, Nebraska

CAUTIONS:

Read Label Before Using.

Keep away from humans, domestic animals and pets. This material may reduce the clotting ability of the blood and cause bleeding. In such cases, intravenous and oral administration of Vitamin K combined with blood transfusions are indicated as in the case of hemorrhage caused by overdoses of bishydroxycoumarin.

Bait should be placed so as to protect humans, pets and domestic animals.

ACCEPTED

43954-4

JUL 19 1967

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
ED UNDER NO. 43954-4 SUBJECT
TO ATTACHED COMMENTS.