

# 7.8% ZINEB

# 10% SEVIN®


## Active Ingredients:

Zineb (Zinc Ethylene Bisdithiocarbamate) ..... 7.8%

Carbaryl (1-Naphthyl N-Methylcarbamate) ..... 10.0%

Inert Ingredients: ..... 82.2%

TOTAL ..... 100.0%



## WARNING

**KEEP OUT OF REACH OF CHILDREN**

## Directions For Use

**TOMATOES:** For the control of Late and Early Blight, Flea Beetle, European Corn Borer, Tomato Fruit Worm, Fall Armyworm. Apply at the rate of 20 pounds per acre. Make first application as soon as the plants are growing well or insects or blight appears. Repeat not less than once a week thereafter. Treatment should be repeated after each rain that is heavy enough to wash the dust off. *Do not apply within 5 days of harvest.*

**WARNING! HARMFUL IF INHALED OR SWALLOWED.** Do not breathe dust. Do not take internally. Skin Contact May be harmful. Avoid Contact. Wash Hands before Eating. Take Shower or Bath after Work. Wear Regular Long Sleeve Work Clothing. Change to Clean Clothing Daily.

Note to Physician: Carbaryl is a moderate reversible Cholinesterase inhibitor. Atropine is Antidotal. **KEEP OUT OF REACH OF CHILDREN.**

E. P. A. Reg. No. 240-154

\* Trade mark of UCC for Carbaryl.

This product is highly toxic to bees exposed to direct treatment or residue on plants while bees are active. Do not apply directly to beehives. May be obtained from your Cooperative Agent or retail dealer.

Dispose of waste by burying in non-crop lands away from water supplies. Containers should be disposed by burning. Keep out of Smoke.

**NOTICE OF WARRANTY** - Seller's guarantee shall be limited to the terms of the label, and subject thereto the buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.

Manufactured By

**DALY-HERRING COMPANY**

Kinston, N.C. 28501

Greensboro, N.C. 27405

Ahoskie, N.C. 27910

Dillon, S.C. 29536