


779-248

FRONT PANEL



# HARLEQUIN VEGETABLE DUST

E.P.A. Reg. No. 240-213

**ACTIVE INGREDIENTS:**

*Endosulfan (Hexachlorohexahydromethano-2,4,3-benzodioxathiepin oxide) .....	3.0%
INERT INGREDIENTS: .....	97.0%
*Thiodan U. S. Pat. No. 2,799,685 .....	100.0%

## CAUTION

### KEEP OUT OF REACH OF CHILDREN


Harmful if swallowed, inhaled or absorbed through skin. Avoid breathing of dust. Avoid contact with skin, eyes, or clothing. In case of contact with skin or eyes, flush with plenty of water for eyes; get medical attention. Wash with soap and water after handling and before eating or smoking; wear clean clothing. Avoid contamination of food and feedstuffs. Do not apply or allow to drift to areas cultivated by or protected humans or beneficial animals.

SEE OTHER WARNINGS ON BACK PANEL

E.P.A. EST. No. 240-NC-1

**NET WEIGHT FOUR POUNDS**

BACK PANEL



## HARLEQUIN VEGETABLE DUST DIRECTIONS FOR USE

Apply the listed amounts per acre when insects first appear and repeat as required to maintain effective control. Coverage of both upper and lower leaf surfaces is essential for good control. Observe use limitations. If insect control is required beyond the following use patterns, supplement the control program with other suitable pesticides. Observe days interval between the last application and harvest indicated by number in ( ) following crop

OR

**COLLARDS** (21) Cabbage Looper, Diamondback Moth Larvae, Harlequin Bug, Imported Cabbage Worm—25 lbs. Do not exceed one application.

**BEANS**, succulent type (3) **DRY BEANS** (3) **SOUTHERN FIELD PEAS** (3); Black Bean Aphid, Bean Leaf Skeletonizer, Cucumber Beetles, Green Stink Bug, Beans Beetle—17 to 33 lbs. Do not exceed 3 applications to bean. Do not feed treated threshings or allow livestock to graze in treated fields. Do not use on lima beans that are to be removed from field for processing.

**BROCCOLI** (7), **CABBAGE** (7), **BRUSSELS SPROUTS** (14), **CAULIFLOWER** (14); Cabbage Aphid, Cabbage Looper, Cross-striped Cabbage Worm, Flea Beetles, Harlequin Bugs, Stink Bugs, Diamond-back Moth Larvae, Imported Cabbage Worm—25 to 33 lbs.

**CUCUMBERS** (0), **SUMMER AND WINTER SQUASH** (0): Aphids, Cucumber Beetles, Striped Cabbage Flea Beetle, Squash Vine Borer—25 to 33 lbs. For Vine Borer control, apply weekly to flower beds, stems and vines beginning when moths first appear.

**EGGPLANTS**: (1): Blister Beetle, Colorado Potato Beetle, Flea Beetle, Green Peach Aphid, Green Stink Bug, Whitefly—20 lbs.

**MUSTARD GREENS** (21), **TURNIP GREENS** (21): Aphids, Cabbage Looper, Imported Cabbage Worm—25 lbs. Do not exceed one application

### USAGE CAUTIONS

Workers entering treated fields within 24 hours of application should wear protective clothing.

Do not plant root crops other than carrots, potatoes, sugar beets, and sweet potatoes as follow-up crops.

This product is toxic to fish and wildlife. Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes. Birds feeding on treated areas may be killed. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Apply this product only as specified on this label.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

**DISPOSE OF WASTES** by burying in non-crop lands away from water supplies. Container should be disposed of by burying with wastes or by burning. Keep out of smoke.

# HARLEQUIN VEGETABLE DUST

E.P.A. Reg. No. 240-213

**ACTIVE INGREDIENTS:**

*Endosulfan (Hexachlorohexahydromethano-2,4,3-benzodioxathiepin oxide) .....	3.0%
INERT INGREDIENTS: .....	97.0%
*Thiodan U.S. Pat. No. 2,799,685	100.0%

**CAUTION**

**KEEP OUT OF REACH OF CHILDREN**

Harmful if swallowed, inhaled or absorbed through skin. Avoid breathing of dust. Avoid contact with skin, eyes, or clothing. In case of contact with skin or eyes, flush with plenty of water for eyes, get medical attention. Wash with soap and water after handling and before eating or smoking; wear clean clothing. Avoid contamination of feed and foodstuffs. Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals.

SEE OTHER WARNINGS ON BACK PANEL

E.P.A. EST. No. 240-NC-1

**NET WEIGHT FOUR POUNDS**

Thiodan is a registered trademark of Canadian Hoechst Limited.

MANUFACTURED BY

**DALY-HERRING COMPANY**

KINSTON, NORTH CAROLINA 28501

is required beyond the following use patterns, supplement the control program with other suitable pesticides. Observe days interval between the last application and harvest indicated by number in ( ) following crop

OR

**COLLARDS** (21) Cabbage Looper, Diamondback Moth Larvae, Harlequin Bug, Imported Cabbage Worm—25 lbs. Do not exceed one application.

**BEANS**, succulent type (3) **DRY BEANS** (3) **SOUTHERN FIELD PEAS** (3); Black Bean Aphid, Bean Leaf Skeletonizer, Cucumber Beetles, Green Stink Bug, Beans Beetle—17 to 33 lbs. Do not exceed 3 applications to bean. Do not feed treated threshings or allow livestock to graze in treated fields. Do not use on lima beans that are to be removed from field for processing.

**BROCCOLI** (7), **CABBAGE** (7), **BRUSSELS SPROUTS** (14), **CAULIFLOWER** (14); Cabbage Aphid, Cabbage Looper, Cross-striped Cabbage Worm, Flea Beetles, Harlequin Bugs, Stink Bugs, Diamond-back Moth Larvae, Imported Cabbage Worm—25 to 33 lbs.

**CUCUMBERS** (0), **SUMMER AND WINTER SQUASH** (0): Aphids, Cucumber Beetles, Striped Cabbage Flea Beetle, Squash Vine Borer—25 to 33 lbs. For Vine Borer control, apply weekly to flower beds, stems and vines beginning when moths first appear.

**EGGPLANTS**: (1): Blister Beetle, Colorado Potato Beetle, Flea Beetle, Green Peach Aphid, Green Stink Bug, Whitefly—20 lbs.

**MUSTARD GREENS** (21), **TURNIP GREENS** (21): Aphids, Cabbage Looper, Imported Cabbage Worm—25 lbs. Do not exceed one application

**USAGE CAUTIONS**

Workers entering treated fields within 24 hours of application should wear protective clothing.

Do not plant root crops other than carrots, potatoes, sugar beets, and sweet potatoes as follow-up crops.

This product is toxic to fish and wildlife. Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes. Birds feeding on treated areas may be killed. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Apply this product only as specified on this label.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

**DISPOSE OF WASTES** by burying in non-crop lands away from water supplies. Container should be disposed of by burying with wastes or by burning. Keep out of smoke.

073-6

MANUFACTURED BY

**DALY-HERRING COMPANY**

KINSTON, NORTH CAROLINA 28501