

STA-GREEN

QUALITY FERTILIZERS SINCE 1924

TREATS 5,000 Sq. Ft.

lawn pest control & fertilizer

FOR CONTROL OF: SOD WEBWORM (LAWN MOTH), ARMYWORMS, ANTS, CHINCH BUGS, BROWN DOG TICKS, FLEAS, CUTWORMS, CHIGGERS, CLOVER MITES, AND CRICKETS.

25-3-3

CONTROLS PESTS AS IT FEEDS GRASS

ACTIVE INGREDIENTS:

Chlorpyrifos [0,0-diethyl 0-(3,5,6-trichloro-2-pyridyl) phosphorothioate].....00.75%

INERT INGREDIENTS:.....99.25%

TOTAL 100.00%

U.S. Patent No. 3,244,589

STATEMENT OF PRACTICAL TREATMENT

CAUTION

KEEP OUT OF REACH OF CHILDREN

IF SWALLOWED, induce vomiting by giving one tablespoon of salt in a glass of warm water.

IF INHALED, remove to fresh air.

IF ON SKIN, Wash with plenty of soap and water.

IF IN EYES, flush with water for 15 minutes.

Get medical attention if irritation persists.

(SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS)

EPA Reg. No. 104-10

EPA Est. No. 8660-AI-1

NET WT. 20 LBS. (9.07 KG)

STA-GREEN PLANT FOOD COMPANY, INC.
CHICAGO, ILL.



lawn pest control & fertilizer

18

RIVERSIDE TOXAPHENE 8

AN EMULSIFIABLE CONCENTRATE CONTAINING 8 POUNDS OF TOXAPHENE PER GALLON. USEFUL FOR THE CONTROL OF CERTAIN INSECT PESTS OF ALFALFA, BEANS, (Dry, Green, Lima), CABBAGE AND OTHER COLE CROPS, CORN, COTTON, COWPEAS, OKRA, PEACHES, PEAS, PEANUTS, PECANS, PIMENTOS, CERTAIN SMALL GRAINS (Barley, Oats, Rice, Rye, Wheat), SORGHUM, SOYBEANS, SPINACH, STRAWBERRIES, AND TOMATOES.

APPROVED
APR 13 1981
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 777-257

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING POISONOUS BY SKIN CONTACT, SWALLOWING, INHALATION OF SPRAY MIST, OR IF ABSORBED THROUGH SKIN

Toxaphene is poisonous by skin contact, inhalation, or swallowing and may be fatal if swallowed. Do not get on skin, in eyes, or on clothing. Toxaphene can be absorbed through the skin. Do not breathe spray mist. Wash thoroughly after using. Wash all contaminated clothing with soap and hot water before re-use. During application keep unprotected persons away from areas being treated or where there may be drift. Do not contaminate feed or foodstuffs.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, shrimp, crab, birds, and other aquatic life. Keep out of any body of water.

Shrimp and crab may be killed at application rates recommended on this label. Do not apply where there are important resources.

Do not apply when weather conditions favor drift from treated areas.

Do not contaminate water by cleaning of equipment or disposal of wastes.

PHYSICAL OR CHEMICAL HAZARDS

DO NOT USE, POUR, OR STORE NEAR HEAT OR OPEN FLAME. In case of spillage, use absorbent clay, hydrated lime, or other solids to soak up any spill. Scrape up excess and dispose away from water's edge. Wash area with a detergent and hot water.

ACTIVE INGREDIENTS:	72%
Toxaphene (Technical chlorinated camphene containing 87.69% toxaphene)	
Inert Range aromatic solvent	28%
WEIGHT INGREDIENTS:	100%
TOTAL:	
Product contains 8 pounds of toxaphene per gallon	

STOP - READ LABEL BEFORE USING!

KEEP OUT OF REACH OF CHILDREN WARNING

STATEMENT OF PRACTICAL TREATMENT

FIRST-AID TREATMENT

INTERNAL: If Riverside Toxaphene 8 is swallowed, do not induce vomiting. See Physicians Note. Never give anything by mouth to an unconscious person. CALL A PHYSICIAN IN ALL CASES OF SUSPECTED POISONING.

EXTERNAL: Remove contaminated clothing and wash skin thoroughly with soap and water. If in eyes, flush with clean water for at least 15 minutes. Call a physician.

PREVENTION: If swallowed, use gastric lavage. Avoid application. Toxaphene produces central nervous system excitation that responds to barbiturate and cholinergic receptor or control compounds with barbiturate, preferably administered intravenously.

If the barbiturate can be given before convulsions start, sodium pentobarbital is the drug of choice. First toxaphene has a long duration of action, a long acting anticholinergic such as phentolamine may be used after initial control of convulsions. Refer to Clinical Handbook on Economic Poisons, which was prepared by the U.S. Department of Health, Education and Welfare, Public Health Service Communicable Disease Center, Atlanta, Georgia.

SEE LEFT SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

NOTICE TO DRUM HANDLERS: IN CASE OF FIRE, LEAKY OR DAMAGED DRUMS, OR OTHER EMERGENCY, REPORT AT ONCE BY TOLL-FREE TELEPHONE TO: 800-424-6266.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

RIVERSIDE TOXAPHENE 8 may be applied by ground sprayers or by airplane using the RIVERSIDE TOXAPHENE 8 emulsion to form an emulsion spray solution. To prepare the spray solution, add the spray tank with water, agitate, add the proper amount of RIVERSIDE TOXAPHENE 8 and then fill the spray tank with water while agitating. Continue to agitate during use.

When applying with ground sprayers, calibrate so as to apply the recommended amount of RIVERSIDE TOXAPHENE 8 in not less than 2 gallons of water per acre. Broadcast spray the recommended amount of RIVERSIDE TOXAPHENE 8 in not less than 1 gallon of water per acre. Always use sufficient water for adequate coverage of the crop foliage.

The amount of spray solution applied should not exceed the rate per acre of RIVERSIDE TOXAPHENE 8 in any one application. When a single crop is grown on the same soil or young plants or with light soil conditions, use the higher rate on top or bottom plants or soil heavy sand conditions.

The labels and directions for handling RIVERSIDE TOXAPHENE 8 in the container are applicable to all other dilutions with water for spray application.

ALFALFA: Broadcast granules with 40% Calcium Cyanide 2 1/2 to 3 to 4 lbs. per acre. Broadcast granules 2 to 3 to 4 lbs. per acre. Broadcast granules 2 to 3 to 4 lbs. per acre. Broadcast granules 2 to 3 to 4 lbs. per acre. Broadcast granules 2 to 3 to 4 lbs. per acre.

BEANS (Dry, Green, Lima): For control of Mexican Bean Beetle and Thrips apply 1 to 2 to 3 to 4 lbs. per acre. Broadcast granules 2 to 3 to 4 lbs. per acre. Broadcast granules 2 to 3 to 4 lbs. per acre. Broadcast granules 2 to 3 to 4 lbs. per acre.

BROAD LEAF GRASSES: Broadcast granules 2 to 3 to 4 lbs. per acre. Broadcast granules 2 to 3 to 4 lbs. per acre. Broadcast granules 2 to 3 to 4 lbs. per acre. Broadcast granules 2 to 3 to 4 lbs. per acre.

CABBAGE AND OTHER COLE CROPS: Broadcast granules 2 to 3 to 4 lbs. per acre. Broadcast granules 2 to 3 to 4 lbs. per acre. Broadcast granules 2 to 3 to 4 lbs. per acre. Broadcast granules 2 to 3 to 4 lbs. per acre.

CORN: Broadcast granules 2 to 3 to 4 lbs. per acre. Broadcast granules 2 to 3 to 4 lbs. per acre. Broadcast granules 2 to 3 to 4 lbs. per acre. Broadcast granules 2 to 3 to 4 lbs. per acre.

Do not use this product on or within 30 days of harvest. Plants treated with RIVERSIDE TOXAPHENE 8 should not be fed to dairy animals or animals being bred for slaughter.

CORN: For control of Corn Earworm and Corn Rootworm apply 2 to 3 to 4 lbs. per acre. Broadcast granules 2 to 3 to 4 lbs. per acre. Broadcast granules 2 to 3 to 4 lbs. per acre. Broadcast granules 2 to 3 to 4 lbs. per acre.

COTTON: For control of Cotton Bollworm, Cotton Rootworm, and Cotton Leafworm apply 2 to 3 to 4 lbs. per acre. Broadcast granules 2 to 3 to 4 lbs. per acre. Broadcast granules 2 to 3 to 4 lbs. per acre. Broadcast granules 2 to 3 to 4 lbs. per acre.

COWPEAS: For control of Cowpea Curculionid apply 2 to 3 to 4 lbs. per acre. Broadcast granules 2 to 3 to 4 lbs. per acre. Broadcast granules 2 to 3 to 4 lbs. per acre. Broadcast granules 2 to 3 to 4 lbs. per acre.

CUCUMBERS: For control of Cucumber Beetle apply 2 to 3 to 4 lbs. per acre. Broadcast granules 2 to 3 to 4 lbs. per acre. Broadcast granules 2 to 3 to 4 lbs. per acre. Broadcast granules 2 to 3 to 4 lbs. per acre.

OKRA: For control of Okra Weevil apply 2 to 3 to 4 lbs. per acre. Broadcast granules 2 to 3 to 4 lbs. per acre. Broadcast granules 2 to 3 to 4 lbs. per acre. Broadcast granules 2 to 3 to 4 lbs. per acre.

PEACHES: For control of Peach Borer apply 2 to 3 to 4 lbs. per acre. Broadcast granules 2 to 3 to 4 lbs. per acre. Broadcast granules 2 to 3 to 4 lbs. per acre. Broadcast granules 2 to 3 to 4 lbs. per acre.

PEAS: For control of Pea Aphid apply 2 to 3 to 4 lbs. per acre. Broadcast granules 2 to 3 to 4 lbs. per acre. Broadcast granules 2 to 3 to 4 lbs. per acre. Broadcast granules 2 to 3 to 4 lbs. per acre.

PECANS: For control of Pecan Weevil apply 2 to 3 to 4 lbs. per acre. Broadcast granules 2 to 3 to 4 lbs. per acre. Broadcast granules 2 to 3 to 4 lbs. per acre. Broadcast granules 2 to 3 to 4 lbs. per acre.

PEANUTS: For control of Peanut Rootworm apply 2 to 3 to 4 lbs. per acre. Broadcast granules 2 to 3 to 4 lbs. per acre. Broadcast granules 2 to 3 to 4 lbs. per acre. Broadcast granules 2 to 3 to 4 lbs. per acre.

Do not apply more than 2 days before harvest. Do not feed treated plants or animals from treated plants to poultry, dairy animals, or animals being bred for slaughter. Do not feed treated plants to livestock or poultry.

PEANUTS: For control of Peanut Rootworm apply 2 to 3 to 4 lbs. per acre. Broadcast granules 2 to 3 to 4 lbs. per acre. Broadcast granules 2 to 3 to 4 lbs. per acre. Broadcast granules 2 to 3 to 4 lbs. per acre.

PEACHES: For control of Peach Borer apply 2 to 3 to 4 lbs. per acre. Broadcast granules 2 to 3 to 4 lbs. per acre. Broadcast granules 2 to 3 to 4 lbs. per acre. Broadcast granules 2 to 3 to 4 lbs. per acre.

PEACHES: For control of Peach Borer apply 2 to 3 to 4 lbs. per acre. Broadcast granules 2 to 3 to 4 lbs. per acre. Broadcast granules 2 to 3 to 4 lbs. per acre. Broadcast granules 2 to 3 to 4 lbs. per acre.

PEACHES: For control of Peach Borer apply 2 to 3 to 4 lbs. per acre. Broadcast granules 2 to 3 to 4 lbs. per acre. Broadcast granules 2 to 3 to 4 lbs. per acre. Broadcast granules 2 to 3 to 4 lbs. per acre.

PEACHES: For control of Peach Borer apply 2 to 3 to 4 lbs. per acre. Broadcast granules 2 to 3 to 4 lbs. per acre. Broadcast granules 2 to 3 to 4 lbs. per acre. Broadcast granules 2 to 3 to 4 lbs. per acre.

PEACHES: For control of Peach Borer apply 2 to 3 to 4 lbs. per acre. Broadcast granules 2 to 3 to 4 lbs. per acre. Broadcast granules 2 to 3 to 4 lbs. per acre. Broadcast granules 2 to 3 to 4 lbs. per acre.

PEACHES: For control of Peach Borer apply 2 to 3 to 4 lbs. per acre. Broadcast granules 2 to 3 to 4 lbs. per acre. Broadcast granules 2 to 3 to 4 lbs. per acre. Broadcast granules 2 to 3 to 4 lbs. per acre.

PEACHES: For control of Peach Borer apply 2 to 3 to 4 lbs. per acre. Broadcast granules 2 to 3 to 4 lbs. per acre. Broadcast granules 2 to 3 to 4 lbs. per acre. Broadcast granules 2 to 3 to 4 lbs. per acre.

PEACHES: For control of Peach Borer apply 2 to 3 to 4 lbs. per acre. Broadcast granules 2 to 3 to 4 lbs. per acre. Broadcast granules 2 to 3 to 4 lbs. per acre. Broadcast granules 2 to 3 to 4 lbs. per acre.

h
18

