

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

Alice C. Walker  
 Manager, Label Development  
 Riverside/Terra Corporation  
 P.O. Box 171376  
 Memphis, TN 38187

Dear Ms. Walker:


Subject: Atrazine 4L; EPA Reg. No. 9779-255  
 Atrazine 80W; EPA Reg. No. ~~9779-254~~  
 Atrazine 90DF; EPA Reg. No. 9779-253  
 Re: Atrazine Voluntary Exposure Reductions Measures  
 Your Amendment Submissions Dated May 21, 1990

The labeling referred to above, submitted in connection with the Federal Insecticide, Fungicide, and Rodenticide Act, as amended is acceptable with the following comments:

1. On Page 10 (all 3 labels), delete references to "Boot Stage" or "3 Leaf Stage", or add the statement, "but no higher than 12 inches".
2. On Page 2 (all 3 labels), under "Statement of Practical Treatment, If On Skin", Add the word "skin" to read: "Remove contaminated clothing and wash skin with soap and water. Also add the statement: "Get medical attention if irritation persists".
3. Delete the pineapple use from the 80W label as discussed in Beverley Neal's phone conversation with Steve Morrill.

Please submit five (5) copies each of your final printed labeling incorporating this amendment and above mentioned comments before you release these products for shipment. All technical and end-use products released for shipment must have amended labels after September 1, 1990.

Sincerely yours,



Robert J. Taylor  
 Product Manager (25)  
 Fungicide-Herbicide Branch  
 Registration Division (H7505C)

CONCURRENCES

SYMBOL	H7505C							
SURNAME	Morrill							
DATE	5/29/90							

RESTRICTED USE PESTICIDE  
(GROUND WATER CONCERNS)

FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR  
PERSONS UNDER THEIR DIRECT SUPERVISION, AND ONLY FOR THOSE USES  
COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION.

THIS PRODUCT IS A RESTRICTED USE HERBICIDE DUE TO GROUND WATER  
CONCERNS. USERS MUST READ AND FOLLOW ALL PRECAUTIONARY STATEMENTS  
AND INSTRUCTIONS FOR USE IN ORDER TO MINIMIZE POTENTIAL FOR  
ATRAZINE TO REACH GROUND WATER.

R®

ATRAZINE 80W

For weed control in corn, sorghum, and other crops

ACCEPTED  
with COMMENTS  
in EPA Letter Dated:

1988 20

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.  
9779-254

ACTIVE INGREDIENTS

Atrazine (2-chloro-4-ethylamino- 6-isopropylamino-s-triazine).....	76.8%
Related Compounds.....	3.2%
INERT INGREDIENTS.....	20.0%
TOTAL	100.0%

KEEP OUT OF REACH OF CHILDREN  
CAUTION  
STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Contact your local poison control center, hospital, or physician.  
If the patient is unconscious, maintain breathing and heartbeat (cardiopulmonary  
resuscitation). If the patient is conscious and alert, induce vomiting (syrup  
of ipecac or stimulate the back of the throat with a finger). Never give  
anything by mouth to an unconscious person.  
IF INHALED: Remove victim to fresh air and apply respiration if indicated.  
IF ON SKIN: Remove contaminated clothing and wash with soap and water.  
IF IN EYES: Flush with plenty of water. Get medical attention if irritation  
persists.

See additional PRECAUTIONARY STATEMENTS on side panel.

EPA Reg. No. 9779-254

EPA Est. No. 9779-

Manufactured For  
RIVERSIDE/TERRA CORPORATION  
Terra Centre, 600 Fourth Street, Sioux City, Iowa 51101  
Riverside Serves Agriculture. Agriculture Serves Everyone.

NET CONTENTS  
LBS.

BEST AVAILABLE COPY

---

**PRECAUTIONARY STATEMENTS**  
**CAUTION**  
**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

Harmful if swallowed, inhaled, or absorbed through skin. Do not breathe dust or spray mist. Avoid contact with eyes, skin or clothing. Remove contaminated clothing and wash before reuse. Wash thoroughly with soap and water after handling.

Users are required to wear long sleeve shirts and long pants or equivalent, chemical resistant gloves, and boots (waterproofed). In addition, persons involved in mixing/loading operations are required to use chemical resistant rubber or neoprene gloves and a face shield or goggles. Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

**NOTE TO PHYSICIANS:** There is no specific antidote for atrazine. If this product is ingested, induce emesis or lavage stomach. The use of an aqueous slurry of activated charcoal may be considered.

**ENVIRONMENTAL HAZARDS**

This pesticide is toxic to aquatic invertebrates. Do not apply directly to water or wetlands. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters.

Atrazine leaches readily and accepted label rates have been found to result in contamination of water supplies by way of groundwater. Therefore users are advised to avoid use of atrazine in well drained soils, particularly in areas having high groundwater tables. Ground water contamination may be reduced by diking and flooring of permanent liquid bulk storage sites with an impermeable material. This product may not be mixed/loaded, or used within 50 feet of all wells including abandoned wells, drainage wells, and sink holes.

---

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product through any type of irrigation system.

**RE-ENTRY STATEMENT:** Do not enter treated areas without protective clothing until sprays have dried. Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

**NOTICE TO CROP OWNERS**

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. In case of accidental exposure follow the information given under STATEMENT OF PRACTICAL TREATMENT and have exposed person(s) see a physician. When oral warnings are

**BEST AVAILABLE COPY**

given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is a reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information: CAUTION. Area treated with Atrazine on (date). Do not enter without appropriate protective clothing until sprays have dried. In case of accidental exposure, refer to the STATEMENT OF PRACTICAL TREATMENT on the label.

---

#### STORAGE AND DISPOSAL

DO NOT CONTAMINATE WATER, FOOD, OR FEED BY STORAGE OR DISPOSAL.

##### STORAGE

Store in a dry location away from children, animals, foods, feeds, seeds, or other agricultural chemicals. Handle in accordance with information given under PRECAUTIONARY STATEMENTS. In the event of spillage, scrape up spillage. If spillage is not suitable for use, dispose of in accordance with information given under DISPOSAL. Repackage and relabel useable product in a sound container. In case of fire or other emergency, report at once by toll-free telephone to 800-424-9300.

##### DISPOSAL

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

---

#### GENERAL INFORMATION

This herbicide controls many annual broadleaf and grass weeds in corn, sorghum, sugarcane, and certain other crops specified on this label. It is also effective in noncrop areas and industrial sites for control of most annual and many perennial broadleaf and grass weeds. This product may be applied before or after weeds emerge.

FOLLOWING MANY YEARS OF CONTINUOUS USE OF THIS PRODUCT AND CHEMICALLY RELATED PRODUCTS, BIOTYPES OF SOME OF THE WEEDS LISTED ON THIS LABEL HAVE BEEN REPORTED WHICH CANNOT BE EFFECTIVELY CONTROLLED BY THIS AND RELATED HERBICIDES. WHERE THIS IS KNOWN OR SUSPECTED, WE RECOMMEND THE USE OF THIS PRODUCT IN COMBINATION WITH OTHER REGISTERED HERBICIDES WHICH ARE NOT TRIAZINES. CONSULT WITH YOUR STATE AGRICULTURAL EXTENSION SERVICE FOR SPECIFIC RECOMMENDATIONS.

Since this product acts mainly through root absorption, its effectiveness depends on moisture to move it into the root zone. If weeds develop, a shallow cultivation or rotary hoeing will generally result in better weed control.

This product is noncorrosive to equipment and metal surfaces, non-flammable, and has low electrical conductivity.

Avoid using near adjacent desirable plants or in greenhouses, or injury may occur.

Where the use directions give a range of rates, use the lower rate on coarse-textured soil and soil high in organic matter.

Note: Riverside/Terra Corporation does not recommend applications in combination with other herbicides or oils, except as specifically described on the label.

**BEST AVAILABLE COPY**

## APPLICATION PROCEDURES

**Ground Application:** Use conventional ground sprayers equipped with nozzles that provide accurate and uniform application. Be certain that nozzles are uniformly spaced and are the same size. Calibrate sprayer before use and recalibrate at the start of each season and when changing carriers. Unless otherwise specified, use a minimum of 10 gallons of spray mixture per acre for all preplant, preemergence, and postemergence applications (with or without oil or surfactant) with ground equipment.

Use a pump with a capacity to (1) maintain 35-40 psi at nozzles, (2) provide sufficient agitation in tank to keep mixture in suspension, and (3) to provide a minimum of 20% bypass at all times. Use centrifugal pumps which provide propeller shear action for dispersing and mixing this product. The pump should provide a minimum of 10 gallons per minute per 100 gallons tank size circulated through a correctly positioned sparger tube or jets.

Use screens to protect the pump and to prevent nozzles from clogging. Screens placed on the suction side of the pump should be 16-mesh or coarser. Do not place a screen in the recirculation line. Use 50-mesh or coarser screens between the pump and boom, and where required, at the nozzles. Check nozzle manufacturer's recommendations.

For band applications, calculate amount to be applied per acre as follows:

$\frac{\text{band width in inches}}{\text{row width in inches}}$	x	broadcast rate	=	amount needed
		per acre		per acre of field

**Aerial Application:** Use aerial application only where broadcast applications are specified. Apply in a minimum of 1 gallon of water for each pound of ATRAZINE 80W applied per acre. For postemergence treatments on corn and sorghum, apply recommended rate in a minimum of 2 gallons of water per acre. Avoid applications under conditions where uniform coverage cannot be obtained or where excessive spray drift may occur.

Avoid application to humans or animals. Flagmen and loaders should avoid inhalation of spray mist and prolonged contact with skin and should wash thoroughly before eating and at the end of each day's operation.

**Application in Water or Liquid Fertilizer:** Nitrogen solution or complete liquid fertilizer may replace all or part of the water as a carrier for preemergence or preplant ground application on corn and sorghum. Before using, check the compatibility of this product with liquid fertilizer and/or nitrogen solution as shown below. Do not apply in nitrogen solution or complete liquid fertilizer after corn or sorghum emerges, except as noted under Lay-by Treatment for corn, or crop injury may occur.

To determine the compatibility of ATRAZINE 80W alone or in tank mixtures with fluid fertilizer, pour the products into a small container of fluid fertilizer in the proportions shown below. For different spray volume or herbicide rate, make appropriate changes in the ingredients.

For each pound of ATRAZINE 80W or other liquid or flowable product to be applied per acre, add 1 level teaspoon or 5 milliliters of the product to one pint of fluid fertilizer (assuming a volume of 25 gallons per acre).

For each pint of a liquid or flowable product to be applied per acre, add 0.5 teaspoon or 2.5 milliliters of the product to one pint of fluid fertilizer (assuming a volume of 25 gallons per acre).

**BEST AVAILABLE COPY**

After thorough mixing, let stand for 5 minutes. If the herbicide/fertilizer combination remains mixed, or can be remixed readily, the mixture is compatible and can be sprayed. If it is not compatible, make a water slurry with the herbicide before adding the fertilizer and use the mixture only if this test shows it is compatible.

Application in Water Plus Emulsifiable Oil or Oil Concentrate: Adding an emulsifiable oil or oil concentrate, such as Prime Oil or Prime Oil II, to postemergence water-based sprays on corn and sorghum may improve weed control. However, under certain conditions, the use of either type of oil may seriously injure the crop. To minimize this possibility, observe the following directions.

Use a crop oil containing 1-2% suitable emulsifier, or a suitable crop oil concentrate containing not more than 20% emulsifier or surfactant blend. Emulsifiable oil and oil concentrate contaminated with water or other materials can cause compatibility problems and/or crop injury.

Mixing Procedures--All Uses: (1) Be sure sprayer is clean and not contaminated with any other materials or crop injury or sprayer clogging may result. (2) Fill tank 1/4 full with clean water, nitrogen solution, or complete liquid fertilizer. (3) Start agitation. (4) Be certain that the agitation system is working properly and creates a rippling or rolling action on the liquid surface. (5) Pour product directly from container into tank. (6) Continue filling tank until 90% full. Increase agitation if necessary to maintain surface action. (7) Add emulsifiable oil, oil concentrate or tank mix herbicide(s) after this product is thoroughly suspended. (8) Finish filling tank. (9) Empty tank as completely as possible before refilling to prevent buildup of oil or emulsifiable concentrate residue. Maintain agitation to avoid separation of materials. (10) If an oil or emulsifiable concentrate film starts to build up in tank, drain it and clean with strong detergent solution or solvent. (11) Clean sprayer thoroughly immediately after use by flushing system with water containing a detergent.

ATRAZINE 80W applied alone--Corn or Grain Sorghum

Preplant Incorporated or Preemergence Weeds Controlled:

annual morningglory	mustards
barnyardgrass	nightshade
(watergrass)	pigweed
cocklebur	purslane
giant foxtail*	ragweed
green foxtail*	velvetleaf
groundcherry	(buttonweed)
jimsonweed	wild oats
kochia	witch grass
lambsquarters	(Panicum capillare)
large (nairy) crabgrass	yellow foxtail*

\*Control of foxtail can be erratic

Postemergence with Emulsifiable Oil or Oil Concentrate in Water--Weeds Controlled:

annual morningglory	pigweed
cocklebur	ragweed
jimsonweed	smartweed
lambsquarters	wild buckwheat
mustards	velvetleaf

**BEST AVAILABLE COPY**

**CORN**

**Preplant (Broadleaf and grass control):** Broadcast in spring after plowing at rate in Table 1. Apply before, during, or after final seedbed preparation. If soil is tilled or worked after application, avoid deep incorporation. For best results, apply within two weeks prior to planting.

**Preemergence (Broadleaf and grass control):** Apply during or shortly after planting before weed emergence at rate in Table 1.

**Postemergence (Broadleaf and grass control):** Apply before weeds EXCEED 1.5 inches in height at rate in Table 1. Postemergence application to corn must be made before corn reaches 12 inches in height. The maximum application rate for corn is 3.75 lbs. per acre per calendar year. Applications for quackgrass suppression in corn are restricted to a spring application only. No fall applications are permitted.

**TABLE 1: Broadleaf and Grass Weed Control on Corn\***

Soil Texture	Broadcast Rate/A
<b>COARSE</b>	
Sand, loamy sand, sandy loam	2.5 lb.
<b>MEDIUM</b>	
Loam, silt loam, silt, clay loam, sandy clay loam, silty clay loam, sandy clay, or silty clay with low organic matter	3.0 lb.
<b>FINE</b>	
Loam, silt loam, silt, clay loam, sandy clay loam, silty clay loam, sandy clay, or silty clay with medium to high organic matter, and clay (including the dark prairie soils of the Corn Belt)	3.75 lb.
Peat, muck and high organic clay (apply postemergence only)	3.75 lb.

**BEST AVAILABLE COPY**

\*Broadleaf control (eastern CO, western KS, western NE, NM, OK Panhandle, west TX and eastern WY). On sand, sandy loam, mild to strongly alkaline soil, and all recently leveled soil, apply no more than 1.5 lb/A, either preplant or preemergence. On the other soils in these areas, apply rate in Table 1 for broadleaf and grass control.

**Postemergence with Emulsifiable Oil or Oil Concentrate in Water:** Add 1 gal. of emulsifiable oil per acre for ground application, or 0.5 gal. per acre for aerial application, or add 1 qt. of oil concentrate, such as Prime Oil or Prime Oil II, per acre for ground application, or 1 pt. per acre for aerial application.

**Broadleaf and Grass Control:** Broadcast 1.5 lb. per acre plus emulsifiable oil or oil concentrate before pigweed and lambsquarters reach 6 inches in height and before all other weeds reach 4 inches in height. A cultivation may be necessary if all weeds are not controlled or if weeds regrow.

**Precautions for Applications with Emulsifiable Oil or Oil Concentrate:**

- (1) Do not apply when crop is under stress from prolonged cold, wet weather, poor fertility, or other factors, or when crop is wet and succulent from recent rainfall, as crop injury may occur.
- (2) Inbred lines or any breeding stock may be severely injured by applications with emulsifiable oil or oil concentrate.
- (3) Adding other insecticides, herbicides, liquid fertilizers, or other materials is not recommended because they may cause compatibility problems or crop injury.
- (4) Store and handle emulsifiable oil and oil concentrate carefully. Oil contaminated with even a small amount of water may not emulsify properly when added to the tank.
- (5) Do not make more than one application per season except as recommended for suppression of yellow nutsedge and Canada thistle.

## Problem Weeds:

**Yellow nutsedge and Canada thistle:** This product will partially control (suppress) yellow nutsedge (*Cyperus esculentus*) and Canada thistle (*Cirsium arvense*) when applied according to directions. For best results, apply each year until yellow nutsedge or Canada thistle is eliminated or reaches a level of infestation where neither weed is an economic problem. If yellow nutsedge or Canada thistle regrow following last application, cultivate once.

Four alternative methods of use for partial control (suppression) of yellow nutsedge and Canada thistle are listed below in order of preference. If other weed species, such as annual grasses, are also expected, use alternative 2 or 4.

1. Broadcast 1.8 lb/A plus 1 gal. of emulsifiable oil or 1 qt. of oil concentrate. Apply after crop and yellow nutsedge or Canada thistle emerge but before yellow nutsedge reaches a height of 3 inches or Canada thistle reaches a height of 6 inches. Repeat application 10-20 days later, but before corn is 12 inches tall.
2. Broadcast 1.8 lb/A preplant incorporated for control of yellow nutsedge, or broadcast 1.8 lb/A during or shortly after planting, but prior to crop or weed emergence for suppression of Canada thistle. Follow with an application of 1.8 lb/A plus 1 gal. of emulsifiable oil or 1 qt. of oil concentrate after corn and weeds emerge, but before yellow nutsedge or Canada thistle reaches height of 6 inches and before the corn is 12 inches high.
3. Broadcast 3.75 lb/A plus 1 gal. of emulsifiable oil or 1 qt. of oil concentrate after crop emerges, but before corn is 12 inches high. Apply after yellow nutsedge and Canada thistle emerge, but before yellow nutsedge is 3 inches tall or Canada thistle is 6 inches tall.
4. Broadcast 3.75 lb/A before, during, or shortly after planting, but before crop or weed emergence (yellow nutsedge only).

**Note:** Do not use emulsifiable oil or oil concentrate when corn is wet or under stress, especially when using 3.75 lb/A of this product. See Precautions for Applications with Emulsifiable Oil or Oil Concentrate in Water for additional directions.

**QUACKGRASS SUPPRESSION ON LAND GOING INTO CORN PRODUCTION:** Broadcast 3.75 lb/A in the spring before corn reaches a height of 12 inches. Plow 1-3 weeks later. Follow up in the fall using a suitable contact herbicide.

**BEST AVAILABLE COPY**

## TANK MIXTURES FOR CORN

This product may be tank mixed with these herbicides for control of certain broadleaf and grass weeds in corn:

Dual* (metolachlor)	Lasso* or Lasso EC (alachlor)
Dual + Gramoxone Super*	Lasso or Lasso EC + Roundup
Dual + Roundup* (glyphosate)	Lasso or Lasso EC + Gramoxone Super
Dual + Princep*	Gramoxone Super
Dual + Princep + Gramoxone Super	Princep
Dual + Princep + Roundup	Princep + Gramoxone Super
Bexton* or Ramrod*	Princep + Roundup
Bladex*	Sutan **

\*Use tank mix directions appearing on the labels of the above herbicides when tank mixing with this product. Observe all precautions and limitations on labeling of products used in a particular tank mix.

**Note:** When labels of the above herbicides refer to Atrazine 4L or Atrazine 90 dry flowable, use equivalent rates of ATRAZINE 80W. One quart of 4L equals 1.25 lb. of 80W. One lb. of 90 dry flowable equals 1.125 lb. of 80W.



Princep 80W, Princep 4L (4LC), or Princep Caliber\* 90

In addition to the weeds listed under ATRAZINE 80W Applied Alone--Corn and Grain Sorghum--Preplant or Preemergence, this combination controls crabgrass, fall panicum, and carpetweed.

Broadcast tank mix before planting, at planting, or after planting, but before weeds emerge, at rates in Table 2. Use the 1:1 ratio for control of most weeds. Use the 1:2 ratio for expected heavy infestations of crabgrass and fall panicum. Cultivate shallowly if weeds develop.

Preplant: Apply in spring during or after final seedbed preparation. If soil is tilled or worked after application, avoid deep incorporation. For best results, apply within two weeks prior to planting.

Preemergence: Apply during or shortly after planting, but before crop and weeds emerge.

Refer to CORN sections of this label and to Princep 80W, Princep Caliber 90, or Princep 4L (4LC) labels for further directions, limitations, and precautions.

Table 2 Tank Mixtures with Princep on Corn

Soil Texture	BROADCAST RATE PER ACRE			
	1:1 Ratio*		1:2 Ratio**	
	This Product	Princep 80W <sup>1</sup>	This Product	Princep 80W <sup>1</sup>
Sand, loamy sand, sandy loam	1.25 lb.	1.25 lb.	0.83 lb.	1.67 lb.
Loam, silt loam, silt, clay loam, sandy clay loam, silty clay loam, sandy clay, or silty clay with low organic matter	1.5 lb.	1.5 lb.	1 lb.	2 lb.
Loam, silt loam, silt, clay loam, sandy clay loam, silty clay loam, sandy clay, or silty clay with medium to high organic matter, and clay (including dark prairie soils of the Corn Belt)	1.8 lb.	1.8 lb.	1.2 lb.	2.4 lb.

\*For control of above weeds.

\*\*For control of expected heavy infestations of crabgrass and fall panicum.

<sup>1</sup>When using Princep Caliber 90 or Princep 4L(4LC), use equivalent rates. One lb. of Princep 80W equals 0.9 lb. of Princep Caliber 90 or 1.6 pt. of Princep 4L(4LC).

Princep 80W, Princep 4L (4LC), or Princep Caliber 90 plus Roundup: Use as a tank mixture for preemergence and postemergence control of certain broadleaf and grass weeds where corn will be planted directly into a cover crop, established sod, or in previous crop residues. Refer to Roundup label for all directions, weeds controlled, precautions, and limitations.

Princep 80W, Princep 4L (4LC), or Princep Caliber 90 plus Gramoxone Super: Use as a tank mixture with Princep and Gramoxone Super to kill existing vegetation and for residual weed control where corn will be planted directly into a cover crop, established sod, or in previous crop residues. Add this product and Princep to water in spray tank, agitating until thoroughly mixed. Then add Gramoxone Super\* and a nonionic surfactant, such as Activate Plus. Continue agitation during application. Broadcast 1.25-2.5 lb. of this product plus

**BEST AVAILABLE COPY**

1.25-2.5 lb. Princep 80W (or 2-4 pt. Princep 4L(4LC), or 1.1-2.2 lb. Princep Caliber 90) plus 1-2 pt. Gramoxone Super in 20-60 gallons of water per acre. Apply before, during, or after planting, but before corn emerges. Add 0.5 pt. of a nonionic surfactant, such as Activate Plus, per 100 gallons of spray mixture. Use the 2 pint rate of Gramoxone Super if existing vegetation is 4-6 inches tall. This mixture will not control weeds taller than 6 inches.

Refer to further limitations and precautions on labels for this product, Princep, and Gramoxone Super.

Precautions for all applications to corn: (1) The maximum application rate for corn is 3.75 lbs. per acre per calendar year. Applications for quackgrass suppression in corn are restricted to a spring application only. No fall applications are permitted. (2) Following harvest, plow (moldboard or disk-plow) and thoroughly till soil in fall or spring to minimize possible injury to rotational spring-seeded crops, regardless of rate used.

Note for all Applications to Corn: Do not graze or feed forage from treated areas for 21 days following application.

ROTATIONAL CROPS: 1) Do not rotate to any crop except corn or sorghum until the following year, or injury may occur. 2) If applied after June 10, do not rotate with crops other than corn or sorghum the next year, or crop injury may occur. 3) In the High Plains and Intermountain areas of the West where rainfall is sparse and erratic or where irrigation is required, use only when corn or sorghum is to follow corn or sorghum or when a crop of untreated corn or sorghum is to precede other rotational crops. 4) In eastern parts of the Dakotas, KS, western MN, and NE., do not rotate to soybeans if the rate applied to corn or sorghum was more than 2.5 lb/A or equivalent band application rate, or soybean injury may occur. 5) Injury may occur to soybeans planted the year following application on soils having a calcareous surface layer. 6) Do not plant sugar beets, tobacco, vegetables (including dry beans), spring-seeded small grains, or small-seeded legumes and grasses the year following application, or injury may occur.

**SORGHUM AND SORGHUM-SUDAN HYBRIDS**  
(Grain and Forage Types)

**BEST AVAILABLE COPY**

Preplant (Broadleaf and grass control): Broadcast in spring after plowing at rate in Table 3. Apply before, during, or after final seedbed preparation. If soil is tilled or worked after application, avoid deep incorporation. For best results, apply within two weeks prior to planting.

Preemergence (Broadleaf and grass control): Apply during or shortly after planting, but prior to weed or crop emergence at rate in Table 3.

Table 3: Preplant and Preemergence Broadleaf and Grass Weed Control in Sorghum\*

SOIL TEXTURE	ORGANIC MATTER	BROADCAST RATE PER ACRE
COARSE Sand, loamy sand, sandy loam	Any level	DO NOT USE (except for preemergence use on bedded sorghum in AZ and CA)
MEDIUM AND FINE Loam, silt loam, silt, clay loam, sandy clay loam, silty clay loam, sandy clay, silty clay, clay	Less than 1%	
	1-1.5%	2-2.5 lb.
	More than 1.5%	2.5-3 lb.

\*Do not apply preplant in AL, AR, FL, GA, LA, MS, NC, NM, OK, SC, TN or TX. Do not apply preemergence in NM, OK, or TX, except in northeast OK, and the TX Gulf Coast.

In case of planting failure, sorghum or corn may be replanted. Do not make a second broadcast application, or injury may occur. If originally applied in a band and sorghum or corn is replanted in untreated row middles, this product may be applied in a band to the second planting. However, the maximum application rate for sorghum is 3.75 lb/A per calendar year.

Preemergence broadleaf weed control in furrow irrigated bedded sorghum (AZ and CA only): For preemergence control of many broadleaf weeds, broadcast 1-1.5 lb. per acre after bed preparation during or after planting, but before sorghum and weeds emerge and before the first furrow irrigation. Follow with several regular irrigations, making sure to thoroughly wet all soil.

Precautions for Preemergence Application to Furrow Irrigated Bedded Sorghum in AZ and CA: To avoid possible sorghum injury, do not use on sand or loamy sand soil or on sorghum planted in furrows. Applications to sorghum growing on alkali soils or where cuts, fills, or erosion have exposed calcareous or alkali subsoils may cause crop injury. In case of crop failure, do not replant sorghum for 8 months following application. Corn may be planted immediately.

Postemergence (Broadleaf and grass control): Apply at rate in Table 4 before weeds exceed 1.5 inches in height. Postemergence application to sorghum must be made before sorghum reaches 12 inches in height. The maximum application rate for sorghum is 3.75 lbs. per acre per calendar year.

TABLE 4: Postemergence Broadleaf and Grass Weed Control in Sorghum

Soil Texture	Minimum Sorghum Height at Treatment	Broadcast Rate Per Acre
Sand or loamy sand	DO NOT USE	
Sandy loam	See Directions For Postemergence Broadleaf Weed Control	
Silt loam to sandy clay loam	Completely Emerged	2.5-3 lb.
Olton and Pullman clay soil	6 inches	2.5-3 lb.
Silty clay loam and finer textured soil	Completely emerged	3.75 lb.

Postemergence Broadleaf Weed Control with Emulsifiable Oil or Oil Concentrate in Water: Broadcast 1.5 lb/A for control of many broadleaf weeds. Apply before pigweed and lambsquarters reach 6 inches in height and before all other weeds reach 4 inches in height. In CO, western KS, NM, OK, TX and desert regions of AZ and CA, apply when sorghum is 6-10 inches in height, but before it reaches boot stage. In all other areas, apply after sorghum reaches 3-leaf stage. Add 1 gal. of emulsifiable oil per acre for ground application and 0.5 gal. per acre for aerial application, or add 1 qt. per acre of oil concentrate, such as Prime Oil or Prime Oil II, for ground application, or 1 pt/A for aerial application. A cultivation may be necessary if all weeds are not controlled or if weeds regrow.

For a list of weeds controlled, see ATRAZINE 80W Applied Alone--Corn or Grain Sorghum--Postemergence with Emulsifiable Oil or Oil Concentrate in Water.

Precautions for Applications with Emulsifiable Oil or Oil Concentrate in Water: See Precautions for Applications with Emulsifiable Oil or Oil Concentrate in Water in Corn section.

Postemergence Broadleaf Weed Control with Surfactant (CO, western KS, NM, OK, TX, and desert regions of AZ and CA only): Broadcast 1.5 lb/A plus 0.75-1.5 pt. of surfactant after sorghum reaches 6 inches in height, but before weeds exceed 1.5 inches in height. Apply only on sandy loam and finer textured soil.

---

**Precautions for All Applications to Sorghum:** (1) Heavy rain immediately following application tends to cause excessive concentration of herbicide in seed furrow, resulting in possible crop injury. Do not apply to furrow-planted sorghum until furrows are leveled (plowed in). Level deep planter marks or seed furrows before application. (2) Application to sorghum growing under stress caused by minor element deficiency or to sorghum growing on highly calcareous soil may result in crop injury. (3) Following harvest, plow (moldboard or disk-plow) and thoroughly till soil in fall or spring to minimize possible injury to spring-seeded rotational crops, regardless of rate used. (4) Injury may occur if both this herbicide, preplant or preemergence, and an at-planting systemic insecticide applied in-furrow are used. (5) The maximum application rate for sorghum is 3.75 lbs. per acre per calendar year. Applications for quackgrass suppression in sorghum are restricted to a spring application only. No fall applications are permitted.

---

**Note:** Do not graze or feed forage from treated areas for 21 days following application.

**Rotational Crops:** See "Rotational Crops" in Corn section

#### TANK MIXTURES FOR GRAIN SORGHUM

**Igran\* 80W:** Use as tank mixture for control of certain broadleaf and grass weeds in grain sorghum. Refer to Igran 80W label for all directions, weeds controlled, precautions, and limitations.

**Dual 8E:** Use as tank mixture with Dual 8E for control of those weeds listed on the Dual label, as well as on this label. Use this tank mixture only on Concep\*-treated sorghum seed. Refer to the Dual 8E label for all directions, precautions, and limitations.

#### ATRAZINE 80W Alone--Chemical Fallow

**Wheat-Sorghum-Fallow:** To control annual broadleaf and grass weeds following wheat harvest and in the following sorghum crop when grown under minimum tillage, broadcast 3.75 lb/A to wheat stubble immediately following wheat harvest. If weeds are present at application, remove them with a sweep plow or other suitable implement after application, or use an approved contact herbicide before or after the application of ATRAZINE 80W. Plant sorghum into wheat stubble the following spring with minimum soil disturbance. Use a surface planter or a planter leaving a shallow furrow. If weeds are present at planting, remove them with a sweep plow or other suitable implement before planting.

For a list of weeds controlled, see ATRAZINE 80W Applied Alone--Corn or Grain Sorghum--Preplant Incorporated or Preemergence

---

**Precautions:** (1) Use only on silt loam or finer textured soil. (2) Wheat-sorghum-fallow cropping sequence must be followed. (3) Do not apply following sorghum harvest.

---

**Note:** Do not graze or feed forage from treated area to livestock. Do not plant any crop other than those listed on this label within 18 months following treatment.

**BEST AVAILABLE COPY**

**Wheat-Corn-Fallow (KS, NE):** This product controls cheatgrass (downy brome, chess), kochia, mustards, pigweed, Russian thistle, wild lettuce, wild sunflower, and volunteer wheat during period after wheat harvest. Weed control may extend into following corn crop grown under minimum tillage.

Follow directions for use, notes and precautions in the Wheat-Sorghum-Fallow section above, substituting corn for references to sorghum.

**Wheat-Fallow-Wheat (CO, KS, MT, NE, ND, SD, and WY):** For preemergence control of cheatgrass (downy brome, chess), common lambsquarters, field pennycress, kochia, mustard, Russian thistle, wild lettuce and suppression of volunteer wheat during fallow period of wheat-fallow-wheat rotations, broadcast 0.6-1.25 lb/A on all soils except those listed under Precautions. For control of pigweed and wild sunflower, use the higher rate. Apply to stubble ground. Treat only once during same fallow period.

#### TANK MIXTURES FOR CHEMICAL FALLOW

**Wheat-Sorghum-Fallow or Wheat-Corn-Fallow (KS, NE):**

**Igran 80W\*:** Use a tank mixture for control of certain broadleaf and grass weeds in wheat-sorghum-fallow or wheat-corn-fallow rotations. Refer to Igran 80W label directions, weeds controlled, precautions, and limitations.

**Gramoxone Super:** If weeds are present at application, a tank mix with Gramoxone Super may be used. Broadcast 3.75 lb of ATRAZINE 80W plus 1-2 pt of Gramoxone Super in 20-60 gal. of water per acre by ground equipment. Add 0.5-1 pt of a nonionic surfactant, such as Activate, per 100 gal. of spray mixture. Add ATRAZINE 80W to spray tank first and thoroughly mix with water. Then add Gramoxone Super, followed by surfactant. Use the 2 pt. rate of Gramoxone Super if weeds are 4-6 inches tall. This mixture will not control weeds taller than 6 inches. Apply to stubble ground. Treat only once during same fallow period. Refer to Gramoxone Super label for further directions, precautions, limitations.

**Wheat-Fallow-Wheat (CO, KS, MT, NE, ND, SD, and WY):**

**Igran 80W:** Use as tank mixture for control of certain broadleaf and grass weeds in wheat-fallow-wheat rotations. Refer to Igran 80W label for directions, weeds controlled, precautions, and limitations.

**Gramoxone Super:** If weeds are present at application, a tank mix with Gramoxone Super may be used. Broadcast 0.6-1.25 lb. of ATRAZINE 80W plus 1-2 pt. of Gramoxone Super in 20-60 gal. of water per acre by ground equipment. Add 0.5-1 pt. of a nonionic surfactant, such as Activate, per 100 gal. of spray mixture. Add ATRAZINE 80W to spray tank first and thoroughly mix with water. Then add Gramoxone Super followed by surfactant. Use the 2 pt. rate of Gramoxone Super if weeds are 4-6 inches tall. This mixture will not control weeds taller than 6 inches. Apply to stubble ground. Treat only once during same fallow period. Refer to Gramoxone Super label for further directions, precautions, and limitations.

if weeds are present at application and this product is used alone, use either an approved contact herbicide before or after treatment, or tillage after treatment.

Use tillage to control weeds which escape during fallow period. Till before planting. For this product applied alone or in tank mixture with Gramoxone Super, plant at least 2 inches deep and 12 months or more after application.

BEST AVAILABLE COPY

Precautions: (1) Do not use on sandy soil. (2) Do not treat eroded hillsides, caliche and rocky outcroppings of exposed calcareous subsoil. (3) Do not treat soils of the Rosebud and Canyon Series in western NE and adjoining counties in CO and WY. (4) Do not treat soils with calcareous surface layers. (5) Avoid spray overlap.

Note: Do not graze treated areas within 6 months after application.

Aerial Application: In order to assure that spray will be controllable within the target area when used according to label directions, make applications at a maximum height of 10 ft. using low drift nozzles at a maximum pressure of 40 psi, and restrict application to periods when wind speed does not exceed 10 mph. To assure that spray will not adversely affect adjacent sensitive nontarget plants, apply ATRAZINE 80W alone by aircraft at a minimum upwind distance of 400 ft. from sensitive plants, or apply ATRAZINE 80W + Igran 80W at a minimum upwind distance of 500 ft. from sensitive plants. When applying ATRAZINE 80W and Igran 80W by aircraft, use a minimum of 5 gal. of water per acre.

#### TURF GRASSES FOR SOD

St. Augustinegrass, Centipedegrass, and Zoysia Grass

Broadcast 2.5-5 lb. per acre according to soil texture to control those weeds listed under ATRAZINE 80W Applied Alone--Corn or Grain Sorghum--Preplant Incorporated or Preemergence.

Muck or Peat	5 lb.	Old beds: Within 2 days after lifting sod.
		New beds: 3-4 days after sprigging or plugging.
Sandy Soil	2.5 lb.	Old beds: Within 2 days after lifting sod.
		New beds: 7-10 days after sprigging or plugging.

If weeds regrow, apply an additional 2.5 lb. per acre on muck or peat, or 1.25 lb. per acre on sandy soil.

Precautions: (1) Do not apply within 30 days prior to cutting or lifting. (2) Do not apply in combination with surfactants or other spray additives, or injury may occur. (3) Do not use north of NC.

#### MACADAMIA NUTS

For preemergence control of many broadleaf and grass weeds, including crabgrass, foxtail, wiregrass, Flora's paintbrush, spanishneedles, and fireweed, broadcast 2.5-5 lb. per acre before harvest and before weeds emerge. Repeat as necessary. Do not spray when nuts are on ground during harvest period. Do not apply by air.

#### PINEAPPLE

For control of purslane, spanishneedles, annual grasses, annual bindweed, ageratum, amaranth, rattlepod, Flora's paintbrush, fireweed, spurge, indigo, and papalo, apply up to 8 lbs./acre as a blanket spray immediately after planting or following harvest. Make additional blanket or interspace applications of up to 2 lbs./acre at 1-2 month intervals as needed prior to differentiation. Apply in 20-40 gal. of water/acre to assure thorough coverage.

Precautions (1) Do not apply more than 37.5 lbs./acre per cycle (2) Repeated monthly applications to plant foliage may slow plant growth and delay fruiting.

**BEST AVAILABLE COPY**

Note: Do not apply within 45 days of fruit harvest or forage harvest if forage is to be fed to livestock.

#### SUGARCANE

For control of many broadleaf grass weeds, including amaranths, crabgrass, fireweed, Flora's paintbrush, foxtails, junglerice and wiregrass, broadcast 2.5-5 lb. per acre of ATRAZINE 80W at time of planting or ratooning, but before sugarcane emerges. Broadcast aerially in a minimum of 5 gal. of spray per acre, or broadcast or band by ground equipment in a minimum of 20 gal. per acre, unless indicated otherwise. One additional application may be made over the sugarcane as it emerges, and two additional aerial applications may be made interline after emergence as directed sprays. Repeat treatments, where needed, may be applied broadcast, band, or interline as suggested with the final application being prior to close-in. Do not exceed the rate of herbicide suggested for any one crop of sugarcane.

Note: Where high rates of ATRAZINE 80W are used alone, apply in a minimum of 1 gal. of water for each 1 lb. of ATRAZINE 80W applied per acre.

Note: Temporary injury to trees may occur following the use of atrazine on coarse-textured soil.

---

Precautions: (1) Do not graze treated areas. (2) Do not apply to seedbeds. (3) Apply only once per year.

---

Aerial application: In order to assure that spray will be controllable within the target area when used according to label directions, make applications at a maximum height of 10 ft. above vegetation, using low drift nozzles at a maximum pressure of 40 psi, and restrict application to periods when wind speed does not exceed 10 mph. To assure that spray will not adversely affect adjacent sensitive nontarget plants, apply ATRAZINE 80W alone by aircraft at a minimum upwind distance of 400 ft. from sensitive plants.

#### Florida

For control of emerged pellitory weed: Apply 0.5-0.75 lb. per acre in at least 40 gal. of water as a directed spray by ground equipment prior to close-in. Add 4 qt. of surfactant for each 100 gal. of spray. Thoroughly cover weed foliage.

For control of alexandergrass, large crabgrass, pellitory (artillery) weed, and spiny amaranth, use one of the following methods at planting or ratooning:

1. Apply 5 lb/A preemergence. Follow with one or two applications, as needed, postemergence to sugarcane and weeds, at 2.5 lb. per acre. Treat before weeds exceed 1.5 inches in height.
2. Apply 1-3 times, as needed, at 2.5 lb/A postemergence to sugarcane and weeds. Treat before weeds exceed 1.5 inches in height.

#### Texas

Use ATRAZINE 80W for control of barnyardgrass, pigweed, purslane and sunflower, in plant or ratoon sugarcane.

**BEST AVAILABLE COPY**

Apply 5 lb. per acre of ATRAZINE 80W preemergence. Follow with one or two applications, as needed, at 3.75 lb. per acre postemergence to sugarcane and weeds.

For best results when weeds are emerged, add a nonionic surfactant at a concentration of 2 qt. per 100 gal. to the spray and apply before weeds exceed 1.5 inches in height.

---

Precautions: (1) Injury to sugarcane may occur when under moisture stress, when soil is of low absorptive capacity, or when land is first cropped to sugarcane. (2) Do not apply after close-in. (3) Do not apply more than 12.5 lb. per acre to any one crop of sugarcane.

---

#### NONSELECTIVE WEED CONTROL ON NONCROP LAND

This product provides long-term weed control on industrial sites, highway medians and shoulders, railroad rights-of-way, lumberyards, petroleum tank farms, and in noncrop areas on farms, such as around buildings, equipment and fuel storage areas, along fences, and lanes. Do not use near adjacent desirable plants or in greenhouses, or injury may occur.

Apply before or soon after weeds begin growth. Make postemergence applications when weeds are young and actively growing. Use the higher rates on fine clay and muck soils. Use sufficient water to assure thorough coverage. Use at least 1 gal. of water for each lb. of product; use more if practical.

Note: Do not allow livestock to graze treated areas.

Annual broadleaf and grass weeds (including barnyardgrass, cheatgrass, crabgrass, lambsquarters, foxtail, ragweed, puncturevine, and turkey mullein): Broadcast 6-12.5 lb. per acre.

Hard-to-kill annual and many perennial broadleaf and grass weeds (including bluegrass, burdock, Canada thistle, dogfennel, orchardgrass, plantain, quackgrass, purpletop, redtop, and smooth brome): Broadcast 12.5 lb. per acre.

For small areas, 4.5 oz. per sq. ft. is equivalent to 12.5 lb. per acre.

#### CONIFERS

For control of annual broadleaf and grass weeds prior to transplanting, after transplanting or in established conifers (including Douglas fir, grand fir, noble fir, white fir, Austrian pine, bishop pine, Jeffrey pine, knobcone pine, loblolly pine, lodgepole pine (shore pine), monterey pine, ponderosa pine, Scotch pine, slash pine, blue spruce, and Sitka spruce): Broadcast 2.5-5 lb. in a minimum of 5 gal. of water per acre by air or 10 gal. by ground before weeds are 1.5 inches tall. Apply to established trees between fall and early spring while trees are dormant. For new transplants, apply during or soon after transplanting. For applications prior to transplanting, allow sufficient precipitation to activate ATRAZINE 80W before transplanting. In areas where spring and summer rainfall is inadequate to activate atrazine, apply during fall prior to spring transplanting.

For the list of weeds controlled, see ATRAZINE 80W Applied Alone--Corn or Grain Sorghum--Preplant Incorporated or Preemergence.

**BEST AVAILABLE COPY**



Quackgrass control: Broadcast 5 lb. in a minimum of 5 gal. of water per acre by air or 10 gal. by ground between fall and early spring while trees are dormant and before quackgrass is more than 1.5 inches tall.

Aerial Application: In order to assure that spray will be controllable within the target area when used according to label directions, make applications at a maximum height of 10 ft. using low drift nozzles at a maximum pressure of 40 psi, and restrict application to periods when wind speed does not exceed 10 mph. To assure that spray will not adversely affect adjacent sensitive nontarget plants, apply ATRAZINE 80W alone by aircraft at a minimum upwind distance of 400 ft. from sensitive plants.

NOTICE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to Seller and Buyer assumes the risk of any such use.

The following are registered trademarks.

Bexton® trademark of Dow Chemical Company for propachlor  
Caliber®, Concep®, Dual® (metolachlor), Igran® (terbutryn), and Princep®  
(simazine) trademarks of Ciba-Geigy  
Gramoxone Super® trademark of ICI Americas for paraquat  
Lasso® (alachlor), Ramrod® (propachlor), and Roundup® (glyphosate) trademarks of  
Monsanto Company  
Sutan® + trademark of Stauffer Chemical Company for butylate