



**For season long weed control in corn and sorghum,  
for weed control in certain other crops, and in non crop areas.**

Riverside Atraside 4L contains 4 lbs. active ingredients per gallon.

**STOP-READ LABEL BEFORE USING!**

## STATEMENT OF PRACTICAL TREATMENT

**IF SWALLOWED:** Contact your nearest poison control center, hospital or physician if the patient is unable to maintain breathing and heart rate (cardio-pulmonary resuscitation). If the patient is conscious and alert, induce vomiting (syringe tip) or stimulate the back of the throat with a finger. NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON.

IF INHALED: ... ..

**IF ON SKIN:** Remove excess oil. Wash with soap and water.

**IF IN EYES** A person with a severe eye injury should not attempt to remove anything from the eye.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA REG NO 5779-255

EPA EST NO 979 A5 13



**RIVERSIDE CHEMICAL CO.**

**RIVERIDE SERIES AGRICULTURE** AGRICULTURE SERIES RIVERIDE

**NET CONTENTS 2½ GALLONS**

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Do not breathe vapors or spray mist. Avoid contact with eyes, skin or clothing. Remove contaminated clothing and wash before reuse. Wash thoroughly with soap and water after handling.

Do not apply this product in such a manner as to directly or indirectly injure persons or other persons. The area being treated must be vacated by unexposed persons.

**NOTE TO PHYSICIANS:** There is no specific antidote for atropine. This product is expected induce effects on the digestive tract. The use of an aqueous slurry of atropine sulfate may be considered.

## ENVIRONMENTAL HAZARDS

1. The following information is being furnished to you for your information only. It is not intended to be used for any other purpose.

### DIRECTIONS FOR USE

1. The first step is to identify the problem or question that needs to be answered. This involves understanding the context and the specific requirements of the task.

## RE-ENTRY STATEMENT

[illegible]

Two dose certain states may require more restrictive controls, such as those being used with this product, and will want State Department of Agriculture to further investigate.

## STORAGE AND DISPOSAL

## DO NOT CONTAMINATE WATER, FOOD, OR FEED BY STORAGE OR DISPOSAL

## STORAGE

Figure 1. The effect of the concentration of the *Agrobacterium* suspension on the transformation efficiency of *Agrobacterium* strains. The concentration of the *Agrobacterium* suspension was 10<sup>6</sup> cells/ml (A), 10<sup>7</sup> cells/ml (B), 10<sup>8</sup> cells/ml (C), and 10<sup>9</sup> cells/ml (D). The concentration of the *Agrobacterium* suspension was 10<sup>6</sup> cells/ml (A), 10<sup>7</sup> cells/ml (B), 10<sup>8</sup> cells/ml (C), and 10<sup>9</sup> cells/ml (D). The concentration of the *Agrobacterium* suspension was 10<sup>6</sup> cells/ml (A), 10<sup>7</sup> cells/ml (B), 10<sup>8</sup> cells/ml (C), and 10<sup>9</sup> cells/ml (D). The concentration of the *Agrobacterium* suspension was 10<sup>6</sup> cells/ml (A), 10<sup>7</sup> cells/ml (B), 10<sup>8</sup> cells/ml (C), and 10<sup>9</sup> cells/ml (D).

1. The first step is to identify the problem or question that needs to be answered. This involves understanding the context and the specific requirements of the task.

[illegible]

## DISPOSAL

PESTICIDE DISPOSAL CONTAINER INFORMATION SHEET  
 USE THIS FORM TO REPORT THE DISPOSITION OF PESTICIDE CONTAINERS TO THE APPROPRIATE AGENCY.  
 IF YOU ARE NOT SURE WHICH AGENCY TO CONTACT, CALL THE NATIONAL PESTICIDE DISPOSAL HOTLINE AT 1-800-456-7766.

CONTAINER DISPOSAL: This container and its contents shall not be disposed of in the trash or sewer. It should be disposed of in accordance with the following instructions: 1. If the container is empty, it should be disposed of in the trash. 2. If the container contains a liquid, it should be disposed of in the sewer. 3. If the container contains a solid, it should be disposed of in the trash. 4. If the container contains a gas, it should be disposed of in the trash. 5. If the container contains a flammable liquid, it should be disposed of in the trash. 6. If the container contains a flammable gas, it should be disposed of in the trash. 7. If the container contains a toxic liquid, it should be disposed of in the trash. 8. If the container contains a toxic gas, it should be disposed of in the trash. 9. If the container contains a radioactive material, it should be disposed of in the trash. 10. If the container contains a hazardous material, it should be disposed of in the trash. 11. If the container contains a biohazardous material, it should be disposed of in the trash. 12. If the container contains a chemical, it should be disposed of in the trash. 13. If the container contains a biological material, it should be disposed of in the trash. 14. If the container contains a physical material, it should be disposed of in the trash. 15. If the container contains a radioactive material, it should be disposed of in the trash. 16. If the container contains a hazardous material, it should be disposed of in the trash. 17. If the container contains a biohazardous material, it should be disposed of in the trash. 18. If the container contains a chemical, it should be disposed of in the trash. 19. If the container contains a biological material, it should be disposed of in the trash. 20. If the container contains a physical material, it should be disposed of in the trash.

## APPLICATION PROCEDURES

Use only ATRASIDE 4L in conventional ground sprayers equipped with a pump, nozzle, agitator and uniform application system. Maintain a minimum spray distance of 20 inches. The spray should be applied in a continuous manner. When spraying, a tank should be filled with water before spraying. The spray should be applied in a continuous manner. When spraying, a tank should be filled with water before spraying.

For best results, apply ATRASIDE 4L in a continuous manner. When spraying, a tank should be filled with water before spraying. The spray should be applied in a continuous manner. When spraying, a tank should be filled with water before spraying.

For best results, apply ATRASIDE 4L in a continuous manner. When spraying, a tank should be filled with water before spraying. The spray should be applied in a continuous manner. When spraying, a tank should be filled with water before spraying.

For best results, apply ATRASIDE 4L in a continuous manner. When spraying, a tank should be filled with water before spraying. The spray should be applied in a continuous manner. When spraying, a tank should be filled with water before spraying.

For best results, apply ATRASIDE 4L in a continuous manner. When spraying, a tank should be filled with water before spraying. The spray should be applied in a continuous manner. When spraying, a tank should be filled with water before spraying.

For best results, apply ATRASIDE 4L in a continuous manner. When spraying, a tank should be filled with water before spraying. The spray should be applied in a continuous manner. When spraying, a tank should be filled with water before spraying.

For best results, apply ATRASIDE 4L in a continuous manner. When spraying, a tank should be filled with water before spraying. The spray should be applied in a continuous manner. When spraying, a tank should be filled with water before spraying.

For best results, apply ATRASIDE 4L in a continuous manner. When spraying, a tank should be filled with water before spraying. The spray should be applied in a continuous manner. When spraying, a tank should be filled with water before spraying.

For best results, apply ATRASIDE 4L in a continuous manner. When spraying, a tank should be filled with water before spraying. The spray should be applied in a continuous manner. When spraying, a tank should be filled with water before spraying.

For best results, apply ATRASIDE 4L in a continuous manner. When spraying, a tank should be filled with water before spraying. The spray should be applied in a continuous manner. When spraying, a tank should be filled with water before spraying.

For best results, apply ATRASIDE 4L in a continuous manner. When spraying, a tank should be filled with water before spraying. The spray should be applied in a continuous manner. When spraying, a tank should be filled with water before spraying.

For best results, apply ATRASIDE 4L in a continuous manner. When spraying, a tank should be filled with water before spraying. The spray should be applied in a continuous manner. When spraying, a tank should be filled with water before spraying.

For best results, apply ATRASIDE 4L in a continuous manner. When spraying, a tank should be filled with water before spraying. The spray should be applied in a continuous manner. When spraying, a tank should be filled with water before spraying.

For best results, apply ATRASIDE 4L in a continuous manner. When spraying, a tank should be filled with water before spraying. The spray should be applied in a continuous manner. When spraying, a tank should be filled with water before spraying.

For best results, apply ATRASIDE 4L in a continuous manner. When spraying, a tank should be filled with water before spraying. The spray should be applied in a continuous manner. When spraying, a tank should be filled with water before spraying.

REV 11-84



# ATRASIDE 4L

## NOTICE

The directions on this product reflect the opinion of experts for intended use and tests. The directions are believed to be safe and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Using this product in a manner not intended or for purposes not intended may result in injury, death, or property damage. The user assumes the responsibility for the use of this product. The user assumes the responsibility for the use of this product.

RIVERSIDE CHEMICAL COMPANY warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. But neither this warranty nor any other warranty of merchantability or fitness for particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not known to RIVERSIDE CHEMICAL COMPANY, and buyer assumes the risk of any such use. RIVERSIDE CHEMICAL COMPANY shall not be responsible for incidental damages, if any, resulting from a breach of warranty.

## GENERAL INFORMATION

This herbicide controls many annual broadleaf and grass weeds in corn, sorghum, sugarcane, and certain other crops specified on this label. It is also effective in noncrop areas and industrial sites for control of most annual and many perennial broadleaf and grass weeds. This product may be applied before or after weeds emerge.

Following many years of continuous use of this product and chemically related products, biotypes of some of the weeds listed on this label have been reported which cannot be effectively controlled by this and related herbicides. Where this is known or suspected, we recommend the use of this product in combination with other registered herbicides which are not triazines. Consult with your State Agricultural Extension Service for specific recommendations.

Since this product acts mainly through soil absorption, its effectiveness depends on moisture to move it into the root zone. Weeds growing in a dry soil, however, may be controlled by generally resulting in better weed control.

This product is nonflammable.

Avoid using near adjacent desirable plants or in greenhouses, etc. may injure.

Where the use directions give a range of rates, use the lower rate on coarse-textured soil and soil low in organic matter; use the higher rate on fine-textured soil and soil high in organic matter.


NOTE: Riverside Chemical Company does not recommend applications in combination with other herbicides or oils, except as specifically described on the label or in literature published by Riverside Chemical Company.

(Continued on back panel)

SEE ATTACHED BOOKLET FOR COMPLETE DIRECTIONS FOR USE

## ATRASIDE 4L

For best results, apply ATRASIDE 4L in a continuous manner. When spraying, a tank should be filled with water before spraying. The spray should be applied in a continuous manner. When spraying, a tank should be filled with water before spraying.



DO NOT DETACH BOOKLET FROM CONTAINER

KEEP OUT OF REACH OF CHILDREN

**CAUTION**

See inside booklet for Precautionary Statements and Statement of Practical Treatment

EPA Reg. No. 4774-255

EPA Est. No. 4775 AR 12

BEST AVAILABLE COPY

BEST BEFORE DATE

150000

9179-255

IF INHALED: Remove to fresh air. If breathing is difficult, use oxygen.  
IF ON SKIN: Remove contaminated clothing and wash with soap and water.  
IF IN EYES: Flush with water for 15 minutes. Seek medical attention.

### PRECAUTIONARY STATEMENTS CAUTION

**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**  
Irritant to the skin. May cause allergic reactions. May cause respiratory irritation. May cause eye irritation. May cause skin irritation. May cause skin dryness. May cause skin cracking. May cause skin peeling. May cause skin blistering. May cause skin ulcers. May cause skin cancer. May cause skin deformities. May cause skin death.

Do not apply to skin. Do not use on children. Do not use on pregnant women. Do not use on nursing infants. Do not use on the face. Do not use on the neck. Do not use on the chest. Do not use on the abdomen. Do not use on the back. Do not use on the arms. Do not use on the legs. Do not use on the feet. Do not use on the hands. Do not use on the fingers. Do not use on the toes. Do not use on the nails. Do not use on the hair. Do not use on the scalp. Do not use on the beard. Do not use on the mustache. Do not use on the eyebrows. Do not use on the eyelashes. Do not use on the lips. Do not use on the mouth. Do not use on the throat. Do not use on the windpipe. Do not use on the trachea. Do not use on the bronchi. Do not use on the lungs. Do not use on the heart. Do not use on the liver. Do not use on the kidneys. Do not use on the bladder. Do not use on the intestines. Do not use on the stomach. Do not use on the pancreas. Do not use on the spleen. Do not use on the gallbladder. Do not use on the bile ducts. Do not use on the pancreas. Do not use on the spleen. Do not use on the gallbladder. Do not use on the bile ducts.

**NOTE TO PHYSICIANS:** There is no specific antidote for atrazine. If this product is ingested, induce emesis or lavage stomach. The use of an aqueous slurry of activated charcoal may be considered.

### ENVIRONMENTAL HAZARDS

Do not apply directly to any body of water. Do not contaminate water. Do not use equipment or disposal of waste.

**BEST AVAILABLE COPY**

### DIRECTIONS FOR USE

## STORAGE AND DISPOSAL

*Journal of Interpersonal Violence* 28(10)

## DISP-05A

**PESTICIDE DISPOSAL:** Excess of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL** These use the container as a receptacle for recycling or for packaging, or production and dispose of in a sanitary landfill or incineration. It is allowed by state and local authorities by burning.

### GENERAL INFORMATION

The first of these methods, which is the most widely used, is the "single-stage" method. This method involves the use of a single stage of treatment, which is typically a combination of a chemical and a physical process. This method is also effective in removing a wide range of pollutants, including organic and inorganic compounds, and is often used in conjunction with other treatment methods.

Do not apply if wind velocity is greater than 15 mph. Do not apply if rain is forecast within 2 hours of application. Do not apply if temperature is below 50°F. Do not apply if relative humidity is greater than 90%. Do not apply if wind velocity is greater than 15 mph. Do not apply if rain is forecast within 2 hours of application. Do not apply if temperature is below 50°F. Do not apply if relative humidity is greater than 90%.

[illegible]

For each pint of RIVERSIDE ATHASIDE 4L or other liquid flowable product to be applied per acre, add 0.5 teaspoon or 1/40th of a liter of the product to one pint of fluid fertilizer (assuming a volume of 25 gallons per acre).

After thorough mixing, let stand for 5 minutes. If the herbicide/fertilizer combination remains mixed or can be removed readily, the mixture is compatible and can be sprayed. If it is not compatible, make a water slurry with the herbicide before adding fertilizer and use the mixture only if the test shows it is compatible.

the fact that the *in vitro* and *in vivo* experiments have shown that the release of the drug from the implant is controlled by the rate of degradation of the polymer. The release of the drug from the implant is controlled by the rate of degradation of the polymer. The release of the drug from the implant is controlled by the rate of degradation of the polymer.

Applied Along Edge of Main Member

### Woods Controlled

imprisoned  
 tochio  
 a mink's quarters  
 annual mox, angkor in  
 mustards  
 nightshade  
 jagged  
 our scene  
 jagged  
 velvet at  
 (a) (b) (c) (d) (e) (f) (g) (h) (i) (j) (k) (l) (m) (n) (o) (p) (q) (r) (s) (t) (u) (v) (w) (x) (y) (z)

100-443887-100

1962, 1963, 1964, 1965, 1966, 1967, 1968, 1969, 1970, 1971, 1972, 1973, 1974, 1975, 1976, 1977, 1978, 1979, 1980, 1981, 1982, 1983, 1984, 1985, 1986, 1987, 1988, 1989, 1990, 1991, 1992, 1993, 1994, 1995, 1996, 1997, 1998, 1999, 2000, 2001, 2002, 2003, 2004, 2005, 2006, 2007, 2008, 2009, 2010, 2011, 2012, 2013, 2014, 2015, 2016, 2017, 2018, 2019, 2020, 2021, 2022, 2023, 2024, 2025, 2026, 2027, 2028, 2029, 2030, 2031, 2032, 2033, 2034, 2035, 2036, 2037, 2038, 2039, 2040, 2041, 2042, 2043, 2044, 2045, 2046, 2047, 2048, 2049, 2050, 2051, 2052, 2053, 2054, 2055, 2056, 2057, 2058, 2059, 2060, 2061, 2062, 2063, 2064, 2065, 2066, 2067, 2068, 2069, 2070, 2071, 2072, 2073, 2074, 2075, 2076, 2077, 2078, 2079, 2080, 2081, 2082, 2083, 2084, 2085, 2086, 2087, 2088, 2089, 2090, 2091, 2092, 2093, 2094, 2095, 2096, 2097, 2098, 2099, 2100, 2101, 2102, 2103, 2104, 2105, 2106, 2107, 2108, 2109, 2110, 2111, 2112, 2113, 2114, 2115, 2116, 2117, 2118, 2119, 2120, 2121, 2122, 2123, 2124, 2125, 2126, 2127, 2128, 2129, 2130, 2131, 2132, 2133, 2134, 2135, 2136, 2137, 2138, 2139, 2140, 2141, 2142, 2143, 2144, 2145, 2146, 2147, 2148, 2149, 2150, 2151, 2152, 2153, 2154, 2155, 2156, 2157, 2158, 2159, 2160, 2161, 2162, 2163, 2164, 2165, 2166, 2167, 2168, 2169, 2170, 2171, 2172, 2173, 2174, 2175, 2176, 2177, 2178, 2179, 2180, 2181, 2182, 2183, 2184, 2185, 2186, 2187, 2188, 2189, 2190, 2191, 2192, 2193, 2194, 2195, 2196, 2197, 2198, 2199, 2200, 2201, 2202, 2203, 2204, 2205, 2206, 2207, 2208, 2209, 2210, 2211, 2212, 2213, 2214, 2215, 2216, 2217, 2218, 2219, 2220, 2221, 2222, 2223, 2224, 2225, 2226, 2227, 2228, 2229, 2230, 2231, 2232, 2233, 2234, 2235, 2236, 2237, 2238, 2239, 2240, 2241, 2242, 2243, 2244, 2245, 2246, 2247, 2248, 2249, 2250, 2251, 2252, 2253, 2254, 2255, 2256, 2257, 2258, 2259, 2260, 2261, 2262, 2263, 2264, 2265, 2266, 2267, 2268, 2269, 2270, 2271, 2272, 2273, 2274, 2275, 2276, 2277, 2278, 2279, 2280, 2281, 2282, 2283, 2284, 2285, 2286, 2287, 2288, 2289, 2290, 2291, 2292, 2293, 2294, 2295, 2296, 2297, 2298, 2299, 2300, 2301, 2302, 2303, 2304, 2305, 2306, 2307, 2308, 2309, 2310, 2311, 2312, 2313, 2314, 2315, 2316, 2317, 2318, 2319, 2320, 2321, 2322, 2323, 2324, 2325, 2326, 2327, 2328, 2329, 2330, 2331, 2332, 2333, 2334, 2335, 2336, 2337, 2338, 2339, 2340, 2341, 2342, 2343, 2344, 2345, 2346, 2347, 2348, 2349, 2350, 2351, 2352, 2353, 2354, 2355, 2356, 2357, 2358, 2359, 2360, 2361, 2362, 2363, 2364, 2365, 2366, 2367, 2368, 2369, 2370, 2371, 2372, 2373, 2374, 2375, 2376, 2377, 2378, 2379, 2380, 2381, 2382, 2383, 2384, 2385, 2386, 2387, 2388, 2389, 2390, 2391, 2392, 2393, 2394, 2395, 2396, 2397, 2398, 2399, 2400, 2401, 2402, 2403, 2404, 2405, 2406, 2407, 2408, 2409, 2410, 2411, 2412, 2413, 2414, 2415, 2416, 2417, 2418, 2419, 2420, 2421, 2422, 2423, 2424, 2425, 2426, 2427, 2428, 2429, 2430, 2431, 2432, 2433, 2434, 2435, 2436, 2437, 2438, 2439, 2440, 2441, 2442, 2443, 2444, 2445, 2446, 2447, 2448, 2449, 2450, 2451, 2452, 2453, 2454, 2455, 2456, 2457, 2458, 2459, 2460, 2461, 2462, 2463, 2464, 2465, 2466, 2467, 2468, 2469, 2470, 2471, 2472, 2473, 2474, 2475, 2476, 2477, 2478, 2479, 2480, 2481, 2482, 2483, 2484, 2485, 2486, 2487, 2488, 2489, 2490, 2491, 2492, 2493, 2494, 2495, 2496, 2497, 2498, 2499, 2500, 2501, 2502, 2503, 2504, 2505, 2506, 2507, 2508, 2509, 2510, 2511, 2512, 2513, 2514, 2515, 2516, 2517, 2518, 2519, 2520, 2521, 2522, 2523, 2524, 2525, 2526, 2527, 2528, 2529, 2530, 2531, 2532, 2533, 2534, 2535, 2536, 2537, 2538, 2539, 2540, 2541, 2542, 2543, 2544, 2545, 2546, 2547, 2548, 2549, 2550, 2551, 2552, 2553, 2554, 2555, 2556, 2557, 2558, 2559, 2560, 2561, 2562, 2563, 2564, 2565, 2566, 2567, 2568, 2569, 2570, 2571, 2572, 2573, 2574, 2575, 2576, 2577, 2578, 2579, 2580, 2581, 2582, 2583, 2584, 2585, 2586, 2587, 2588, 2589, 2590, 2591, 2592, 2593, 2594, 2595, 2596, 2597, 2598, 2599, 2600, 2601, 2602, 2603, 2604, 2605, 2606, 2607, 2608, 2609, 2610, 2611, 2612, 2613, 2614, 2615, 2616, 2617, 2618, 2619, 2620, 2621, 2622, 2623, 2624, 2625, 2626, 2627, 2628, 2629, 2630, 2631, 2632, 2633, 2634, 2635, 2636, 2637, 2638, 2639, 2640, 2641, 2642, 2643, 26

[illegible]

Prescriptions for Application with 1 March 2006

[illegible]

### Problem Words

Yellow Nuttall's and Canada Thistle. The plots with the yellow nuttall's *Cyperus micranthus* and Canada thistle *Cirsium arvense* were applied with the best results. The best results were obtained with the yellow nuttall's and Canada thistle. The yellow nuttall's and Canada thistle were neither applied to the plots with the yellow nuttall's and Canada thistle. The yellow nuttall's and Canada thistle were neither applied to the plots with the yellow nuttall's and Canada thistle.

6.           [illegible]

...the ... .. the ... ..  
 ... ..  
 ... ..  
 ... ..

1. Broadleaf ... .. of oil  
 ... .. Canada  
 ... .. height  
 ... .. inches

... ..  
 ... ..  
 ... ..  
 ... ..

... ..  
 ... ..  
 ... ..  
 ... ..

... ..  
 ... ..  
 ... ..

Note: ... ..  
 ... ..  
 ... ..

#### Quackgrass Control on Land going into Corn Production.

**Split Application** ... ..  
 ... ..  
 ... ..  
 ... ..

**Single Application** ... ..  
 ... ..

**TANK MIXTURES FOR CORN**  
 This product may be tank mixed with these herbicides for control  
 of certain broadleaf and grass weeds in corn  
 Dual® (metolachlor) Bladex®

... ..  
 ... ..  
 ... ..  
 ... ..

... ..  
 ... ..  
 ... ..

... ..  
 ... ..  
 ... ..

... ..  
 ... ..  
 ... ..

**Preplant** ... ..  
 ... ..

**Preemergence** ... ..  
 ... ..

Refer to Corn sections of this label and to Princeps 60W, Princeps  
 Control 30, or Princeps 4, 4.7, label for further corn herbicide  
 information and directions.

**BEST AVAILABLE COPY**



Preemergence (Broadleaf and grass control) - High water table

TABLE 3. Preemergence Broadleaf and Grass Weed Control in Sorghum

| State | Rate (lb/A) | Remarks                       |
|-------|-------------|-------------------------------|
| AZ    | 2.4         | Apply before sorghum emerges. |
| CA    | 2.4         | Apply before sorghum emerges. |
| CO    | 2.4         | Apply before sorghum emerges. |
| KS    | 2.4         | Apply before sorghum emerges. |
| NM    | 2.4         | Apply before sorghum emerges. |
| OK    | 2.4         | Apply before sorghum emerges. |
| TX    | 2.4         | Apply before sorghum emerges. |

Preemergence Broadleaf and Grass Weed Control in Sorghum

In case of planting failure, sorghum can be replanted 30 days after emergence of weeds. If weeds are not controlled, sorghum can be replanted 30 days after emergence of weeds.

Preemergence Broadleaf weed control in furrow irrigated bedded sorghum (AZ and CA only) - High water table. Apply before sorghum emerges. Apply 2.4 lb/A of emulsifiable oil or oil concentrate in water. Apply 0.75-1.5 pt of surfactant after sorghum reaches 6 inches in height. In case of crop failure, do not replant sorghum for 8 months following application. Corn may be planted immediately.

**Precautions for Preemergence Application to Furrow Irrigated Bedded Sorghum in AZ and CA**  
In AZ and CA, sorghum injury may occur on sand or silty sand soils when sorghum is planted in furrows. Applications to sorghum growing on alkali soils or where cuts, fills, or erosion have exposed calcareous or alkali subsoils may cause crop injury. In case of crop failure, do not replant sorghum for 8 months following application. Corn may be planted immediately.

Postemergence (Broadleaf and grass control) - High water table. Apply before weeds exceed 1.5 inches in height.

TABLE 4. Postemergence Broadleaf and Grass Weed Control in Sorghum

| State | Rate (lb/A) | Remarks                       |
|-------|-------------|-------------------------------|
| AZ    | 2.4         | Apply before sorghum emerges. |
| CA    | 2.4         | Apply before sorghum emerges. |
| CO    | 2.4         | Apply before sorghum emerges. |
| KS    | 2.4         | Apply before sorghum emerges. |
| NM    | 2.4         | Apply before sorghum emerges. |
| OK    | 2.4         | Apply before sorghum emerges. |
| TX    | 2.4         | Apply before sorghum emerges. |

Postemergence Broadleaf Weed Control with Emulsifiable Oil or Oil Concentrate in Water - Broadcast 2.4 lb/A of emulsifiable oil or oil concentrate in water. Apply before pigweed and white squatters reach 6 inches in height and before all other weeds reach 4 inches in height. In CO, western KS, NM, OK, TX, and desert regions of AZ and CA, apply when sorghum is 6 inches in height. Do not replant sorghum for 8 months following application. Corn may be planted immediately.

Postemergence Broadleaf Weed Control with Emulsifiable Oil or Oil Concentrate in Water - Broadcast 2.4 lb/A of emulsifiable oil or oil concentrate in water. Apply before pigweed and white squatters reach 6 inches in height and before all other weeds reach 4 inches in height. In CO, western KS, NM, OK, TX, and desert regions of AZ and CA, apply when sorghum is 6 inches in height. Do not replant sorghum for 8 months following application. Corn may be planted immediately.

Postemergence Broadleaf Weed Control with Emulsifiable Oil or Oil Concentrate in Water - Broadcast 2.4 lb/A of emulsifiable oil or oil concentrate in water. Apply before pigweed and white squatters reach 6 inches in height and before all other weeds reach 4 inches in height. In CO, western KS, NM, OK, TX, and desert regions of AZ and CA, apply when sorghum is 6 inches in height. Do not replant sorghum for 8 months following application. Corn may be planted immediately.

**Precautions for Applications with Emulsifiable Oil or Oil Concentrate in Water:** See Precautions for Applications with Emulsifiable Oil or Oil Concentrate in Water in Corn section.

**Postemergence Broadleaf Weed Control with Surfactant (CO, western KS, NM, OK, TX, and desert regions of AZ and CA only):** Broadcast 2.4 lb/A plus 0.75-1.5 pt. of surfactant after sorghum reaches 6 inches in height but before weeds exceed 1.5 inches in height. Apply only on sandy loam and finer textured soil.

**BEST AVAILABLE COPY**

**Rotation Crops** \_\_\_\_\_

2014-2015

Dear Sir,

14

• Surinam Paper Co. V. United States, 100 F. 2d 1001, 1002 (2d Cir. 1937).

1990

15

BEST AVAILABLE COPY

Wheat-Fallow Wheat (CA, ID, NV, OR, UT, and WY)  
Igran 80W use as directed on the label for the broadcast

Wheat-Fallow Wheat (CA, ID, NV, OR, UT, and WY)  
Igran 80W use as directed on the label for the broadcast

#### Parasol Control

Parasol Control is a selective herbicide for the control of annual and biennial weeds in wheat-fallow wheat. It is applied as a broadcast or spot treatment. It is effective against a wide range of weeds, including annual and biennial weeds, and is safe for wheat.

Parasol Control is a selective herbicide for the control of annual and biennial weeds in wheat-fallow wheat. It is applied as a broadcast or spot treatment. It is effective against a wide range of weeds, including annual and biennial weeds, and is safe for wheat.

Parasol Control is a selective herbicide for the control of annual and biennial weeds in wheat-fallow wheat. It is applied as a broadcast or spot treatment. It is effective against a wide range of weeds, including annual and biennial weeds, and is safe for wheat.

#### Precautions

Do not graze treated areas until the risk of injury to livestock is eliminated. Do not graze treated areas until the risk of injury to livestock is eliminated. Do not graze treated areas until the risk of injury to livestock is eliminated.

**Note:** Do not graze treated areas until the risk of injury to livestock is eliminated. Do not graze treated areas until the risk of injury to livestock is eliminated. Do not graze treated areas until the risk of injury to livestock is eliminated.

#### PROSE-MALLET

PROSE-MALLET is a selective herbicide for the control of annual and biennial weeds in wheat-fallow wheat. It is applied as a broadcast or spot treatment. It is effective against a wide range of weeds, including annual and biennial weeds, and is safe for wheat.

#### PROSE-MALLET

PROSE-MALLET is a selective herbicide for the control of annual and biennial weeds in wheat-fallow wheat. It is applied as a broadcast or spot treatment. It is effective against a wide range of weeds, including annual and biennial weeds, and is safe for wheat.

#### PROSE-MALLET

PROSE-MALLET is a selective herbicide for the control of annual and biennial weeds in wheat-fallow wheat. It is applied as a broadcast or spot treatment. It is effective against a wide range of weeds, including annual and biennial weeds, and is safe for wheat.

**Aerial Application:** To assure that spray will be confined within the target area when used according to label directions, apply PROSE-MALLET at a height of 10 ft. above the ground, at a maximum pressure of 40 psi, and restrict application to periods when wind speed does not exceed 10 mph.

PROSE-MALLET is a selective herbicide for the control of annual and biennial weeds in wheat-fallow wheat. It is applied as a broadcast or spot treatment. It is effective against a wide range of weeds, including annual and biennial weeds, and is safe for wheat.

PROSE-MALLET is a selective herbicide for the control of annual and biennial weeds in wheat-fallow wheat. It is applied as a broadcast or spot treatment. It is effective against a wide range of weeds, including annual and biennial weeds, and is safe for wheat.

**Rehabilitation of existing stands of perennial range grasses:** Central Great Plains (CO, KS, MT, NE, ND, SD, WY). Broadcast 1-2 pt. per acre in fall before ground freezes.

**Southern Great Plains (OK, TX):** Broadcast 2 pt. per acre in spring before April 30.

**West (CA, ID, NV, OR, UT, WA):** Do not graze area to be treated for 6 months prior to application. Broadcast 1-1.5 pt. per acre in fall before ground freezes.

PROSE-MALLET is a selective herbicide for the control of annual and biennial weeds in wheat-fallow wheat. It is applied as a broadcast or spot treatment. It is effective against a wide range of weeds, including annual and biennial weeds, and is safe for wheat.

BEST AVAILABLE COPY

Apply to the soil surface, after the grass has been mowed, or to the foliage of the grass.

Established stands of grass

Apply to the soil surface, after the grass has been mowed, or to the foliage of the grass.

Apply to the soil surface, after the grass has been mowed, or to the foliage of the grass.

Precautions: A.

Note: Do not graze treated area with livestock for 14 days after application. Do not plant treated area for other crops for 14 months.

#### TURF GRASSES FOR SOO

St. Augustinegrass, Centipedegrass, and Zoysia Grass

Apply to the soil surface, after the grass has been mowed, or to the foliage of the grass.

Apply to the soil surface, after the grass has been mowed, or to the foliage of the grass.

#### 4. ADAMIA NUTS

Apply to the soil surface, after the grass has been mowed, or to the foliage of the grass.

#### ARE

Apply to the soil surface, after the grass has been mowed, or to the foliage of the grass.

Apply to the soil surface, after the grass has been mowed, or to the foliage of the grass.

Note: Where high rates of RIVERSIDE ATRASIDE 4L are used, apply in a minimum of 1 qt. of water for each 1 qt. of ATRASIDE 4L applied per acre.

Application: In order to assure that spray will be controllable to target area when used according to label directions, operate at a maximum height of 10 ft. using low drift nozzles with pressure of 40 psi and restrict application

BEST AVAILABLE COPY

to be made when wind speed for the spray will not exceed 10 mph. Do not spray if wind speed is 10 mph or more. Do not spray if wind speed is 10 mph or more. Do not spray if wind speed is 10 mph or more.

#### Florida

For control of annual broadleaf and grass weeds in sugarcane, apply REVER-side ATRASIDE 4L at 4 pt. per acre. Apply 2 to 3 times as needed at 4 pt. per acre. For control of annual broadleaf and grass weeds in sugarcane, apply REVER-side ATRASIDE 4L at 4 pt. per acre. Apply 2 to 3 times as needed at 4 pt. per acre.

For control of annual broadleaf and grass weeds in sugarcane, apply REVER-side ATRASIDE 4L at 4 pt. per acre. Apply 2 to 3 times as needed at 4 pt. per acre. For control of annual broadleaf and grass weeds in sugarcane, apply REVER-side ATRASIDE 4L at 4 pt. per acre. Apply 2 to 3 times as needed at 4 pt. per acre.

#### Texas

For control of annual broadleaf and grass weeds in sugarcane, apply REVER-side ATRASIDE 4L at 4 pt. per acre. Apply 2 to 3 times as needed at 4 pt. per acre. For control of annual broadleaf and grass weeds in sugarcane, apply REVER-side ATRASIDE 4L at 4 pt. per acre. Apply 2 to 3 times as needed at 4 pt. per acre.

**Precautions:** (1) Injury to sugarcane may occur when the mixture is applied when the sugarcane is in the seedling stage. (2) Do not apply after the sugarcane has reached the seedling stage. (3) Do not apply more than 2.5 pt. per acre to any one crop of sugarcane.

#### CONIFERS

For control of annual broadleaf and grass weeds prior to transplanting, after transplanting or in established conifers (including Douglas fir, grand fir, noble fir, white fir, Austrian pine, bishop pine, Jeffrey pine, knobcone pine, loblolly pine, lodgepole pine (shore pine), Monterey pine, ponderosa pine, Scotch pine, slash pine, blue spruce, and Sitka spruce) Broadcast 4-8 pt. in a minimum of 5 gal. of water per acre by air or 10 gal. by ground before weeds are 1.5 inches tall. Apply to established trees between fall and early spring while trees are

For control of annual broadleaf and grass weeds in sugarcane, apply REVER-side ATRASIDE 4L at 4 pt. per acre. Apply 2 to 3 times as needed at 4 pt. per acre. For control of annual broadleaf and grass weeds in sugarcane, apply REVER-side ATRASIDE 4L at 4 pt. per acre. Apply 2 to 3 times as needed at 4 pt. per acre.

For control of annual broadleaf and grass weeds in sugarcane, apply REVER-side ATRASIDE 4L at 4 pt. per acre. Apply 2 to 3 times as needed at 4 pt. per acre. For control of annual broadleaf and grass weeds in sugarcane, apply REVER-side ATRASIDE 4L at 4 pt. per acre. Apply 2 to 3 times as needed at 4 pt. per acre.

#### INSELECTIVE WEED CONTROL NONCROP LAND

For control of annual broadleaf and grass weeds in sugarcane, apply REVER-side ATRASIDE 4L at 4 pt. per acre. Apply 2 to 3 times as needed at 4 pt. per acre. For control of annual broadleaf and grass weeds in sugarcane, apply REVER-side ATRASIDE 4L at 4 pt. per acre. Apply 2 to 3 times as needed at 4 pt. per acre.

For control of annual broadleaf and grass weeds in sugarcane, apply REVER-side ATRASIDE 4L at 4 pt. per acre. Apply 2 to 3 times as needed at 4 pt. per acre. For control of annual broadleaf and grass weeds in sugarcane, apply REVER-side ATRASIDE 4L at 4 pt. per acre. Apply 2 to 3 times as needed at 4 pt. per acre.

For control of annual broadleaf and grass weeds in sugarcane, apply REVER-side ATRASIDE 4L at 4 pt. per acre. Apply 2 to 3 times as needed at 4 pt. per acre. For control of annual broadleaf and grass weeds in sugarcane, apply REVER-side ATRASIDE 4L at 4 pt. per acre. Apply 2 to 3 times as needed at 4 pt. per acre.

For control of annual broadleaf and grass weeds in sugarcane, apply REVER-side ATRASIDE 4L at 4 pt. per acre. Apply 2 to 3 times as needed at 4 pt. per acre. For control of annual broadleaf and grass weeds in sugarcane, apply REVER-side ATRASIDE 4L at 4 pt. per acre. Apply 2 to 3 times as needed at 4 pt. per acre.

For control of annual broadleaf and grass weeds in sugarcane, apply REVER-side ATRASIDE 4L at 4 pt. per acre. Apply 2 to 3 times as needed at 4 pt. per acre. For control of annual broadleaf and grass weeds in sugarcane, apply REVER-side ATRASIDE 4L at 4 pt. per acre. Apply 2 to 3 times as needed at 4 pt. per acre.

For control of annual broadleaf and grass weeds in sugarcane, apply REVER-side ATRASIDE 4L at 4 pt. per acre. Apply 2 to 3 times as needed at 4 pt. per acre. For control of annual broadleaf and grass weeds in sugarcane, apply REVER-side ATRASIDE 4L at 4 pt. per acre. Apply 2 to 3 times as needed at 4 pt. per acre.

For control of annual broadleaf and grass weeds in sugarcane, apply REVER-side ATRASIDE 4L at 4 pt. per acre. Apply 2 to 3 times as needed at 4 pt. per acre. For control of annual broadleaf and grass weeds in sugarcane, apply REVER-side ATRASIDE 4L at 4 pt. per acre. Apply 2 to 3 times as needed at 4 pt. per acre.

14 of 14

**RIVERSIDE CHEMICAL CO.**  
 2595 ARCADE DRIVE • ARLINGTON, VA 22204  
 (703) 521-1100 • FAX (703) 521-1101 • WWW.RIVERSIDECHEM.COM  
 RIVERSIDE SERVES AGRICULTURE AGRICULTURE SERVES RIVERSIDE

BEST AVAILABLE COPY