

2,4-D LV6 (9779-256)
Riverside/Terra Corporation

**FOR USE IN CROP RESIDUE MANAGEMENT SYSTEMS
IN SOYBEANS (Preplant only)**

DIRECTIONS FOR USE

It is a violation of federal law to use this product
in a manner inconsistent with its labeling.

Weeds	Amount of Product to Use Per Acre	Directions
Postemergence	1/2 to 2/3 pint	Apply not less than 7 days prior to planting soybeans when weeds are small and actively growing. Use the higher rate on larger weeds and when perennials are present.
	2/3 to 1 1/3 pts	Apply not less than 30 days prior to planting soybeans when weeds are actively growing.

In addition to those weeds found on the WEED LIST, this product will suppress or control the following broadleaf weeds frequently encountered in reduced tillage soybean production systems: bullnettle, smallflowered bittercress, Carolina geranium, smallflowered buttercup, common and rough cinquefoil, red clover,* horseweed or maretail, mousetail, wild mustard, field pennycress, cutleaf evening primrose, common purslane, speedwell, and Virginia copperleaf.

*These weeds are only partially controlled.

Apply no more than 1 1/3 pints of this product in one season prior to planting soybeans. After applying, plant soybean seed as deep as practical or at least 1-1/2 to 2 inches deep. Adjust the planter press wheel, if necessary, to ensure that planted seed is completely covered.

ACCEPTED
12 APR 1993
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 9779-256

BEST AVAILABLE COPY

Aerial Application: When applying aerially, use 2 or more gallons of total spray volume per acre.

Ground Application: With ground equipment, use 10 or more gallons of total spray volume per acre.

If desired, this product may be applied preplant to soybeans in tank mixtures with other herbicides such as Poast, Poast Plus, Roundup, Roundup D-Pak, Honcho, Gramoxone Extra, Prowl, Pursuit Plus, Scepter 70 DG, Squadron and others that are registered for preplant soybean use.

NOTE: Unacceptable injury to soybeans planted in fields previously treated with this product may occur. Whether or not soybean injury occurs and the extent of the injury will depend on weather and agronomic factors such as the amount of weed vegetation and previous crop residue present that may be in effect between the time of application and the emergence of the soybean plant.

Not registered for use in California.

RESTRICTIONS AND LIMITATIONS FOR USE IN SOYBEANS (PREPLANT)

BEFORE USING 2,4-D LV4, READ AND CAREFULLY OBSERVE ALL APPLICABLE DIRECTIONS, RESTRICTIONS, PRECAUTIONS, AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.

Do not apply this product prior to planting soybeans if you are not prepared to accept the results of soybean injury including possible loss of stand and yield.

Do not apply this product when weather conditions such as temperature, air inversions, or wind favor drift from treated areas to susceptible plants.

Apply no more than 1 1/3 pints of this product per acre in one season prior to planting soybeans.

Do not mow or cultivate weeds prior to treating with this product as poor control may result.

Do not feed treated hay, forage, or fodder or graze treated soybeans to livestock.

Only one application of this product may be made, regardless of application rate, per growing season.

Do not feed or graze treated cover crops to livestock.

BEST AVAILABLE COPY