

2,4-D LV4 (9779-257)  
Riverside/Terra Corporation

FOR USE IN CROP RESIDUE MANAGEMENT SYSTEMS  
IN SOYBEANS (Preplant only)

DIRECTIONS FOR USE


It is a violation of federal law to use this product  
in a manner inconsistent with its labeling.

Weeds	Amount of Product to Use Per Acre	Directions
Postemergence	3/4 to 1 pint	Apply not less than 7 days prior to planting soybeans wher weeds are small and actively growing. Use the higher rate on larger weeds and when perennials are present.
	1 to 2 pints	Apply not less than 30 days prior to planting soybeans when weeds are active- ly growing.

In addition to those weeds found on the  
WEED LIST, this product will suppress or  
control the following broadleaf weeds fre-  
quently encountered in reduced tillage -  
soybean production systems: bullnettle,  
smallflowered bittercress, Carolina geran-  
ium, smallflowered buttercup, common and  
rough cinquefoil, red clover,\* horseweed  
or mareetail, mousetail, wild mustard,  
field pennycress, cutleaf evening prim-  
rose, common purslane, speedwell, and Vir-  
ginia copperleaf.

\*These weeds are only partially con-  
trolled.

Apply no more than 2 pints of this  
product in one season prior to planting  
soybeans. After applying, plant soybean  
seed as deep as practical or at least 1-  
1/2 to 2 inches deep. Adjust the planter  
press wheel, if necessary, to ensure that  
planted seed is completely covered.



**BEST AVAILABLE COPY**

2 9 2

**Aerial Application:** When applying aerially, use 2 or more gallons of total spray volume per acre.

**Ground Application:** With ground equipment, use 10 or more gallons of total spray volume per acre.

If desired, this product may be applied preplant to soybeans in tank mixtures with other herbicides such as Poast, Poast Plus, Roundup, Roundup D-Pak, Honcho, Gramoxone Extra, Prowl, Pursuit Plus, Scepter 70 DG, Squadron and others that are registered for preplant soybean use.

**NOTE:** Unacceptable injury to soybeans planted in fields previously treated with this product may occur. Whether or not soybean injury occurs and the extent of the injury will depend on weather and agronomic factors such as the amount of weed vegetation and previous crop residue present that may be in effect between the time of application and the emergence of the soybean plant.

Not registered for use in California.

#### RESTRICTIONS AND LIMITATIONS FOR USE IN SOYBEANS (PREPLANT)

**BEFORE USING 2,4-D LV4, READ AND CAREFULLY OBSERVE ALL APPLICABLE DIRECTIONS, RESTRICTIONS, PRECAUTIONS, AND ALL OTHER INFORMATION APPEARING ON THE PRODUCT LABEL.**

Do not apply this product prior to planting soybeans if you are not prepared to accept the results of soybean injury including possible loss of stand and yield.

Do not apply this product when weather conditions such as temperature, air inversions, or wind favor drift from treated areas to susceptible plants.

Apply no more than 2 pints of this product per acre in one season prior to planting soybeans.

Do not replant fields treated with this product in the same growing season with crops other than those labeled for 2,4-D preplant use.

Do not mow or cultivate weeds prior to treating with this product as poor control may result.

Do not feed treated hay, forage, or fodder or graze treated soybeans to livestock.

Only one application of this product may be made, regardless of application rate, per growing season.

Do not feed or graze treated cover crops to livestock.

**BEST AVAILABLE COPY**