

PM21 9779-277
9779-277

87.0

Riverside Trademark®

MANEB 8% DUST

POTATO SEED-PIECE FUNGICIDE DUST

ACTIVE INGREDIENT	
Maneb (manganese ethylenebisdithiocarbamate).....	8.00%
INERT INGREDIENTS.....	92.00%
	Total 100.00%
Manganese, equivalent as metallic 1.37%	

STOP-READ LABEL BEFORE USING!

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center at once.
 IF IN EYES: Flush eyes immediately with water for at least 15 minutes. Call a physician.
 IF ON SKIN: Wash affected area with soap and water.
 IF INHALED: Remove victim to fresh air. Apply respiration if indicated.

SEE LEFT PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

EPA REG. NO. 9779-277


EPA EST. NO. _____

Riverside
Trademark®

MANUFACTURED FOR
 RIVERSIDE/TERRA CORPORATION
 A SUBSIDIARY OF TERRA INTERNATIONAL, INC.
 TERRA CENTRE, 600 FOURTH STREET, SIOUX CITY, IOWA 51101
 RIVERSIDE SERVES AGRICULTURE. AGRICULTURE SERVES EVERYONE.

NET CONTENTS _____ POUNDS

ACCEPTED
 AUG 21 1987
 Under the Federal Insecticide, Fungicide, and Plant Disease Act
 and related laws, this pesticide
 registration is for
 EPA Reg. No. 9779-277



(Left Panel)

PRECAUTIONARY STATEMENTS

CAUTION

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

May cause irritation to nose, throat, eyes, and skin. Do not breathe dust or spray mist. In case of contact with eyes, flush immediately with plenty of water; for skin, wash thoroughly. Get medical attention if irritation persists.

Wash thoroughly after using and before eating or smoking. Remove and wash contaminated clothing before reusing.

Avoid contamination of feed or foodstuffs. Do not store with food.

Do not store with feed or forage to be fed to dairy animals being finished for slaughter.

Treated seed must not be used for or mixed with food or animal feed. Treat only those seed pieces needed for immediate use, minimizing the interval between treatment and planting.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

PHYSICAL OR CHEMICAL HAZARDS

Keep away from fire and sparks. Store in a cool, dry place. Do not allow to become wet or overheated in storage. This may bring on chemical changes which will impair the fungicidal effectiveness of the product and may also generate flammable vapors. Keep container closed when not in use.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

20/4

STORAGE AND DISPOSAL

DO NOT CONTAMINATE WATER, FOOD, OR FEED BY STORAGE OR DISPOSAL.

STORAGE

Store in a cool, dry location away from children, animals foods, feeds, seeds, or other agricultural chemicals. Do not store in direct sunlight. Protect from excessive heat. Handle in accordance with information given under "Precautionary Statements". In the event of spillage or leakage, scrape up and dispose of in accordance with information given under "DISPOSAL." Repackage and relabel useable product in a sound container. In case of fire or other emergency, report at once by toll-free telephone to 800-424-9300.

DISPOSAL

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.
CONTAINER DISPOSAL: Plastic Containers-Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

GENERAL INFORMATION

MANEB 8% DUST controls fusarium seed-piece decay, the most common seed- and soil-borne fungus disease of potatoes.

MANEB 8% DUST helps prevent seed-piece rot under adverse sprouting conditions (cold and wet, or dry soil conditions).

MANEB 6% DUST protects seed which is held for planting.

POTATO SEED-PIECE: Cutting and Treating - Before planting, cutting machines or knives, trays, tables, barrels or sacks, equipment, trucks and planters should be thoroughly cleaned and sanitized. If not sprouted, subject seed potatoes to temperature of 50° to 60°F for about 10 to 15 days before handling them.

Dosage and Application: Use 1 pound of MANEB 8% DUST per 100 pounds of cut seed-pieces. Apply it so that the entire surface of the seed-piece is thoroughly covered by the fungicide dust. Place treated seed in clean, disease-free containers or store in bulk at proper temperature and humidity. Prevent inhalation of dust by use of exhaust fans during treatment.

The cut and treated seed may be planted immediately, but preferably should be held for a week or more at cool temperature and high relative humidity. This holding period promotes suberization and healing of the cut surfaces. Good suberization reduces the susceptibility of the seed-piece to infection by other disease organisms, especially in cold, wet soil.

Do not use excess treated seed-pieces for food or feed.

3/7/4

NOTICE: RIVERSIDE/TERRA CORPORATION warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to RIVERSIDE/TERRA CORPORATION and buyer assumes the risk of any such use. RIVERSIDE/TERRA CORPORATION shall not be responsible for incidental or consequential damages, if any, resulting from a breach of warranty.

4/0/01