

NOV 30 1989

Beverley Neale  
Label Development Specialist  
Riverside/Terra Corporation  
P.O. box 171376  
Memphis, TN 38187

Dear Ms. Neale:

Subject: Sodium Chlorate 6  
EPA Registration No. 9779-305  
RE: Label Amendment To Add Dry Beans and Southern Peas  
Your Submission Dated October 17, 1989

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended is acceptable with the following changes:

- \* Under PRECAUTIONARY STATEMENTS add the phrase: "Wear goggles or face shield."
- \* Under STORAGE AND DISPOSAL delete the phrase: "Repackage and relabel usable product in a sound container." Add the phrase: "Store in original containers only."

Please submit five (5) copies of your final printed labeling for the above mentioned product incorporating the amendment before you release the product for shipment. A stamped copies of the reviewed draft label is enclosed for your records.

Sincerely yours,

*Robert J. Taylor*  
 Robert J. Taylor  
 Product Manager (25)  
 Fungicide-Herbicide Branch  
 Registration Division (P7505C)

**BEST AVAILABLE COPY**

CONCURRENCES							
SYMBOL	M 117505C						
SURNAME	M Marshall						
DATE	M 11/20/89						

296

R•

Sodium Chlorate 6

Cotton defoliant and desiccant

---

<b>ACTIVE INGREDIENT</b>	
Sodium Chlorate.....	47.2%
<b>INERT INGREDIENTS.....</b>	<b>52.8%</b>
	<b>TOTAL 100.0%</b>

Product contains 6 pounds of sodium chlorate per gallon.

STOP--READ LABEL BEFORE USING.

KEEP OUT OF REACH OF CHILDREN  
WARNING/AVISO

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Flush with water. Get medical attention.  
 IF SWALLOWED: Drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Avoid alcohol. Get medical attention.  
 IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists.  
 NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

See additional PRECAUTIONARY STATEMENTS on side panel.

EPA Reg. No. 9779-305

EPA Est. No. 9779-

Manufactured For  
 RIVERSIDE/TERRA CORPORATION  
 Terra Centre, 600 Fourth Street, Sioux City, Iowa 51101  
 Riverside Serves Agriculture. Agriculture Serves Everyone.

NET CONTENTS  
GALS.

ACCEPTED  
with COMMENTS  
in EPA Letter Dated:

NOV 30 1989

Use of this Federal Insecticide,  
Fungicide and Rodenticide Act  
is authorized, for the pesticide  
registered under EPA Reg. No.

9779-305

396

---

**PRECAUTIONARY STATEMENTS  
WARNING  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

Causes eye irritation. Do not get in eyes, on skin, or on clothing. Harmful if swallowed. Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

**ENVIRONMENTAL HAZARDS**

Do not apply directly to water or wetlands. Do not contaminate water when disposing of equipment washwaters.

---

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product through any type of irrigation system.

**REENTRY STATEMENT**

Do not enter treated areas without protective clothing until sprays have dried. Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

**NOTICE TO CROP OWNERS**

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is a reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information: **WARNING.** Area treated with Sodium Chlorate or (date of application). Do not enter without appropriate protective clothing until sprays have dried. In case of accidental exposure refer to **STATEMENT OF PRACTICAL TREATMENT** on the Sodium Chlorate 6 label.

**STORAGE AND DISPOSAL**  
**DO NOT CONTAMINATE WATER, FOOD OR FEED BY STORAGE OR DISPOSAL.**  
**STORAGE**

Store in a dry location away from children, animals, foods, feeds, seeds, or other agricultural chemicals. Handle in accordance with information given under PRECAUTIONARY STATEMENTS. In the event of spillage or leakage, soak up material with absorbent clay, sand, sawdust, or other absorbent material. Scrape up and dispose of in accordance with information given under DISPOSAL. Repackage and relabel usable product in a sound container. In case of fire or other emergency, report at once by toll-free telephone to 800-424-9300.

**DISPOSAL**

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.  
**CONTAINER DISPOSAL:** Plastic Containers-Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke. Metal Containers-Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedure approved by state and local authorities.

**GENERAL INFORMATION**

Do not allow spray to contact adjacent crops or other plants that may be defoliated. Clean and wash spray equipment before and after use and do not allow wash water to contact desirable plants. To avoid fire or explosion, do not mix with insecticides.

**BEANS (DRY):** To facilitate the harvest of edible dry beans, apply one gallon of Sodium Chlorate 6 in enough water to give thorough coverage, either by aircraft or ground equipment. For air application, use a minimum of 5 gallons of ready-to-apply material per acre; for ground equipment use minimum of 20 gallons of total solution per acre. Make applications 7 to 10 days before anticipated harvest date when beans are mature and ready for harvest.

**CHILIPEPPERS:** Apply 1/2 to 1-1/4 gallons per acre in 5 to 15 gallons of water by aircraft and 20 to 40 gallons of water by ground rig. Use on peppers for processing only. Apply 10 days before harvest. When grown for processing, consult a processor before applying.

**CORN:** Sodium Chlorate 6 may be used as a harvest aid for desiccation of trashy weeds in early maturing corn. Use 1 gallon Sodium Chlorate 6 in 5-7 gallons of water per acre by air. Apply at least 14 days before anticipated harvest date on clear days when temperature is expected to go above 70°F. Use the 5 gallon rate when weeds are small and the crop canopy is fairly open. Do not graze treated fields or feed fodder, forage or residual seeds within 14 days of application.

**COTTON DEFOLIATION:** Plants must be mature to get results with a defoliant. The usual signs of maturity include a slight yellowing of the foliage and a reddish tinge on the leaves with a clean break by a slight downward pull. Plants should not be drought-stricken because the tough wilted leaves do not permit uptake of the chemical, or growth of an abscission layer. When defoliants are applied to drought-stricken plants, leaves will die but usually not drop, thereby causing extra trash to be picked up in harvesting.

Weather conditions that favor good defoliation are high temperatures, high humidity, low wind velocity and high to adequate soil moisture. Plant conditions which favor good defoliation are ample fertility and moisture, complete insect control, heavy boll load and bolls mature, vegetable growth ceased, and little or no second growth.

Apply 1/2 to 3/4 gallons of Sodium Chlorate 6 per acre in sufficient water to give thorough coverage. When using ground equipment, apply 15 to 20 gallons of spray solution per acre, and by air, 5 gallons of spray solution per acre. Thorough coverage is essential.

Apply Sodium Chlorate 6 to mature cotton plants after the earliest bolls expected to make cotton are at least 30 days old. Do not apply later than 7 days before harvest. Do not graze treated areas or feed gin waste to livestock.

FLAX: For use as a harvest aid: To desiccate trashy weeds and desiccate the mature crop, apply 1 gallon Sodium Chlorate 6 per acre in enough water to give thorough coverage. Make application on bright, sunny days (preferably when temperature is above 60°F) 7-10 days before anticipated harvest date. For air application, use a minimum of 5 gallons ready-to-apply material per acre. Activity of the spray will be enhanced by the addition of a good non-ionic surfactant at the rate of 1 quart per 100 gallons of spray. Do not graze treated fields or feed treated foliage and fodder for 14 days after application.

GRAIN SORGHUM: Sodium Chlorate 6 may be used to reduce the moisture content in grain sorghum prior to harvest. It may be applied by aircraft or ground equipment. Thorough coverage of the grain sorghum is essential.

To reduce the moisture content of grain while the crop is standing, apply 3/4 to 1 gallon Sodium Chlorate 6 per acre in sufficient water to give thorough coverage. Apply from 7 to 10 days before harvest. Apply a total of 5 to 10 gallons of spray per acre by airplanes and 10 to 20 gallons per acre by ground equipment. Use the lower rate when grain moisture is low and the weather is clear and dry. Use the higher rate when conditions for desiccation are poor.

GUAR BEANS: To facilitate harvest and reduce bean seed moisture, apply 1 gallon of Sodium Chlorate 6 in enough water to provide thorough coverage either by aircraft or ground equipment. Make application 7-10 days before anticipated harvest date. Do not apply under conditions of extreme heat during the middle of the day. For air application, use 5-10 gallons of total solution per acre; for ground application, use a minimum of 20 gallons of total solution per acre. Do not graze treated fields or feed treated fodder or forage to livestock.

RICE: To facilitate harvest and reduce moisture in the fully mature seed heads and to desiccate immature red rice in mature white rice, apply by aircraft 3/4 to 1 gallon of Sodium Chlorate 6 per acre in 5 or more gallons of water per acre. Use the lower rate when crop conditions and weather are right to enhance rapid drying. Apply 7 to 10 days before anticipated harvest. To avoid drift, do not apply during windy weather. Thorough coverage of the crop is essential to achieve good results. Do not graze treated fields or feed treated straw or crop residues within 14 days of application.

SAFFLOWER: To facilitate harvest and reduce moisture in the fully mature seed heads, apply 1 gallon of Sodium Chlorate 6 in sufficient water to give good coverage, 7 or more days before normal harvest when safflower heads are fully mature. Apply in 5 to 10 gallons of water per acre by air or in 20 to 30 gallons of water per acre by ground. Do not apply later than 7 days before harvest. Do not graze treated fields or feed treated forage.

**SOUTHERN PEAS:** To defoliate plants and facilitate harvest, apply one gallon of Sodium Chlorate 6 in enough water to provide thorough coverage either by aircraft or ground equipment. Make application 5 to 7 days before anticipated harvest date. Do not apply under conditions of extreme heat during the middle of the day. For air application, use 5 to 10 gallons of total solution per acre; for ground application, use a minimum of 20 gallons of total solution per acre.

**SOYBEANS:** As a soybean harvest aid to dry weeds in early maturing soybeans and to facilitate the harvest, apply 1 gallon of Sodium Chlorate 6 per acre in enough water to give thorough coverage; application may be made by ground equipment or aircraft. Make application 7 to 10 days before anticipated harvesting date when beans are mature and ready for harvest. For air application use a minimum of 5 gallons ready-to-apply spray solution per acre; for ground equipment apply a minimum of 20 gallons of ready-to-apply spray solution per acre. Do not graze treated fields or feed treated soybean foliage.

**SUNFLOWER:** To facilitate harvest and reduce moisture in the fully mature seed heads, apply 1 gallon Sodium Chlorate 6 in sufficient water to give good coverage, 7 or more days before normal harvest. Apply in 5 to 10 gallons of water per acre by air or in 20 to 30 gallons of water per acre by ground with a high clearance sprayer. Do not apply later than 7 days before harvest. Do not graze treated fields or feed treated forage.

**NOTICE:** Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to Seller and Buyer assumes the risk of any such use.

.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....