

Maxon

With plant growth regulators

ACTIVE INGREDIENTS.....	0.0126%
Gibberellic Acid.....	0.0030% ✓
Cytokinin (as kinetin).....	0.0096%
INERT INGREDIENTS.....	99.9874%

Contains 113 mg/gal. gibberellic acid and 363 mg/gal. cytokinin.

GUARANTEED ANALYSIS

Sulfur (S) (combined).....	1.0000%
Boron (B).....	0.0005%
Cobalt (Co).....	0.0005%
Copper (Cu) (chelated).....	0.0500%
Iron (Fe) (chelated).....	0.0500%
Molybdenum (Mo).....	0.0005%
Zinc (Zn) (chelated).....	0.0500%

Derived from sodium borate, cobalt sulfate, copper EDTA, iron HEDTA, sodium molybdate, and zinc EDTA.

ACCEPTED
 Feb 13 1991
 Under the Federal Insecticide, Fungicide, and Plant Pesticide Act, as amended, for the pesticide registration number: EPA Reg. No. 9779-314

STOP - READ LABEL BEFORE USING.

KEEP OUT OF REACH OF CHILDREN

CAUTION

BEST AVAILABLE COPY

STATEMENT OF PRACTICAL TREATMENT

- IF INHALED: Remove victim to fresh air and apply respiration if indicated.
- IF ON SKIN: Remove contaminated clothing and wash with soap and water.
- IF IN EYES: Flush with plenty of water. Get medical attention if irritation persists.

See additional PRECAUTIONARY STATEMENTS below:

EPA Reg. No. 9779-

EPA Est. Nos. 9779-TX-3, 9779-MS-1

Manufactured For
 RIVERSIDE/TERRA CORPORATION
 Terra Centre, 600 Fourth Street, Sioux City, Iowa 51101
 Riverside Serves Agriculture. Agriculture Serves Everyone.

NET CONTENTS 1 GAL.

NET WT. 8.34 LBS.

PRECAUTIONARY STATEMENTS
CAUTION
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Avoid contact with skin, eyes, or clothing. Wash hands thoroughly after handling. Wash all clothing with soap and water before reuse.

ENVIRONMENTAL HAZARDS

Do not apply directly to water. Do not apply when weather conditions favor drift from treated areas. Do not apply where runoff is likely to occur. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product through any type of irrigation system.

RE-ENTRY STATEMENT

Do not enter treated areas without protective clothing until sprays have dried. Because certain states may require more restrictive re-entry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

BEST AVAILABLE COPY

STORAGE AND DISPOSAL

DO NOT CONTAMINATE WATER, FOOD, OR FEED BY STORAGE OR DISPOSAL.

STORAGE

Do not store below 40°F. Store out of direct sunlight. Handle in accordance with information given under PRECAUTIONARY STATEMENTS. In the event of spillage or leakage, soak up material with absorbent clay, sand, sawdust, or other absorbent material. Scrape up and dispose of in accordance with information given under DISPOSAL. Repackage and relabel useable product in a sound container. In case of fire or other emergency, report at once by toll-free telephone to 800-424-9300.

DISPOSAL

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: For one gallon plastic containers or larger, triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke. For containers less than one gallon, wrap in newspaper after emptying and dispose in household trash.

GENERAL INFORMATION

Maxon is a combination of plant growth hormones and fertilizer, formulated to stimulate seedling vigor, flower and fruit set, and to increase crop yields. All recommendations are for foliar (broadcast) applications unless otherwise stated. The use of a foliar fertilizer following or in combination with Maxon applications enhances its activity.

PREPARATION AND APPLICATION OF SPRAY SOLUTIONS

Fill spray tank half full with clean water. Start agitation. If compatibility problems are suspected, add an approved compatibility agent at labeled rates. Shake Maxon well and pour directly from the container into the partially filled spray tank. Add the rest of the water. Provide agitation during mixing and application to maintain a uniform suspension.

Maxon may be applied either by ground or air equipment. When applied alone or with foliar fertilizer, use an approved non-ionic surfactant to enhance the wetting and spreading characteristics of the spray solution. When used with other pesticides, follow the label requirements of the pesticide used.

CROPS

BEANS, PEAS (Green, Lima, Pinto, Blackeye, Mung):

If mechanically harvested: Make a broadcast application of 6 oz/acre at the first sign of bloom.

If handpicked: Make a broadcast application of 1 to 2 oz/acre at time of early bloom. Wait 7 to 14 days, then broadcast 1 to 2 oz/acre. Continue to broadcast 1 to 2 oz/acre on a 7 to 14 day schedule throughout the growing season.

BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER: Make an application of 3 to 6 oz/acre broadcast 14 days after transplanting. Follow with a second application of 3 to 6 oz/acre broadcast 28 days after transplanting.

COTTON: Apply 4 oz/acre broadcast at matchhead square. Make a second application of 4 oz/acre at first bloom.

CUCUMBERS, SQUASH: At the 2 to 4 leaf stage, apply 3 oz/acre broadcast. Follow with 3 oz/acre broadcast at early bloom. Wait 14 to 21 days and broadcast 2 oz/acre. Continue to broadcast 1 to 2 oz/acre every 14 to 21 days throughout the growing season.

LETTUCE, SPINACH: Make an application of 6 oz/acre at the 3 to 4 leaf stage. For spinach, broadcast 2 oz/acre after each cutting.

MELONS (Cantaloupes, Honeydews, Watermelons): Make an application of 3 oz/acre broadcast at first sign of running and again two weeks after running.

MILO (Grain Sorghum): Apply 8 oz/acre broadcast over the plants at the 4 to 6 leaf stage.

PEANUTS: Apply 6 oz/acre broadcast at initial pegging.

BEST AVAILABLE COPY

PEPPERS (Bell, Jalapeno, Serrano): Apply 2 oz/acre broadcast two weeks after transplanting, followed by 2 oz/acre at early bloom. Wait 21 days, then broadcast 2 oz/acre. Continue to broadcast 1 to 2 oz/acre on a 21 day schedule up to 2 weeks prior to harvest.

RICE: Make one application of 6 to 8 oz/acre at the 3 to 5 leaf stage.

SOYBEANS: Apply 6 oz/acre broadcast over the plants at first sign of bloom.

STRAWBERRIES: Apply 3 oz/acre broadcast at early bloom.

TOMATOES:

If machine harvested: Apply 6 oz/acre broadcast slightly before or at first bloom.

If hand picked: Apply 1 to 2 oz/acre broadcast slightly before or at first bloom. Continue to broadcast 1 to 2 oz/acre on a 7 to 14 day schedule throughout the growing season.

YOUNG TREES, ORNAMENTALS, SOD, LAWNS, TURF

ORANGE TREES: Spray foliage of young trees to drip or run-off two to three times per year. Use 1 oz. per 2 gallons of water.

SHRUBS (Established): Spray foliage with 1 oz. Maxon in 2 gallons of water. Spray to point of run-off, two or three times a year.

SOD: During initial sod production (4-6 weeks prior to harvest), broadcast 8 oz/acre. For regrowth enhancement after harvest, broadcast 4 oz/acre. Repeat application of 4 oz/acre six weeks later.

LAWNS (Established): Use 1 oz. broadcast per 5,000 square feet.

TURF (Immediately after laying): Broadcast 2 oz/5,000 square feet and water in. Thirty days later, make a second application of 2 oz/5,000 square feet and water in.

GOLF COURSES

GREENS: Use 2 oz. of Maxon per 5,000 sq. ft. of golf green area 60 days prior to first killing frost. Apply 1 oz. per 5,000 sq. ft. 30 days later. In spring, resume applications of 1 oz. per 5,000 sq. ft. every 30 days.

TEES: Use 2 oz. of Maxon per 5,000 sq. ft. every 30 days.

FAIRWAYS: Apply 8 oz. per acre in early to mid spring. Repeat application mid to later summer.

BEST AVAILABLE COPY

NOTICE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to Seller and Buyer assumes the risk of any such use.