

9779-314

11/04/2005

1/5

1 of 5

Maxon™

With plant growth regulators

ACTIVE INGREDIENTS

Gibberellic Acid (GA ₄ + GA ₇)	0.0033%
Cytokinin (as kinetin)	0.0097%

OTHER INGREDIENTS	99.9870%
TOTAL	100.0000%

Contains 29.9 mg/L gibberellic acid and 95.9 mg/L cytokinin

FERTILIZER GUARANTEED ANALYSIS 0-0-0

Sulfur (S) (combined)	1.0000%
Boron (B)	0.0005%
Cobalt (Co)	0.0005%
Copper (Cu)	0.0500%
.05% chelated copper	
Iron (Fe)	0.0500%
0.05% chelated iron	
Molybdenum (Mo)	0.0005%
Zinc (Zn)	0.0500%
.05% chelated zinc	

ACCEPTED

NOV 04 2005

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 9779-314

Derived from sodium borate, cobalt sulfate, copper EDTA, iron HEDTA, sodium molybdate, and zinc EDTA.

KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID	
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance; then give artificial respiration, preferably by mouth-to-mouth if possible. • Call a poison control center or doctor for treatment advice.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment. In case of a health emergency call 1-877-424-7452.</p>	

Read additional PRECAUTIONARY STATEMENTS on page two.

EPA Reg. No. 9779-314

Manufactured For
Agrilience, LLC
P.O. Box 64089, St. Paul, MN 55164-0089
Web: www.agrilience.com

EPA Est. No.

Net Contents

0/02013

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

Caution: Harmful if inhaled or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Avoid breathing spray mist. Wash thoroughly with soap and water after handling. Remove and wash contaminated clothing before reuse. Wear the appropriate Personal Protective Equipment (PPE).

Personal Protective Equipment:

Mixers, loaders, applicators and other handlers must wear:

- long-sleeved shirt and long pants,
- chemical-resistant gloves, and
- shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

For terrestrial uses: Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Do not apply where runoff is likely to occur. Do not contaminate water when disposing of equipment washwater or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product through any type of irrigation system.

For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply

to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours unless wearing appropriate PPE.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- coveralls,
- chemical-resistant gloves, and
- shoes plus socks.

Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Keep unprotected persons or pets out of treated areas until sprays have dried.

GENERAL INFORMATION

Maxon is a combination of plant growth hormones and fertilizer, formulated to stimulate seedling vigor, flower and fruit set, and to increase crop yields. All instructions are for foliar (broadcast) applications unless otherwise stated. The use of a foliar fertilizer following or in combination with Maxon applications enhances its activity.

Where a range of dosage rates is given, use the higher rate for maximum yield potential. For a more economical program, use the lower rate.

PREPARATION AND APPLICATION OF SPRAY SOLUTIONS

Fill spray tank half full with the desired amount of clean water. Start agitation. If compatibility problems are suspected, add an approved compatibility agent at labeled rates. Shake Maxon well and pour the required amount directly from the container into the partially filled spray tank. Add the rest of the water. Provide agitation during mixing and application to maintain a uniform suspension.

Apply Maxon either by ground in 5 to 50 gallons of water or by air in 1 to 10 gallons of water per acre. Use sufficient water to obtain uniform coverage without excessive run-off. Use higher volumes of water within the ranges given above in dense crop canopies or where increased coverage is needed. When applied alone or with foliar fertilizer, use an approved non-ionic surfactant to enhance the wetting and spreading characteristics of the spray solution. When used with other pesticides, follow the label requirements of the pesticide used.


CROPS

(Rates are given in fluid ounces of product before dilution).

BEANS, PEAS (Green, Lima, Pinto, Blackeye, Mung):

If mechanically harvested: Make a broadcast application of 6 fl. oz. of Maxon per acre at the first sign of bloom.

If handpicked: Make a broadcast application of 1 to 2 fl. oz. Maxon per acre at time of early bloom. Wait 7 to 14 days, then broadcast 1 to 2 fl. oz. per acre on a 7 to 14 day schedule throughout the growing season.



BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER: Make an application of 3 to 6 fl. oz. of Maxon per acre broadcast 14 days after transplanting. Follow with a second application of 3 to 6 fl. oz. per acre broadcast 28 days after transplanting.

COTTON: Apply 4 fl. oz. Maxon per acre broadcast at matchhead square. Make a second application of 4 fl. oz. per acre at first bloom.

CUCUMBERS, SQUASH: At the 2 to 4 leaf stage, apply 3 fl. oz. of Maxon per acre broadcast. Follow with 3 oz. per acre broadcast at early bloom. Wait 14 to 21 days and broadcast 2 fl. oz. per acre. Continue to broadcast 1 to 2 fl. oz. per acre every 14 to 21 days throughout the growing season.

LETTUCE, SPINACH: Make an application of 6 fl. oz. Maxon per acre at the 3 to 4 leaf stage. For spinach, broadcast 2 fl. oz. per acre after each cutting.

MELONS (Cantaloupes, Honeydews, Watermelons): Make an application of 3 fl. oz. of Maxon per acre broadcast at first sign of running and again two weeks after running.

MILO (Grain Sorghum): Apply 8 fl. oz. Maxon per acre broadcast over the plants at the 4 to 6 leaf stage.

PEANUTS: Apply 6 fl. oz. of Maxon per acre broadcast at initial pegging.

PEPPERS (Bell, Jalapeno, Serrano): Apply 2 fl. oz. of Maxon per acre broadcast two weeks after transplanting, followed by 2 fl. oz. per acre at early bloom. Wait 21 days, then broadcast 2 fl. oz. per acre. Continue to broadcast 1 to 2 fl. oz. per acre on a 21 day schedule up to 2 weeks prior to harvest.

RICE: Make one application of 6 to 8 fl. oz. of Maxon per acre at the 3 to 5 leaf stage.

SOYBEANS: Apply 6 fl. oz. Maxon per acre broadcast over the plants at first sign of bloom.

STRAWBERRIES: Apply 3 fl. oz. of Maxon per acre broadcast at early bloom.

TOMATOES: If machine harvested: Apply 6 fl. oz. of Maxon per acre broadcast slightly before or at first bloom.

If hand picked: Apply 1 to 2 fl. oz. of Maxon per acre broadcast slightly before or at first bloom. Continue to broadcast 1 to 2 fl. oz. per acre on a 7 to 14 day schedule throughout the growing season.

YOUNG TREES, ORNAMENTALS, SOD, LAWNS, TURF

ORANGE TREES: Spray foliage of young trees to drip or run-off two to three times per year. Use 1 fl. oz. of Maxon per 2 gallons of water.

SHRUBS (Established): Spray foliage with 1 fl. oz. Maxon in 2 gallons of water. Spray to point of run-off, two or three times a year.

SOD: During initial sod production (4-6 weeks prior to harvest), broadcast 8 fl. oz. Maxon per acre. For regrowth enhancement after harvest, broadcast 4 fl. oz. per acre. Repeat application of 4 fl. oz. per acre six weeks later.

LAWNS (Established): Use 1 fl. oz. Maxon broadcast per 5,000 square feet.

TURF (Immediately after laying): Broadcast 2 fl. oz. Maxon per 5,000 square feet and water in. Thirty days later, make a second application of 2 fl. oz. per 5,000 square feet and water in.

GOLF COURSES

GREENS: Use 2 fl. oz. of Maxon per 5,000 sq. ft. of golf green area 60 days prior to first killing frost. Apply 1 fl. oz. per 5,000 sq. ft. 30 days later. In spring, resume applications of 1 fl. oz. per 5,000 sq. ft. every 30 days.

TEES: Use 2 fl. oz. of Maxon per 5,000 sq. ft. every 30 days.

FAIRWAYS: Apply 8 fl. oz. of Maxon per acre in early to mid spring. Repeat application mid to late summer.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Storage

Store the product between 40°F and 120°F. Store out of direct sunlight. Handle in accordance with information given under PRECAUTIONARY STATEMENTS. In the event of spillage or leakage, soak up material with absorbent clay, sand, sawdust, or other absorbent material. Scrape up and dispose of in accordance with information given under DISPOSAL. In case of fire or other emergency, report at once by toll-free telephone to 800-424-9300.

Disposal

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

WARRANTY STATEMENT: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, NOR IS ANY REPRESENTATIVE OF SELLER AUTHORIZED TO MAKE ANY SUCH WARRANTY OR MODIFY THESE TERMS. This warranty does not extend to the storage, handling or use of this product contrary to label instructions.

