



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

September 1, 2016

Lisa Mathias
Product Registration Specialist
Winfield Solutions LLC
PO Box 64589
St. Paul, MN 55164

Subject: Label Amendment: Adds Me-Too crops: Napa cabbage, Chinese Mustard, Cavalo Broccoli, Kohlrabi, Cucurbit, Chinese waxgourd, Gourds, Balsam Apple, Bitter Melon; Eggplant, Groundcherry, Okra, Pepino, Pepper (includes bell pepper, chili pepper, cooking pepper, pimento, sweet pepper), Tomatillo, Ginseng, Horseradish, Lupine, Lentil; Persimmon (Florida and Hawaii only), Rhubarb, Yam, Boston Fern, Florida Ruffle Fern, Leatherleaf Fern. Other minor changes throughout the label

Product Name: Terranil 6L
EPA Registration Number: 9779-320
Application Date: March 17, 2016
Decision Number: 517622

Dear Ms. Mathias:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false

Page 2 of 2
EPA Reg. No. 9779-320
Decision No. 517622

or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6.

If you have any questions, please contact Tony Kish by phone at 703 308-9443, or via email at kish.tony@epa.gov; or Craig Reeves by phone at 703 347-0486, or via email at reeves.craig@epa.gov

Sincerely,

A handwritten signature in black ink that reads "Tony Kish". The signature is written in a cursive style with a large, looped initial "T".

Tony Kish, Product Manager 22
Fungicide Branch
Registration Division (7505P)
Office of Pesticide Programs

Enclosure: Stamped Accepted Label

ACCEPTED

09/01/2016

Under the Federal Insecticide, Fungicide and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 9779-320

TERRANIL® 6L

Flowable Agricultural Fungicide

GROUP	M5	FUNGICIDE
-------	----	-----------

ACTIVE INGREDIENT

Chlorothalonil (tetrachloroisophthalonitrile)..... 54.0%

OTHER INGREDIENTS..... 46.0%

TOTAL 100.0%

Contains 6 pounds chlorothalonil per gallon.

KEEP OUT OF REACH OF CHILDREN

WARNING/AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand this label, find someone to explain it to you in detail.)

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING

Causes substantial but temporary eye injury. May be fatal if inhaled. Harmful if absorbed through skin. Avoid contact with skin, eyes, or clothing. Harmful if swallowed. Do not breathe vapor or spray mist.

FIRST AID	
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For additional information in case of medical emergency, call toll free 1-877-424-7452.	
SEE INSIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS	

EPA Reg. No. 9779-320

EPA Est. No. _____

Distributed By:
Winfield Solutions LLC
P.O. Box 64589
St. Paul, MN 55164-0589

NET CONTENTS
1/0517/6

PERSONAL PROTECTIVE EQUIPMENT (PPE)**WPS Uses**

Mixers, loaders, applicators and all other handlers must wear: - Long-sleeved shirt and long pants, - Chemical-resistant gloves made of any waterproof material – Category A (e.g., barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinyl chloride (PVC) or viton)- Shoes plus socks,- Protective eyewear, and

- A dust/mist filtering respirator (MSHA/NIOSH approval number prefix TC-21C), or a NIOSH approved respirator with any N, R, P or HE filter.

Non-WPS Uses

Applicators and other handlers who handle this pesticide for any use NOT covered by the Worker Protection Standard (40 CFR Part 170) – in general, only agricultural plant uses are covered – must wear:

- Long-sleeved shirt and long pants,
- Chemical-resistant gloves made of any waterproof material – Category A (e.g., barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinyl chloride (PVC) or viton),
- Shoes plus socks, and
- Protective eyewear.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Control Statements

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to aquatic invertebrates and wildlife. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean highwater mark. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate.

This chemical is known to leach through soil into groundwater under certain conditions as a result of label use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

This chemical can contaminate surface water through spray drift. Under some conditions, it may also have a high potential for runoff into surface water for several days to weeks after application. These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas overlaying extremely shallow groundwater, areas with in-field canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips, and areas overlying tile drainage systems that drain to surface water.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Do not apply when wind speed favors drift beyond the area intended for treatment.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard (WPS), 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, chemical-resistant gloves made of any waterproof material, shoes plus socks and protective eyewear.

Special Eye Irritation Provisions: This product is a severe eye irritant. Although the restricted entry interval expires after 12 hours, for the next 6.5 days entry is permitted only when the following safety measures are provided:

(1) At least one container designed specifically for flushing eyes must be available in operating condition at the WPS required decontamination site intended for workers entering the treated area.

(2) Workers must be informed in a manner they can understand:

- that residues in the treated area may be highly irritating to their eyes,
- that they should take precautions, such as refraining from rubbing their eyes, to keep the residues out of their eyes,
- that if they do get residues in their eyes, they should immediately flush their eyes using the eyeflush container that is located at the decontamination site or using other readily available clean water, and
- how to operate the eyeflush container.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

DO NOT enter or allow others to enter area until sprays have dried.

STORAGE AND DISPOSAL

DO NOT CONTAMINATE WATER, FOOD, OR FEED BY STORAGE OR DISPOSAL

Pesticide Storage: Store in a cool place. Protect from excessive heat.

Pesticide Disposal: Pesticide wastes are toxic. Improper disposal of excess pesticide, pesticide spray, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Disposal: Use label language appropriate for container size and type.

Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying.

Nonrefillable container equal to or less than 5 gallons. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Nonrefillable container greater than 5 gallons. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth

several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke. **Refillable container.** Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident, call
CHEMTREC 1-800-424-9300**

GENERAL INSTRUCTIONS AND INFORMATION

Application and Calibration Techniques for Sprinkler Irrigation

Apply this product only through the following types of irrigation systems. Do not apply this product through any other type of irrigation system. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from nonuniform distribution of treated water. If you have questions about calibration, you should contact State Experiment Station specialists, equipment manufacturers, or other experts. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

A. Center Pivot, Traveler, Big Gun, Motorized Lateral Move, End Tow, and Side (Wheel) Roll Irrigation Equipment: Operate system and injection equipment at normal pressures recommended by the manufacturer of injection equipment used. Fill tank of injection equipment with water. Operate system for one complete circle for center pivot or one complete run for the other recommended equipment, measuring time required, amount of water injected, and acreage contained in circle or run. Mix recommended amount of TERRANIL 6L for acreage to be covered into same amount of water used during calibration and inject into system continuously for one revolution or run, but continue to operate irrigation system until TERRANIL 6L has been cleared from last sprinkler head. Spray mixture in the chemical supply tank must be agitated at all times, otherwise settling and uneven application may occur.

B. Solid Set and Hand Move Irrigation Equipment: Determine acreage covered by sprinkler. Fill tank of injection equipment with water and adjust flow to use contents over a thirty to forty-five minute period. Mix desired amount of TERRANIL 6L for acreage to be covered into quantity of water used during calibration and operate entire system at normal pressures recommended by the manufacturer of injection equipment used for amount of time established during calibration. Provide constant mechanical agitation in the mix tank to insure that TERRANIL 6L will remain in suspension during the injection cycle. TERRANIL 6L can be injected at the beginning or end of the irrigation cycle or as a separate application. Stop injection equipment after treatment is completed and continue to operate irrigation system until the product is cleared from last sprinkler head.

Safety Devices

(1) The systems designated above must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow. (2) All pesticide injection pipelines must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. (3) The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. (4) The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. (5) The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely

affected. (6) Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock. (7) Do not apply when wind speed favors drift beyond the area intended for treatment.

Systems Connected to Public Water Sources

Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.

Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.

For additional instructions on safety precautions, refer to statements (2), (3), (4), (6), and (7) in the section on SAFETY DEVICES.

Spray Drift Precautions

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment-and-weather-related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions. The following drift management requirements must be followed to avoid off-target drift movement from aerial applications to agricultural field crops. These requirements do not apply to forestry applications, public health uses or to applications using dry formulations.

1. The distance of the outer most nozzles on the boom must not exceed $\frac{3}{4}$ the length of the wingspan or rotor.
2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.

Where states have more stringent regulations, they should be observed. The applicator should be familiar with and take into account the information covered in the **Aerial Drift Reduction Advisory Information**.

Aerial Drift Reduction Advisory Information

[This section is advisory in nature and does not supersede mandatory label requirements.]

Information on Droplet Size: The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see Wind, Temperature and Humidity, and Temperature Inversions).

Volume - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.

Pressure - Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.

Number of Nozzles - Use the minimum number of nozzles that provide uniform coverage.

Nozzle Orientation - Orienting nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential.

Nozzle Type - Use nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift.

Boom Length: For some use patterns, reducing the effective boom length to less than $\frac{3}{4}$ of the wingspan or rotor length may further reduce drift without reducing swath width.

Application Height: Applications should not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

Swath Adjustment: When applications are made with a crosswind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase, with increasing drift potential (higher wind, smaller drops, etc.).

Wind: Drift potential is lowest between wind speeds of 2-10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. **NOTE:** Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

Temperature and Humidity: When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

Temperature Inversions: Applications should not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

Sensitive Areas: The pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g. residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g. when wind is blowing away from the sensitive areas).

General Information

TERRANIL 6L is an excellent fungicide when used according to label directions for control of a broad spectrum of plant diseases. TERRANIL 6L can be used effectively in dilute or concentrate sprays. Thorough, uniform coverage is essential for disease control.

Slowly invert container several times to assure uniform mixture. The required amount of TERRANIL 6L should be added slowly into the spray tank during filling. With concentrate sprays, pre-mix the required amount of TERRANIL 6L in a clean container and add to the spray tank as it is being filled. Keep agitator running when filling spray tank and during spray operations.

Dosage rates on this label indicate pints of TERRANIL 6L per acre unless otherwise stated. Under conditions favoring disease development, the high rate specified and shortest application interval should be used. Applications should be made in sufficient water to obtain adequate coverage of foliage. Gallonage to be used will vary with crop and amount of plant growth. Spray volume usually will range from 20 to 150 gallons (approximately 80 to 600 liters) per acre for dilute sprays and 5 to 10 gallons (approximately 20 to 40 liters) per acre for concentrate ground sprays and aircraft applications. Both ground and aircraft methods of application are recommended unless specific directions for ground application only are given for a crop. Application through sprinkler irrigation systems is not recommended unless specific directions are given for a crop. See application and calibration instructions above.

This product may be tank mixed with other products at specified rates as long as tank mixing is not prohibited by the label(s) of the tank mix partner products, or as otherwise noted on this label, and the tank mix partner products are labeled for the timing and method of application for the use site to be treated. Do not combine TERRANIL 6L in the spray tank with pesticides, surfactants or fertilizers, unless prior use has shown the combination physically compatible, effective and noninjurious to your conditions of use. If compatibility with another product is not known, perform a (jar) test to determine compatibility. Follow all applicable directions, restrictions and precautions that appear on the respective tank mix product labels.

General Precautions and Restrictions

This product must not be applied within 150 feet (for aerial and air-blast applications) or 25 feet (for ground applications) of marine/estuarine water bodies unless there is an untreated buffer area of that width between the area to be treated and the water body.

Do not combine TERRANIL 6L with Dipel® 4L, Latron® AG-98, or Latron B-1956 as phytotoxicity may result from the combination when applied to some crops on this label.

Do not use on home lawns and turf sites associated with apartment buildings, daycare centers, playgrounds, playfields, recreational park athletic fields, athletic fields located on or next to schools (i.e., elementary, middle and high schools), campgrounds, churches, and theme parks.

Do not use on greenhouse grown food crops.

APPLICATION RATES

Dosage rates on this label indicate pints of TERRANIL 6L per acre, unless otherwise stated. Under conditions favoring disease development, the high rate specified and the shortest application interval should be used.

For each listed crop, the maximum total amount chlorothalonil active ingredient (lbs. a.i./A) which may be applied per acre of that crop (or crop group) during each growing season is stated in the Specific Use Restrictions section associated with each crop (or crop group). For each crop use situation listed below, the listed maximum individual and seasonal application rates must not be exceeded and the listed minimum retreatment intervals must not be decreased.

CROP	DISEASES (Pathogen)	PTS. PRODUCT/ A (lbs. a.i./A)	APPLICATION DIRECTIONS
Asparagus	Rust (<i>Puccinia asparagi</i>) Purple Spot (<i>Pleospora herbarum</i>) Cercospora blight (<i>C. asparagi</i>)	2 to 4 (1.5 to 3.0)	Use water volumes of 25-50 gallons per acre. Begin applications following final harvest of spears. Repeat applications at 14-28 day intervals (the minimum retreatment interval is 14 days), depending on disease pressure. Use the higher rate and shorter interval if disease severity begins to increase during the season or weather conditions are conducive for severe epidemics. Apply by ground.
Specific Use Restrictions: Do not apply more than 12 pints TERRANIL 6L (9.0 lbs. a.i.) per acre during each growing season. Do not apply within 190 days (120 days in CA and AZ) of the harvest of spears in the following season.			

CROP	DISEASES (Pathogen)	PTS. PRODUCT/ A (lbs. a.i./A)	APPLICATION DIRECTIONS
Bean (Snap)	Rust (<i>Uromyces appendiculatus</i>)	1 $\frac{3}{8}$ to 3 (1.0 to 2.25)	Use in sufficient water to obtain adequate coverage. Begin applications during early bloom stage or when disease first threatens and repeat (the minimum retreatment interval is 7 days).
	Botrytis blight (gray mold) (<i>B. cinerea</i>)	3 (2.25)	
Specific Use Restrictions: Do not apply more than 12 pints TERRANIL 6L (9.0 lbs. a.i.) per acre during each growing season. Do not apply within 7 days of harvest.			
Dried shelled pea and bean (except soybeans) bean, adzuki bean, broad bean, dry bean, lablab bean, navy bean, kidney bean, lima bean, moth bean, mung bean, pink bean, pinto bean, tepary bean, urd bean, yardlong catjang chickpea (garbanzo) cowpea lupin, grain lupin bean, rice bean, runner bean, jackbean pea, blackeyed pea, southern	Rust (<i>Uromyces appendiculatus</i>) Anthracnose (<i>Colletotrichum lindemuthianum</i>) Downy mildew (<i>Phytophthora nicotianae</i>) Cercospora leaf blotch (<i>C. cruenta</i>) Ascochyta blight (<i>A. phaseolorum</i>)	1 $\frac{3}{8}$ to 2 (1.0 to 1.5)	Use in sufficient water to obtain adequate coverage. Begin applications at first onset of disease, which may occur as early as 2 to 4 weeks before flowering. Repeat applications at 7 to 10 day intervals (the minimum retreatment interval is 7 days). For use only on beans to be harvested dry with pods removed. Apply by ground, air or chemigation.
Specific Use Restrictions: Do not apply more than 8 pints TERRANIL 6L (6 lbs. a.i.) per acre during each growing season. Do not apply within 14 days before harvest.			

CROP	DISEASES (Pathogen)	PTS. PRODUCT/ A (lbs. a.i./A)	APPLICATION DIRECTIONS
Blueberries	Suppression: Anthracnose (ripe rot) <i>(C. gloeosporoides)</i> Mummy Berry <i>(M. vaccinicorymbosi)</i>	3 to 4 (2.25 to 3.0)	TERRANIL 6L should be integrated into an overall disease management strategy which includes alternation with a fungicide with a different mode of action. Diseases may only be suppressed and russetting may occur under heavy disease pressure or unfavorable environmental conditions. Apply in sufficient water to obtain adequate coverage, normally 20-100 gallons per acre. Begin applications at budbreak (green tip) and repeat at 10-day intervals through early bloom (the minimum retreatment interval is 10 days). Under heavy disease pressure, use the higher rate. Apply by ground or air.
	Septoria leaf spot <i>(Septoria albopunctata)</i> Rust <i>(Pucciniastrum vaccinii)</i>	3 to 4 (2.25 to 3.0)	Foliar Use After Harvest (after all berries are harvested): To maintain healthy leaves for the following season, apply in sufficient water to obtain adequate coverage (normally 20-100 gallons per acre). Repeat at 10-14 day intervals (the minimum retreatment interval is 10 days). Apply by ground or air.
Specific Use Restrictions: Do not apply more than 12 pints TERRANIL 6L (9.0 lbs. a.i.) per acre during each growing season. Do not apply after full bloom (except for foliar use after harvest) or within 42 days of harvest.			
Brassica, head and Stem: Broccoli Broccoli, Chinese Brussels Sprouts Cabbage Cabbage, Chinese (tight-headed varieties and Napa) Cabbage, Chinese Mustard Cauliflower Cavalo, broccoli Kohlrabi	Alternaria leaf spot, <i>(Alternaria spp.)</i> Downy mildew <i>(Peronospora parasitica)</i>	1½ (1.125)	Use in sufficient water to obtain adequate coverage. Begin applications after transplants are set in field, or shortly after emergence of field-seeded crop, or when conditions favor disease development. Repeat at 7 to 10 day intervals (the minimum retreatment interval is 7 days). Apply by ground, air or chemigation.
	Ring spot (California only)	2 (1.5)	For field-seeded Brussels sprouts, begin applications at time of early sprout development or when conditions favor disease development. Repeat at 7 to 10 day intervals (the minimum retreatment interval is 7 days) to maintain control.
Specific Use Restrictions: Do not apply more than 11.7 pints TERRANIL 6L (8.8 lbs. a.i.) per acre during each growing season. Do not apply within 7 days of harvest.			

CROP	DISEASES (Pathogen)	PTS. PRODUCT/ A (lbs. a.i./A)	APPLICATION DIRECTIONS
Carrot	Cercospora leaf spot (<i>C. carotae</i>) Alternaria leaf blight (<i>A. dauci</i>)	1½ to 2 (1.125 to 1.5)	Use in sufficient water to obtain adequate coverage. Start applications when disease threatens and repeat at 7 to 10 day intervals (the minimum retreatment interval is 7 days) to maintain control. Apply by ground, air or chemigation.
Specific Use Restrictions: Do not apply more than 20 pints TERRANIL 6L (15 lbs. a.i.) per acre during each growing season. TERRANIL 6L may be applied the day of harvest.			
Celery	Early blight (<i>Cercospora apii</i>) Late blight (<i>Septoria apicola</i>) Basal stalk rot (<i>Rhizoctonia solani</i>)	2 to 3 (1.5 to 2.25)	Use in sufficient water to obtain adequate coverage. Start applications when transplants are set in the field and repeat at a 7-day interval as needed to maintain control (the minimum retreatment interval is 7 days). Apply by ground, air or chemigation.
	Suppression (7-day schedule): Pink rot (<i>Sclerotinia sclerotiorum</i>)	3 (2.25)	
	Early blight (<i>Cercospora apii</i>) Late blight (<i>Septoria apicola</i>)	1½ to 2 (1.125 to 1.5) per 100 gals.	
Specific Use Restrictions: Do not apply more than 24 pints TERRANIL 6L (18 lbs. a.i.) per acre during each growing season. Do not apply within 7 days of harvest.			
Corn (Sweet) Corn (grown for seed)	Helminthosporium leaf blights Rust (<i>Puccinia</i> spp.)	¾ to 2 (0.6 to 1.5)	Use in sufficient water to obtain adequate coverage. Begin applications when conditions favor disease development and repeat at a 7 day interval as required to maintain control (the minimum retreatment interval is 7 days). Under severe disease conditions, use 1½ to 2 pints TERRANIL 6L per acre. Apply by ground, air or chemigation.
Specific Use Restrictions: Do not apply more than 12 pints TERRANIL 6L (9 lbs. a.i.) per acre during each growing season. Do not apply within 14 days of harvest. Do not apply to sweet corn to be processed. Do not allow livestock to graze in treated fields. Do not ensile treated corn or use as livestock forage.			

CROP	DISEASES (Pathogen)	PTS. PRODUCT/ A (lbs. a.i./A)	APPLICATION DIRECTIONS
Cranberry	Fruit rots Lophodermium leaf/twig blight (<i>L. hypophyllum</i>)	4 to 6½ (3.0 to 4.9)	Apply at early bloom and repeat at 10 to 14 day intervals (the minimum retreatment interval is 10 days). Under severe disease conditions, use the 6½ pint/acre rate on a 10 day schedule. Apply by ground, air or chemigation. When applying by chemigation, use 300 gallons of water per acre through solid set systems only.
	Upright Dieback (<i>Phomopsis vaccinii</i>)	4 to 6½ (3.0 to 4.9)	Apply in sufficient water to obtain coverage of uprights and runners. Make the first application before bloom, at the time shoots begin growth in the spring. Make additional applications at 10-14 day intervals. Apply by ground, air or chemigation. When applying by chemigation, use 300 gallons of water per acre through solid set systems only.

Specific Use Restrictions: Do not apply more than 20 pints TERRANIL 6L (15 lbs. a.i.) per acre during each growing season. Do not apply within 50 days of harvest. Do not apply to beds when flooded or allow release of irrigation water from beds for at least 3 days following application.

Cucurbits: Cantaloupe Chinese waxgourd (Chinese preserving melon) Cucumber Gourds Honeydew melon <i>Momordica</i> spp. (balsam apple, bitter melon) Muskmelon Pumpkin Squash Watermelon Zucchini	Anthracnose (<i>Colletotrichum</i> spp.) Downy mildew (<i>Pseudoperonospora cubensis</i>) Target spot (<i>Corynespora cassiicola</i>)	1½ to 2 (1.125 to 1.5)	Use in sufficient water to obtain adequate coverage. Begin applications when plants are in first true leaf stage or when conditions are favorable for disease development. Repeat applications at 7 day intervals (the minimum retreatment interval is 7 days). Note: Spraying mature watermelons may result in sunburn of the upper surface of the fruit. Do not apply TERRANIL 6L to watermelons when any of the following conditions are present: <ol style="list-style-type: none"> 1. Intense heat and sunlight 2. Drought conditions 3. Poor vine canopy 4. Other crop and environmental conditions which may be conducive to increased natural sunburn.
Including cultivars and/or hybrids of these.	Cercospora leaf spot (<i>C. citrullina</i>) Gummy stem blight/vine decline (<i>Didymella bryoniae</i>) Alternaria leaf blight (<i>A. cucumerina</i>) Alternaria leaf spot (<i>A. alternata</i>) Scab (<i>Cladosporium cucumerinum</i>) Powdery mildew (<i>Sphaerotheca</i> only)	2 to 3 (1.5 to 2.25)	Do not combine TERRANIL 6L with anything except water for application to watermelons unless your prior use has shown the combination to be non-injurious to watermelons under your conditions of use. Apply by ground, air or chemigation.

Specific Use Restrictions: Do not apply more than 21 pints TERRANIL 6L (15.75 lbs. a.i.) per acre during each growing season. TERRANIL 6L may be applied the day of harvest.

CROP	DISEASES (Pathogen)	PTS. PRODUCT/ A (lbs. a.i./A)	APPLICATION DIRECTIONS
Fruiting Vegetables (except tomatoes): Eggplant Groundcherry Okra Pepino Pepper (includes bell pepper, chili pepper, cooking pepper, pimento, sweet pepper) Tomatillo	Anthracnose (<i>Colletotrichum</i> spp.) Botrytis Leaf mold (<i>Botrytis cinera</i>) Cercospora leaf spot (<i>Cercospora</i> spp) Powdery mildew (<i>Leveillula taurica</i>)	$1\frac{1}{2}$ (1.125)	Use in sufficient water to obtain adequate coverage. Begin application when disease first threatens, and repeat at 7 to 10 day intervals as disease pressure warrants. Apply by ground, air or chemigation
Specific Use Restrictions: Do not apply more than 12 pints of TERRANIL 6L (9.0 lbs. a.i.) per acre during each growing season. Do not apply within 3 days of harvest.			
Ginseng	Alternaria blight (<i>Alternaria panax</i>) Grey mold (<i>Botrytis cinerea</i>)	2 (1.5)	Use in sufficient water to obtain adequate coverage. Begin applications when disease first threatens, and repeat at 7 to 10 day intervals as disease pressure warrants.
Specific Use Restrictions: Do not apply more than 16 pints of TERRANIL 6L (12.0 lbs. a.i.) per acre during each growing season. Do not apply within 14 days of harvest.			
Grasses Grown for Seed	Stem rust Leaf rust Stripe rust Septoria leaf spot Glume blotch Bipolaris and Drechslera leaf spots Selenophoma (eyespot)	1 to $1\frac{1}{2}$ (0.75 to 1.125) 1 to 2 (0.75 to 1.5)	Use in sufficient water to obtain adequate coverage. Begin applications during stem elongation when conditions favor disease development. Re-apply at flag (top) leaf emergence and repeat applications at 14 day intervals (the minimum retreatment interval is 14 days). Apply by ground, air or chemigation.
Specific Use Restrictions: Do not apply more than 6 pints TERRANIL 6L (4.5 lbs. a.i.) per acre during each growing season. Do not apply within 14 days of harvest. Do not allow livestock to graze in treated areas or feed hay produced before harvest. Feeding of treated plant parts after harvest of seed is allowed.			
Horseradish	Ramularia stem and leaf spot (<i>Ramularia armoraciae</i>)	3 (2.25)	Use is sufficient water to obtain adequate coverage. Begin applications when disease first threatens, and repeat at 7 to 10 day intervals as disease pressure warrants.
Specific Use Restrictions: Do not apply more than 24 pints of TERRANIL 6L (18 lbs. a.i.) per acre during each growing season. Do not apply within 14 days of harvest.			
Lupine, Lentil	Anthracnose (<i>Colletotrichum gloeosporioides</i>) Ascochyta (<i>Ascochyta pisi</i>)	1 to 1.5 (0.75 to 1.125)	Use in sufficient water to obtain adequate coverage. Begin applications when disease first threatens, and repeat at 7 to 10 day intervals as disease pressure warrants.
Specific Use Restrictions: Do not apply more than 8 pints of TERRANIL 6L (6 lbs. a.i.) per acre during each growing season. Do not apply within 14 days of harvest.			

CROP	DISEASES (Pathogen)	PTS. PRODUCT/ A (lbs. a.i./A)	APPLICATION DIRECTIONS
Mango	Anthracnose (<i>Colletotrichum</i> spp.)	2 to 3½ (1.5 to 2.6)	Use a water volume of 20 to 300 gallons per acre. Begin applications at early bloom and repeat on a 7-14 day interval until early fruit development. Begin the season with the 2 pint rate on a 14-day interval (the minimum retreatment interval is 7 days). If disease pressure is severe, use the higher rate and shorter interval. Apply by ground or air.
Specific Use Restrictions: Do not apply more than 32 pints TERRANIL 6L (24 lbs. a.i.) per acre during each growing season. Do not apply within 21 days of harvest.			
Mint (Indiana, Michigan and Wisconsin only)	Rust (<i>Puccinia menthae</i>) Septoria leaf spot (<i>S. menthae</i>)	1⅜ (1.0)	Use in sufficient water to obtain adequate coverage, normally 20 to 150 gallons per acre for dilute sprays and 5 to 10 gallons per acre for concentrate ground and aircraft applications. Begin applications when emerging plants are 4-8 inches high. Repeat applications at 7 to 10 day intervals to maintain control (the minimum retreatment interval is 7 days).
Specific Use Restrictions: Do not apply more than 4 pints TERRANIL 6L (3 lbs. a.i.) per acre during each growing season. Do not apply within 80 days of harvest. Do not feed fresh or extracted mint hay from treated fields to livestock.			
Mushrooms	Verticillium brown spot and dry bubble (<i>Verticillium fungicola</i>)	Rate per 1,000 sq. ft. of bed surface 2.75 to 5.5 fl. oz.	Apply as a drench to the mushroom bed surface in at least 12.5 gallons of water per 1000 sq. ft. of mushroom bed. Make two applications as follows: <ul style="list-style-type: none"> • First application- apply 5.5 fl. oz. of this product within two days of top-dressing the spawn-colonized mushroom compost with a casing layer. • Second application- apply 2.75 fl. oz. of this product at pinning.
Specific Use Restrictions: Make no more than two applications per cropping cycle. Do not apply more than 8.25 fl. oz. of this product per cropping cycle. Do not apply within 5 days of first harvest.			

CROP	DISEASES (Pathogen)	PTS. PRODUCT/ A (lbs. a.i./A)	APPLICATION DIRECTIONS												
Onion (dry bulb) and Garlic	Botrytis leaf blight -(<i>Botrytis</i> spp.) Purple blotch (<i>Alternaria porri</i>) Suppression: Botrytis neck rot Downy mildew (<i>Peronospora destructor</i>)	1 to 3 (0.75 to 2.25)	Apply in sufficient water to obtain adequate thorough coverage of tops. TERRANIL 6L is recommended for use with disease monitoring systems which adjust fungicide rates and frequency of application according to disease hazard. Apply as follows: <table border="1" data-bbox="1036 499 1515 810"> <thead> <tr> <th></th> <th>Low Disease Hazard & Prior to Infection</th> <th>Low Disease Hazard & Some Disease Present</th> <th>High Disease Hazard</th> </tr> </thead> <tbody> <tr> <td>Rate/Acre</td> <td>1 pt.</td> <td>1½ pts.</td> <td>3 pts.</td> </tr> <tr> <td>Frequency</td> <td>10 days</td> <td>7-10 days</td> <td>7 days</td> </tr> </tbody> </table> For suppression of neck rot (<i>Botrytis</i> spp.) during storage, a minimum of three weekly applications prior to lifting, using 1½ to 3 pints of TERRANIL 6L per acre, is recommended The minimum retreatment interval is 7 days. Apply by ground, air or chemigation.		Low Disease Hazard & Prior to Infection	Low Disease Hazard & Some Disease Present	High Disease Hazard	Rate/Acre	1 pt.	1½ pts.	3 pts.	Frequency	10 days	7-10 days	7 days
	Low Disease Hazard & Prior to Infection	Low Disease Hazard & Some Disease Present	High Disease Hazard												
Rate/Acre	1 pt.	1½ pts.	3 pts.												
Frequency	10 days	7-10 days	7 days												
Specific Use Restrictions: Do not apply more than 20 pints TERRANIL 6L (15 lbs. a.i.) per acre during each growing season. Do not apply within 7 days of harvest.															
Onion (green bunching) Leek, Shallots, Onion and Garlic (grown for seed)	Botrytis leaf blight (<i>Botrytis</i> spp.) Purple blotch (<i>Alternaria porri</i>) Suppression: Downy mildew (<i>Peronospora destructor</i>)	1½ to 3 (1.125 to 2.25)	Use in sufficient water to obtain thorough coverage of tops. Begin applications prior to favorable infection periods, and repeat at 7 to 10 day intervals for as long as conditions favor disease (the minimum retreatment interval is 7 days). Use the high rate and a 7 day schedule of applications when heavy dew or rain persist. Apply by ground, air or chemigation.												
Specific Use Restrictions: Do not apply more than 9 pints TERRANIL 6L (6.75 lbs. a.i.) per acre during each growing season. Do not apply within 7 days of harvest on garlic. Do not apply within 14 days of harvest on green bunching onions, leeks or shallots.															
Papaya	Alternaria fruit spot (<i>A. alternata</i>) Anthracnose (<i>Colletotrichum</i> spp.) Stem end rot (<i>A. alternata</i> , <i>Colletotrichum</i> spp.)	1½ to 3 (1.125 to 2.25)	Apply with ground equipment only, in sufficient water to obtain adequate coverage of fruit and leaves. Begin treatment when conditions favor development of disease and continue treatments at 14 day intervals until weather conditions no longer favor disease development (the minimum retreatment interval is 14 days.).												
Specific Use Restrictions: Do not apply more than 9 pints TERRANIL 6L (6.75 lbs. a.i.) per acre during each growing season. TERRANIL 6L may be applied the day of harvest.															

CROP	DISEASES (Pathogen)	PTS. PRODUCT/ A (lbs. a.i./A)	APPLICATION DIRECTIONS
Parsnip	Alternaria leaf spot (<i>Alternaria</i> spp.) Downy mildew (<i>Plasmopara crustosa</i>) Anthracnose (<i>Colletotrichum</i> spp.) Botrytis blight (gray mold) (<i>B. cinerea</i>) Bottom rot (<i>Rhizoctonia</i>)	1½ to 2 (1.125 to 1.5)	Apply in sufficient water to obtain adequate coverage. Make the first application at the first sign of disease or when conditions are favorable for infection. Continue applications on a 7 to 10 day schedule (the minimum retreatment interval is 7 days).
Specific Use Restrictions: Do not apply more than 8 pints TERRANIL 6L (6 lbs. a.i.) per acre during each growing season. Do not apply within 10 days of harvest.			
Passion Fruit (Hawaii only)	Alternaria fruit and leaf spot (<i>Alternaria</i> spp.) Anthracnose (<i>Colletotrichum</i> spp.) Cercospora fruit spot	2 (1.5)	Apply with ground equipment in sufficient water to obtain adequate coverage of fruit and leaves. Begin applications during late bloom and repeat at 14 day intervals until weather conditions no longer favor disease development (the minimum retreatment interval is 14 days).
Specific Use Restrictions: Do not apply more than 10 pints TERRANIL 6L (7.5 lbs. a.i.) per acre during each growing season. Do not apply within 7 days of harvest.			
Peanut	Early leaf spot (<i>Cercospora arachidicola</i>) Late leaf spot (<i>Cercosporidium personatum</i>) Pepper spot (<i>Leptosphaerulina crassiasca</i>) Rust (<i>Puccinia arachidis</i>) Web blotch (<i>Phoma arachidicola</i>)	1 to 1½ (0.75 to 1.125)	Apply in sufficient water for coverage when leaf wetness first occurs or 30 to 40 days after planting. Repeat at 14 day intervals (the minimum retreatment interval is 14 days). When conditions favor late leaf spot or when rust or web blotch occur, apply 1½ pints TERRANIL 6L per acre at 14 day intervals for the remainder of the season. Apply by ground, air, or chemigation. If applying by chemigation, use 1½ pints TERRANIL 6L per acre. It is recommended to alternate chemigation applications with ground or aerial applications.
Specific Use Restrictions: Do not apply more than 12 pints TERRANIL 6L (9 lbs. a.i.) per acre during each growing season. Do not apply within 14 days of harvest. Do not allow livestock to graze in treated areas. Do not feed hay or threshings from treated fields to livestock.			
Persimmon (Florida and Hawaii only)	Cercospora leaf spot (<i>Cercospora fuliginosa</i>)	1.25 (0.94)	Use in sufficient water to obtain adequate coverage. Begin applications when disease first threatens, and repeat at 14 day intervals as disease pressure warrants. Aerial applications require the use of a minimum of 10 gallons per acre.
Specific Use Restrictions: Do not apply more than 6.25 pints of TERRANIL 6L (4.7 lbs. a.i.) per acre during each growing season. Do not apply within 14 days of harvest.			

CROP	DISEASES (Pathogen)	PTS. PRODUCT/ A (lbs. a.i./A)	APPLICATION DIRECTIONS
Potato	Late blight (<i>Phytophthora infestans</i>) Early blight (<i>Alternaria solani</i>) Botrytis vine rot (<i>B. cinerea</i>) Black dot (<i>Colletotrichum coccodes</i>)	$\frac{3}{4}$ (0.6) -then- 1 to 1½ (0.75 to 1.125)	<p>Begin applications at the low rate when vines are first exposed and leaf wetness occurs. Repeat applications at 5 to 10 day intervals (the minimum retreatment interval is 5 days).</p> <p>Begin applying the higher label rates at 5 to 10 day intervals when any one of the following events occurs:</p> <ul style="list-style-type: none"> - Vines close within the row; - Late blight forecasting measures 18 disease severity values (DSV); - The crop reaches 300 P-days. <p>Increase water spray volume as canopy density increases. Use the highest rate and shortest interval when plants are rapidly growing and disease conditions are severe.</p> <p>Apply by ground, air, or chemigation. Do not exceed a 10 day interval between applications when using chemigation.</p>
<p>Specific Use Restrictions: Do not apply more than 15 pints of TERRANIL 6L (11.25 lbs. a.i.) per acre during each growing season. Do not apply within 7 days of harvest.</p>			
Rhubarb	Ramularia leaf spot (<i>Ramularia rhei</i>) Ascochyta blight (<i>Ascochyta rhei</i>)	3 (2.25)	<p>Use sufficient water to obtain adequate coverage. Begin applications when disease first threatens, and repeat at 7 to 10 day intervals as disease pressure warrants.</p>
<p>Specific Use Restrictions: Do not apply more than 18 pints of TERRANIL 6L (13.5 lbs. a.i.) per acre during each growing season. Do not apply within 30 days of harvest.</p>			

CROP	DISEASES (Pathogen)	PTS. PRODUCT/ A (lbs. a.i./A)	APPLICATION DIRECTIONS
Soybean	Anthracnose (<i>Colletotrichum truncatum</i>) Diaporthe pod and stem rot (<i>D. phaseolorum</i>) Frogeye leaf spot (<i>Cercospora sojina</i>)		Apply in sufficient water to obtain complete coverage, using at least five gallons water per acre for aerial application. Use the three application program in areas having a history of moderate to severe disease intensity. The minimum retreatment interval is 14 days. Apply by ground, air, or chemigation.
	Purple seed stain (<i>C. kikuchii</i>) Cercospora leaf blight (<i>C. kikuchii</i>), Septoria brown spot (<i>S. glycines</i>)	1½ to 2¼ (1.125 to 1.7)	Two application program – For determinate varieties, make the first application at R3 stage (early pod set) and the second application at R5 (seed formation). For indeterminate varieties, make the first application when largest pods are 1-1¼ inches in length. Make the second application 14 days later.
	Suppression: Rust (<i>Phakopsora pachyrhizi</i>)	1 to 2 (0.75 to 1.5)	Three application program: For determinate varieties, make the first application at the beginning of flowering (R1), the second at early pod set (R3), and the third at beginning of seed formation (R5). For indeterminate varieties, make the first application one week after first flowering and continue applications at 14 day intervals.
	Stem canker (<i>Diaporthe phaseolorum</i>)	1 (0.75)	Apply in 10 to 20 gallons of water per acre, as a band treatment directing spray to provide coverage of entire plant. Make the first application at time of emergence of the second trifoliolate leaves (V2). If conditions favor stem canker disease, make a second and a third application. Make all applications at 14 day intervals.
Specific Use Restrictions: Do not apply more than 6 pints TERRANIL 6L (4.5 lbs. a.i.) per acre during each growing season. Do not apply within 6 weeks of harvest. Do not feed hay or threshings from treated fields to livestock.			

CROP	DISEASES (Pathogen)	PTS. PRODUCT/ A (lbs. a.i./A)	APPLICATION DIRECTIONS
Tomato	FOLIAGE: Early blight <i>(Alternaria solani)</i> Late blight <i>(Phytophthora infestans)</i> Gray leaf spot <i>(Stemphyllium botryosum)</i> Gray leaf mold <i>(Fluvia fluva;</i> <i>Cladosporium)</i> Septoria leaf spot <i>(S. lycopersici)</i> Target spot <i>(Corynespora cassicola)</i>	1 $\frac{3}{8}$ to 2 (1.0 to 1.5)	Apply in sufficient water to obtain adequate coverage. Begin applications when dew or rain occur and disease threatens. Apply on a 7-10 day interval for foliage diseases. For fruit diseases, begin at fruit set and apply on a 7-14 day interval. Use the highest rate and shortest interval specified when disease conditions are severe. The minimum retreatment interval is 7 days. Apply by ground, air, or chemigation.
	FRUIT: Anthracnose <i>(Colletotrichum spp.)</i> Alternaria fruit rot (black mold) <i>(A. alternata)</i> Botrytis gray mold <i>(B. cinerea)</i> Late blight fruit rot <i>(P. infestans)</i> Rhizoctonia fruit rot <i>(R. solani)</i>	2 to 2 $\frac{3}{4}$ (1.5 to 2.1)	
Specific Use Restrictions: Do not apply more than 20 pints TERRANIL 6L (15 lbs. a.i.) per acre during each growing season. TERRANIL 6L may be applied the day of harvest.			
Yam	Anthracnose <i>(Colletotrichum gloeosporioides)</i>	1 to 1.25 (0.75 to 1.125)	Use in sufficient water to obtain adequate coverage. Begin applications when disease first threatens, and repeat at 10 to 14 day intervals as disease pressure warrants.
Specific Use Restrictions: Do not apply more than 15 pints of TERRANIL 6L (11.25 lbs. a.i.) per acre during each growing season. Do not apply within 7 days of harvest.			

Tree and Orchard Crops

Apply TERRANIL 6L in sufficient water and with proper calibration to obtain uniform coverage of tree canopy. For fruit and nut bearing crops, the maximum volume is 300 gallons per acre unless indicated otherwise in the specific use directions. For conifers, the maximum volume is 100 gallons per acre.

Application with ground equipment is preferable to aerial application because ground applications generally give better coverage of the tree canopy. If application with ground equipment is not feasible, TERRANIL 6L may be applied with aircraft

using at least 20 gallons of spray per acre. The minimum volume for application by aircraft to conifer stands and Christmas trees is 10 gallons per acre.

When concentrate sprays are used or when treating non-bearing or immature trees, the lower rate of TERRANIL 6L listed may be used. Do not allow livestock to graze in treated areas.

		PTS. PRODUCT PER (lbs. a.i. per)		
CROP	DISEASES (Pathogen)	ACRE	100 GAL.*	APPLICATION DIRECTIONS
Almonds	Anthracnose (<i>Colletotrichum acutatum</i>) Blossom blight/brown rot (<i>Monilinia</i> spp.) Shot hole (<i>Wilsonomyces carpophilus</i>) Scab (<i>Venturia carpophila</i>)	4 (3.0)	1.33 (1.0)	Use water volumes of 20-300 gallons per acre. For blossom blight, begin application at popcorn (pink bud) and follow with an application at full bloom. If weather is still conducive for disease development, another application may be made at petal fall. For control of shot hole, make an application in the autumn at leaf fall. In the spring, make the first application at budbreak, followed by an application at shuck split to control nut infections and to control scab. Apply by ground or air.
Specific Use Restrictions: Do not apply more than 25 pints TERRANIL 6L (18.75 lbs. a.i.) per acre during each growing season (leaf fall through shuck split). Do not apply within 150 days of harvest.				
Filberts (Hazel-nuts)	Eastern filbert blight (<i>Anisogramma anomala</i>)	4 (3.0)	1.33 (1.0)	Use a water volume of 20 to 300 gallons per acre. Begin applications at the onset of disease or when weather conditions favor disease development. Make applications on a 14-28 day schedule, using the shorter interval under heavy disease pressure (the minimum retreatment interval is 14 days.)
Specific Use Restrictions: Do not apply more than 12 pints TERRANIL 6L (9 lbs. a.i.) per acre during each growing season. Do not apply within 120 days of harvest. Do not apply through irrigation. Do not apply with oils, other pesticides, surfactants or fertilizers. Do not apply within one week of an oil-based pesticide application.				

		PTS. PRODUCT PER (lbs. a.i. per)		
CROP	DISEASES (Pathogen)	ACRE	100 GAL.*	APPLICATION DIRECTIONS
Peach Nectarine Apricot Cherry Plum Prune	Leaf curl (<i>Taphrina deformans</i>) Shot hole (<i>Wilsonomyces carpophilus</i>)	3 $\frac{1}{8}$ to 4 $\frac{1}{8}$ (2.3 to 3.1)	1 to 1 $\frac{3}{8}$ (0.75 to 1.0)	For best control of both diseases apply at leaf fall in late autumn, using sufficient water and proper sprayer calibration to obtain uniform coverage. When conditions favor high disease levels, use the high rate of application and apply once or twice more in mid to late winter before bud swell. If the leaf fall application is not practical, application of TERRANIL 6L for control of leaf curl may be made at any time prior to bud swell the following spring. Where shot hole occurs, also apply at budbreak to protect newly emerging leaves and at shuck split to prevent fruit infections. Apply by ground or air.
	Lacy (russet) scab (plum/prune) Brown rot blossom blight (<i>Monilinia</i> spp.)	3 $\frac{1}{8}$ to 4 $\frac{1}{8}$ (2.3 to 3.1)	1 to 1 $\frac{3}{8}$ (0.75 to 1.0)	Make one application at popcorn (pink, red, or early white bud) and a second application at full bloom. If weather conditions favor disease development, make an additional application at petal fall.
	Cherry leaf spot (<i>Blumeriella jaapii</i>) Scab (<i>Cladosporium carpophilum</i>) Black knot (cherry, plum) (<i>Apiosporina morbosa</i>)	3 $\frac{1}{8}$ to 4 $\frac{1}{8}$ (2.3 to 3.1)	1 to 1 $\frac{3}{8}$ (0.75 to 1.0)	In addition to the bloom applications listed above, make one application at shuck-split. Do not apply TERRANIL 6L after shuck-split and before harvest. If additional disease control is needed before harvest, use another registered fungicide. For control of cherry leaf spot after harvest, make one application to foliage within 7 days after fruit is removed. In orchards with a history of high leaf spot incidence, make a second application 10-14 days later. Apply by ground or air.
Specific Use Restrictions: Do not apply more than 20 $\frac{1}{2}$ pints TERRANIL 6L (15.4 lbs. a.i.) per acre during each growing season. TERRANIL 6L may be applied the day of harvest. The minimum retreatment interval is 10 days.				
Pistachio	Botryosphaeria blight (<i>B. dothidea</i>) Suppression: Alternaria late blight (<i>A. alternata</i>)	6 (4.5)	3 (2.25)	Use a water volume of 20 to 200 gallons per acre. Make the first application at the beginning of the blossom period followed by an application at full bloom. Make additional applications as required on a 28-day schedule. (The minimum retreatment interval is 28 days). For Septoria and Botrytis, use the higher rate if disease pressure is severe.
	Septoria leaf spot (<i>S. pistacina</i>) Botrytis blight (<i>B. cinerea</i>)	4 to 6 (3.0 to 4.5)	2 to 3 (1.50 to 2.25)	NOTE: Use of this product may result in speckling or reddening of the fruit hull (epicarp). This effect is superficial and has not resulted in any change in nut quality. Apply by ground or air.
Specific Use Restrictions: Do not apply more than 30 pints TERRANIL 6L (22.5 lbs. a.i.) per acre during each growing season. Do not apply within 14 days of harvest.				

		PTS. PRODUCT PER (lbs. a.i. per)		
CROP	DISEASES (Pathogen)	ACRE	100 GAL.*	APPLICATION DIRECTIONS
Conifers (pines, spruces)	Swiss needlecast (<i>Phaeocryptopus gaeumannii</i>)	2¾ to 5½ (2.1 to 4.125)	2¾ to 5½ (2.1 to 4.125)	Single application technique: In Christmas tree plantations or conifer stands, make one application in the spring when new shoot growth is ½ to 2 inches in length. Make the first application in spring when new shoot growth is ½ to 2 inches in length. Make additional applications at 3 to 4 week intervals until conditions no longer favor disease development. For use in nursery beds, apply the highest rate specified on a 3 week schedule.
	Scleroderris canker (pines) (<i>Gremmeniella abietina</i>) Swiss needlecast (<i>P. gaeumannii</i>)	1½ to 2¾ (1.125 to 2.1)	1½ to 2¾ (1.125 to 2.1)	
	Sirococcus tip blight (<i>S. conigenus</i>)	2 to 3½ (1.5 to 2.6)	2 to 3½ (1.5 to 2.6)	
	Rhizosphaera needlecast (spruces) (<i>Rhizosphaera</i> spp.) Scirrhia brown spot (pines) (<i>Mycosphaerella dearnessii</i>)	5½ (4.125)	5½ (4.125)	
	Cyclaneusma and Lophodermium needlecast (pines)	2¾ to 5½ (2.1 to 4.125)	2¾ to 5½ (2.1 to 4.125)	Apply in early spring prior to budbreak. Repeat applications at approximately 6 to 8 week intervals, until spore release ceases in late fall. Apply monthly during periods of frequent rainfall, and where Lophodermium infections occur during dormancy (Pacific NW). During drought periods, applications may be suspended, then resumed upon next occurrence of needle wetness.
	Rhabdocline needlecast (Douglas fir)	1½ to 2¾ (1.125 to 2.1)	1½ to 2¾ (1.125 to 2.1)	Apply at budbreak and repeat at 3 to 4 week intervals until needles are fully elongated and conditions no longer favor disease development. In plantations of mixed provenance, or when irregular budbreak occurs, apply weekly until all trees have broken bud, then every 3 to 4 weeks as specified above. In nursery beds, use the high rate on a 3 week schedule.
	Botrytis seedling blight Phoma twig blight	1½ to 2¾ (1.125 to 2.1)	1½ to 2¾ (1.125 to 2.1)	Begin applications in nursery beds when seedlings are 4 inches tall and when cool, moist conditions favor disease development. Make additional applications at 7 to 14 day intervals as long as favorable disease conditions persist.
	Autoecious needle rust (Weir's cushion) (spruce)	5½ (4.125)	5½ (4.125)	Begin applications when 10% of buds have broken and twice thereafter at 7-10 day intervals.

Specific Use Restrictions: Do not apply more than 22 pints TERRANIL 6L (16.5 lbs. a.i.) per acre during each growing season. The minimum retreatment interval for established trees is 21 days. The minimum retreatment interval in nursery beds is 7 days. Do not use on forest.

*Volumetric rates to be used only with full dilute spray volume specified on this label for tree and orchard crops.

DIRECTIONS FOR USE ON TURF AND ORNAMENTALS

Turf

Group A. Golf Course Fairways, Sod Farms, Lawns (around institutional, commercial and industrial buildings), and Ornamental Turfgrass:

NOTE: Do not use on home lawns and turf sites associated with apartment buildings, daycare centers, playgrounds, playfields, recreational park athletic fields, athletic fields located on or next to schools (ie., elementary, middle and high schools), campgrounds, churches, and theme parks.

NOTE: Sod farm turf treated with chlorothalonil prior to harvest must be mechanically cut, rolled and harvested. Do not use for sodfarms at application rates greater than 13 pounds of active ingredient, per acre, per year.

Do not apply more than 34.7 pints/acre (12.7 fl. ozs./1000 sq. ft.) of TERRANIL 6L per growing season (26 lbs. a.i./acre/growing season). For sodfarms, do not apply more than 17.4 pints/acre (6.4 fl. ozs./1000 sq. ft.) of TERRANIL 6L per growing season (13 lbs. ai./acre/growing season). The minimum growing retreatment interval for single application rates **up to** 9.75 pints/acre (3.6 fl. ozs./1000 sq. ft.) of TERRANIL 6L (7.3 lbs. a.i./acre) is 7 days. Do not apply more than one application of a rate greater than 9.75 pints/acre (3.6 fl. oz./1000 sq. ft.) of TERRANIL 6L (7.3 lb. a.i./acre) per growing season. The maximum single application rate is 15.1 pints/acre (5.5 fl. ozs./1000 sq. ft.) of TERRANIL 6L (11.3 lbs. a.i./acre).

Apply TERRANIL 6L in 30 to 40 gallons of water per acre. Begin applications when conditions favor disease development and repeat applications as long as these conditions persist. Under severe disease conditions use the highest rate and shortest interval corresponding with the application schedule selected from the table below.

DO NOT mow or water after treatment until spray deposited on turfgrass is thoroughly dry; TERRANIL 6L should always be used in conjunction with good turf management practices.

Group B. Golf Course Tees and Greens.

Golf Course Tees: Do not apply more than 69.3 pints/acre (25.4 fl. ozs./1000 sq. ft.) of TERRANIL 6L (52 lbs. a.i./acre) per growing season. The minimum retreatment interval for single application rates **up to** 9.75 pints/acre (3.6 fl. ozs./1000 sq. ft.) of TERRANIL 6L (7.3 lbs. a.i./acre) is 7 days. The minimum retreatment interval after an application of a rate **greater than** 9.75 pints/acre (3.6 fl. ozs./1000 sq. ft.) of TERRANIL 6L (7.3 lbs. a.i./acre) is 14 days. Do not apply more than two applications of a rate greater than 9.75 pints/acre (3.6 fl. ozs./1000 sq. ft.) of TERRANIL 6L (7.3 lbs. a.i./acre) per growing season. The maximum single application rate is 15.1 pints/acre (5.5 fl. ozs./1000 sq. ft.) of TERRANIL 6L (11.3 lbs. a.i./acre).

Golf Course Greens: Do not apply more than 97.3 pints/acre (35.7 fl. ozs./1000 sq. ft.) of TERRANIL 6L (73 lbs. a.i./acre) per growing season. The minimum retreatment interval for single application rates **up to** 9.75 pints/acre (3.6 fl. ozs./1000 sq. ft.) of TERRANIL 6L (7.3 lbs. a.i./acre) is 7 days and the minimum retreatment interval after an application of a rate **greater than** 9.75 pints/acre (3.6 fl. ozs./1000 sq. ft.) of TERRANIL 6L (7.3 lbs. a.i./acre) is 14 days. Do not apply more than two applications of a rate greater than 9.75 pints/acre (3.6 fl. ozs./1000 sq. ft.) of TERRANIL 6L (7.3 lbs. a.i./acre) per growing season. The maximum single application rate is 15.1 pints/acre (5.5 fl. ozs./1000 sq. ft.) of TERRANIL 6L (11.3 lbs. a.i./acre).

Apply TERRANIL 6L in an adequate amount of water to provide complete coverage. This amount may vary from 90 to 450 gallons per acre. See table below for suggested rates and timing. Under severe disease conditions use the highest rate and shortest interval corresponding with the application schedule selected from the table below.

DO NOT mow or water after treatment until spray deposited on turfgrass is thoroughly dry: TERRANIL 6L should always be used in conjunction with good turf management practices.

Diseases Controlled*	Application Interval (days)	Pre-Disease Rates ^a			Post-Disease Rates ^a		
		fl. oz. product/1000 sq. ft.	pints product/acre	lbs. a.i./acre	fl. oz. product/1000 sq. ft.	pints product/acre	lbs. a.i./acre
Dollar Spot	7 to 10	1.0 ^b to 2.0	2.8 ^b to 5.0	2.1 ^b to 4.1	-	-	-
	7 to 21	2.0 to 3.6	5.5 to 9.75	4.1 to 7.3	-	-	-
	14	-	-	-	4.0 to 5.5	11 to 15.1	8.25 to 11.3
Leafspot Melting-out Brown Blight	7 to 10	2.0	5.5	4.1	-	-	-
	7 to 21	2.0 to 3.6	5.5 to 9.75	4.1 to 7.3	-	-	-
	14	-	-	-	4.0 to 5.5	11 to 15.1	8.25 to 11.3
Brown patch	7 to 14	2.0 to 3.6	5.5 to 9.75	4.1 to 7.3	-	-	-
	14	-	-	-	4.0 to 5.5	11 to 15.1	8.25 to 11.3
Gray Leafspot	7 to 10	2.0 to 3.6	5.5 to 9.75	4.1 to 7.3	-	-	-
	14	-	-	-	4.0 to 5.5	11 to 15.1	8.25 to 11.3
Red Thread	7 to 10	2.0 to 3.6	5.5 to 9.75	4.1 to 7.3	-	-	-
	14	3.6 to 5.5	9.9 to 15.1	7.4 to 11.3	5.5	15.1	11.3
Anthracnose	7 to 14	3.0 to 3.6	8.3 to 9.75	6.2 to 7.3	-	-	-
	14	3.6 to 5.5	9.9 to 15.1	7.4 to 11.3	-	-	-
Copper Spot	14	4.0 to 5.5	11 to 15.1	8.25 to 11.3	5.5	15.1	11.3
Stem Rust (Bluegrass)	14	4.0 to 5.5	11 to 15.1	8.25 to 11.3	5.5	15.1	11.3
DICHONDRA: Leafspot (CA Only)	14	4.0 to 5.5	11 to 15.1	8.25 to 11.3	5.5	15.1	11.3
Gray Snow Mold ^c	30	5.5	15.1	11.3	-	-	-
Fusarium (Gerlachia) Patch ^c	21 to 28	5.5	15.1	11.3	-	-	-
Algae ^c	7 to 14	2.0 to 3.6	5.5 to 9.75	4.1 to 7.3	2.0 to 3.6	5.5 to 9.75	4.1 to 7.3
	14	-	-	-	4.0 to 5.5	11 to 15.1	8.25 to 11.3

^a **Group A Turf:** Limit of one application per season at rates greater than 7.3 lbs. a.i./acre (9.75 pints/acre or 3.6 fl. oz./1000 sq. ft. of TERRANIL 6L).

Group B Turf: Limit of two applications per season at rates greater than 7.3 lbs. a.i./acre (9.75 pints/acre or 3.6 fl. oz./1000 sq. ft. of TERRANIL 6L).

^b Low rate is not effective on intensively mowed turfgrasses such as golf course tees and greens.

^c See specific use directions below.

* Diseases listed are caused by fungi, some of which are named as follows:

- Dollar spot: *Sclerotinia homeocarpa*; *Lanzia* or *Moellerodiscus* spp.
- Leafspots, Melting-out, Brown blight: *Drechslera* spp. (including *D. poae*, *D. siccans*), *Bipolaris sorokiniana*, *Curvularia* spp.
- Brown patch: *Rhizoctonia solani*, *R. zaeae*, *R. cerealis*
- Gray leafspot: *Pyricularia grisea*, *P. oryzae*
- Red Thread: *Laetisaria fuciformis*
- Anthracnose: *Colletotrichum graminicola*
- Copper spot: *Gloeocercospora sorghi*
- Stem rust: *Puccinia graminis*
- Dichondra leaf spot: *Alternaria* spp.
- Gray Snow Mold: *Typhula* spp.
- Fusarium (Gerlachia) Patch
- Algae

Gray snow mold caused by *Typhula* spp. – Group A and B – Turf: Apply in sufficient water to obtain adequate coverage (2 to 10 gallons per 1000 sq. ft.). Apply one application 15.1 pints/acre (5.5 fl. ozs./1000 sq. ft.) of TERRANIL 6L (11.3 lbs. a.i./acre). Application must be made before snow cover in autumn. Group B Turf: If snow cover is intermittent or lacking during the winter, a second application of TERRANIL 6L at 15.1 pints/acre (5.5 fl. ozs./1000 sq. ft.) may be applied one month after the first application.

Fusarium (Gerlachia) Patch: Group A and B Turf: In areas where pink snow mold (Gerlachia or Fusarium patch) is likely to occur, apply TERRANIL 6L at 15.1 pints/acre (5.5 fl. ozs./1000 sq. ft.) (11.3 lbs. a.i./acre) in combination with products containing iprodione at 88 ozs. a.i./acre (2 ozs. a.i./1000 sq. ft.) of turf area. Read and observe all label directions for products containing these active ingredients. For control of Fusarium patch only in areas where snow cover is intermittent or lacking during the winter, apply 15.1 pints/acre of TERRANIL 6L (5.5 fl. ozs./1000 sq.

ft.)(11.3 lbs. a.i./acre). Make application in late autumn. Group B Turf: Apply a second application of 15.1 pints/acre (5.5 fl. ozs./1000 sq. ft.) of TERRANIL 6L 21 to 28 days after the first application unless conditions favorable for Fusarium patch no longer prevail.

Algae: Group A and B Turf: For prevention of algae on turfgrasses, apply TERRANIL 6L at the rate of 5.5 to 9.75 pints/acre (2.0 to 3.6 fl. ozs./1000 sq. ft.) (4.1 to 7.3 lbs. a.i./acre) on a 7 to 14 day schedule. Under severe algae conditions use the 9.75 pints/acre (3.6 fl. oz./1000 sq. ft.) rate and apply on a 7 day schedule.

When algae is well established, every attempt should be made to dry out the afflicted area. Once dry, spiking or verticutting should be done to enhance turfgrass recovery in conjunction with a TERRANIL 6L application at the rate of 11 to 15.1 pints/acre (4.0 to 5.5 fl. ozs./1000 sq. ft.). Group B Turf: A second application at the 15.1 pints/acre (5.5 fl. ozs./1000 sq. ft.) rate may be made 14 days after the first application.

Group A and B Turf: Following applications of the 15.1 pints/acre (5.5 fl. ozs./1000 sq. ft.) rate, several applications of TERRANIL 6L at a rate of 5.5 to 9.75 pints/acre (2.0 to 3.6 fl. ozs./1000 sq. ft.)(4.1 to 7.3 lbs. a.i./acre) on a 7 to 14 day interval may be necessary for turfgrass recovery. Only a preventative spray program with TERRANIL 6L will prevent a recurrence of the algae when environmental conditions are favorable.

Ornamental Plants

Apply TERRANIL 6L at a rate of 1 $\frac{3}{8}$ pints (1.0 lb. a.i.) per 100 gallons of water unless other directions are given in the tables below. DO NOT apply more than 48.5 pints of TERRANIL 6L (36.4 lbs. a.i./acre) per growing season to field grown ornamentals. Apply in a spray to run-off, when conditions are favorable for disease development. Repeat applications at 7 to 14 day intervals until conditions are no longer favorable. During periods when conditions favor severe disease incidence, generally cloudy or wet weather, apply TERRANIL 6L at 7 day intervals. The minimum retreatment interval is 7 days. TERRANIL 6L should be applied to plants when both foliage and flowers are dry, or nearly dry.

DO NOT combine TERRANIL 6L in the spray tank with pesticides, surfactants or fertilizers, unless your prior use has shown the combination to be physically compatible, effective and noninjurious under your conditions of use.

TERRANIL 6L may be used in greenhouses for ornamental plants. DO NOT use mistblowers or high pressure spray equipment when making applications of TERRANIL 6L in greenhouses.

Use of TERRANIL 6L is recommended for control of fungal diseases referred to by numbers in parentheses following each ornamental. Ornamentals listed on this label have been tested and found to tolerate applications of TERRANIL 6L at the recommended rates. The user should test for possible phytotoxic response, using recommended rates on ornamental plants on a small area prior to commercial use. Applications made during bloom may damage flowers and/or fruits.

Fruits and other structures which may be borne on treated plants MUST NOT BE EATEN.

Ornamentals recommended for treatment with TERRANIL 6L

Broadleaf Shrubs and Trees:

Andromeda (Pieris) (4)	Holly (1)
Ash (Fraxinus) (1)	Lilac (5)
Aspen (1)	Magnolia (1)
Azalea (1,2,4)	Maple (1)
Buckeye, Horsechestnut (1)	Mountain Laurel (1)
Cherry-Laurel (1)	Oak (red group only) (1,7)
Crabapple (1,6,8)	Oregon-Grape (Mahonia) (6)
Dogwood (1)	Photinia (1)
Eucalyptus (3)	Poplar (1)
Euonymus (1)	Privet (Ligustrum) (1)
Firethorn (Pyracantha) (1)	Rhododendron (1,2,4)
Flowering Almond (1,2)	Sand Cherry (1,2)
Flowering Cherry (1,2)	Sequoia (1)
Flowering Peach (1,2)	Spiraea (1)
Flowering Plum (1,2)	Sycamore, Planetree (1)
Flowering Quince (1,2)	Viburnum (5)
Hawthorn (1,6)	Walnut (Juglans) (1)

Flowering Plants^a and Bulbs

Arabian Violet (2)
 Begonia (1)
 Camellia (2)
 Carnation (1,2)
 Chrysanthemum (1,2)
 Crocus (1)
 Daffodil (1)
 Daisy (1)
 Geranium (1,6)
 Gladiolus (1,2)
 Hollyhock (6)
 Hydrangea (foliage only) (1,6)
 Iris (1,2)

Iris, bulbous (1)
 Lily (1)
 Lily, Asiatic (1)
 Marigold (1)
 Narcissus (1)
 Pansy (1)
 Petunia (1,4)
 Phlox (1)
 Poinsettia^b (1)
 Rose^c (1)
 Statice (1)
 Tulip (1)
 Zinnia (1,5)

^a Avoid applications during bloom period on plants where flower injury is unacceptable.

^b Discontinue applications prior to bract formation; phytotoxicity is possible on the bracts.

^c Use 1 pint TERRANIL 6L (.75 lbs. a.i.) per 100 gallons of water.

Foliage Plants

Aglaonema (1)
 Areca palm (1)
 Artemisia (1)
 Boston Fern (*Nephrolepis*) (1)
 Dumbcane (*Diffenbachia*) (1)
 Dracaena (1)
 Fatsia (*Aralia*) (1)
 Ficus (1)
 Florida ruffle fern (1)
 Leatherleaf fern (1)
 Lipstick plant (1)

Ming aralia (1)
 Oyster plant (*Rhoeo*) (1)
 Pachysandra^d (1)
 Parlor Palm (*Chamaedorea*) (1)
 Peperomia (1)
 Philodendron (1,4)
 Prayer plant (*Maranta*) (1)
 Syngonium (1)
 Zebra plant (*Aphelandra*) (1)

^d Use 2¾ pints TERRANIL 6L (2.1 lbs. a.i.) per 100 gallons of water.

Diseases controlled with TERRANIL 6L:**1. Leafspots/Foliar Blights:**

Achtinopelte leafspot
 Alternaria leafspot/leaf blight
 Anthracnose leaf blotch, spot
 Anthracnose (*Discula*) blight
 Ascochyta blight
 Bipolaris (*Helminthosporium*) leafspot
 Black spot on roses
 Botrytis leafspot, leaf blight
 Cephalosporium leafspot
 Cercospora leafspot
 Cercosporidium leafspot
 Corynespora leafspot
 Coryneum blight (shothole)
 Curvularia leafspot
 Cylindrosporium leafspot
 Dactylaria leafspot
 Didymellina leafspot
 Drechslera leafspot

Fabraea (*Entomosporium*) leafspot
 Fusarium leafspot
 Gloeosporium black leafspot
 Ink spot (*Dreschlera*)
 Marssonina leafspot
 Monilinia blossom blight, twig blight
 Mycosphaerella ray blight
 Myrothecium leafspot, brown rot
 Nematostoma leaf blight
 Phyllosticta leafspot
 Ramularia leafspot
 Rhizoctonia web blight
 Septoria leafspot
 Sphaeropsis leafspot
 Stagonospora leaf scorch
 Tan leaf spot (*Curvularia*)
 Volutella leaf blight

2. Flower spots/blights:

Botrytis flower spot, flower blight
 Curvularia flower spot
 Monilinia blossom blight
 Ovulinia flower blight
 Rhizopus blossom blight
 Sclerotinia flower blight

3. Cylindrocladium stem canker**4. Phytophthora leaf blight, dieback****5. Powdery mildews***Erysiphe cichoracearum**Microsphaera* spp.**6. Rusts***Gymnosporangium* spp.*Pucciniastrum hydrangeae**Puccinia* spp.**7. Taphrina blister****8. Scab** (*Venturia inaequalis*)

The following ornamental plant species which have been tested with TERRANIL 6L at recommended rates did not exhibit phytotoxicity:

Botanical Name	Common Name
<i>Aechmea fasciata</i>	Aechmea
<i>Araucaria heterophylla</i>	Norfolk Island Pine
<i>Bougainvillea</i> spp.	Bougainvillea
<i>Caladium</i> spp.	Caladium
<i>Calathea makoyana</i>	Peacock Plant
<i>Calistephus chinensis</i>	Aster
<i>Carissa grandiflora</i>	Natal Plum
<i>Clerodendron thomsonae</i>	Bleeding Heart
<i>Codiaeum</i> spp.	Croton
<i>Cordylina terminalis</i>	Ti Plant
<i>Crassula argentea</i>	Jade Plant
<i>Dionaea muscipula</i>	Venus Fly Trap
<i>Dizygotheca elegantissima</i>	False Aralia
<i>Epipremnum aureum</i>	Golden Pothos, Scindapsus
<i>Episcia cupreata</i>	Flame Violet
<i>Fittonia</i> spp.	Silver-nerve plant
<i>Gerbera jamesonii</i>	Gerbera Daisy
<i>Gynura sarmentosa</i>	Purple Passion Vine
<i>Gypsophilia paniculata</i>	Baby's Breath
<i>Hoya</i> spp.	Wax Plant
<i>Ilex cornuta</i>	Chinese Holly
<i>Ilex crenata</i>	Japanese Holly
<i>Impatiens</i> spp.	Impatiens
<i>Pilea cadierei</i>	Aluminum Plant
<i>Sansevieria trifasciata</i> "Hahnii"	Birdsnest Sansevieria
<i>Tolmeia menziesii</i>	Piggy-back plant
<i>Yucca elephantipes</i>	Spineless Yucca
<i>Zygocactus truncatus</i>	Christmas Cactus

NOTE: DO NOT apply TERRANIL 6L to either green or variegated Pittosporum or to Schefflera, as multiple applications have been demonstrated to cause phytotoxic responses.

TERRANIL 6L® is a registered trademark of Winfield Solutions, LLC.

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE

REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THE NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.