

9779 - 335

7/3/2012

1/15



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON D C 20460

JUL 03 2012

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

Alice Walker Ph D
Registration Specialist
Winfield Solutions LLC
3094 Country Club Rd
Senatobia, MS 38668

Subject ASCEND
EPA Registration No 9779 335
Label Amendment to revise the application options for corn (sweet pop and field) and adjust the application timing for soybeans
Decision # 462533
Application Dated March 07 2012

Dear Dr Walker

The amendment referred to above submitted in connection with registration under FIFRA section 3(c)(5) is acceptable provided that you

- 1 Submit and/or cite all data required for registration/reregistration of your product under FIFRA section 3(c)(5) when the Agency requires all registrants of similar products to submit such data
- 2 Submit three (3) copies of your final printed labeling before you release the product for shipment Final printed labeling means the label or labeling of the product when distributed or sold Clearly legible reproductions or photo reductions will be accepted for unusual labels such as those silk screened directly onto glass or metal containers or large bags or drum labels

If these conditions are not complied with the registration will be subject to cancellation in accordance with FIFRA section 6(b) Your release for shipment of the product bearing the amended labeling constitutes acceptance of these conditions

Should you have any questions you may contact Ms Menyon Adams directly at 703 347 8496 or via email at adams.menyon@epa.gov

Sincerely

Lisa A. Smith
Chief
Biochemical Pesticides Branch
Biopesticides and Pollution
Prevention Division (7511P)

MAIL ROOM	7511P			
US MAIL	Adams			
DATE	07/02/12			

EPA Form 1328-1A (1/99)

2/15



ASCEND®

PLANT GROWTH REGULATOR

*Hormone compounds to stimulate plant growth
Concentrations based on biological activity*

ACTIVE INGREDIENTS

Cytokinin as Kinetin	0.090%
*Gibberellic Acid	0.030%
*Indole Butyric Acid	0.045%

OTHER INGREDIENTS

	<u>99.835%</u>
TOTAL	100.000%

- *Contains 0.03 oz cytokinins/qt
- *Contains 0.015 oz indole butyric acid/qt
- *Contains 0.01 oz gibberellic acid/qt

KEEP OUT OF REACH OF CHILDREN

CAUTION

FIRST AID	
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes • Remove contact lenses if present after the first 5 minutes then continue rinsing eye • Call a poison control center or doctor for treatment advice
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing • Rinse skin immediately with plenty of water for 15-20 minutes • Call a poison control center or doctor for treatment advice
Have the product container or label with you when calling a poison control center or doctor or going for treatment. HOTLINE NUMBER In case of medical emergency call 1-877-424-7452	

Read additional precautionary statements found inside booklet

SHAKE WELL BEFORE USING

EPA Reg No 9779-335

EPA Est No 63603 KS 1
NET CONTENTS 1 Gal

Distributed by
WINFIELD SOLUTIONS LLC
P O Box 64589 St Paul MN 55164

ACCEPTED

JUL 03 2012

Under the Federal Insecticide Fungicide
and Rodenticide Act as amended for
the pesticide registered under
EPA Reg No

9779-335

1/0307/2

Lot No _____

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Causes eye irritation Harmful if absorbed through skin Avoid contact with skin eyes or clothing
Wash thoroughly with soap and water after handling Remove and wash contaminated clothing
before reuse Wear the appropriate Personal Protective Equipment (PPE)

Personal Protective Equipment

Mixers loaders applicators and other handlers must wear

- long sleeved shirt and long pants
- shoes plus socks and
- chemical resistant gloves

Follow manufacturer's instructions for cleaning and maintaining PPE If no instructions for
washables use detergent and hot water Keep and wash PPE separately from other laundry

USER SAFETY RECOMMENDATIONS

Users should

- Wash hands before eating drinking chewing gum using tobacco or using the toilet
- Remove clothing immediately if pesticide gets inside Then wash thoroughly and put on clean clothing
- Remove PPE immediately after handling this product Wash the outside of gloves before removing As soon as possible wash thoroughly and change into clean clothing

ENVIRONMENTAL HAZARDS

For terrestrial uses Do not apply directly to water or to areas where surface water is present or to
intertidal areas below the mean high water mark Do not contaminate water by cleaning of
equipment or disposal of equipment washwater or rinsate

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling Do not
apply this product in a way that will contact workers or other persons either directly or through drift
Only protected handlers may be in the area during application

For any requirements specific to your State or Tribe consult the State or Tribal agency responsible
for pesticide regulation

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard 40
CFR part 170 This standard contains requirements for the protection of workers on farms forests
nurseries and greenhouses and handlers of agricultural pesticides It contains requirements for
training decontamination notification and emergency assistance It also contains specific
instructions and exceptions pertaining to the statements on this label about personal protective
equipment and restricted entry intervals The requirements in this box only apply to uses of this
product that are covered by the Worker Protection Standard

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4
hours unless wearing appropriate PPE

EXCEPTION If the product is soil incorporated the Worker Protection Standard under certain
circumstances allows workers to enter the treated area if there will be no contact with anything that
has been treated

4/15

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants soil or water is

- coveralls over long sleeved shirt and long pants
- chemical resistant gloves and
- shoes plus socks

GENERAL CHEMIGATION INSTRUCTIONS

Apply this product only through sprinkler including center pivot lateral move side (wheel) roll traveler big gun solid set hand move or furrow irrigation systems Do not apply this product through any other type of irrigation system

Crop injury lack of effectiveness or illegal pesticide residues in the crop can result from non uniform distribution of treated water

If you have questions about calibration you should contact State Extension Service specialists equipment manufacturers or other experts

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label prescribed safety devices for public water systems are in place

A person knowledgeable of the chemigation system and responsible for its operation or under supervision of the responsible person shall shut the system down and make necessary adjustments should the need arise

Maintain agitation in the supply tank while adding the required amount of ASCEND and throughout the application ASCEND should be added to the supply tank at the end of water application (prior to last complete cycle in moving systems)

The correct amount of ASCEND to add is calculated as the rate in fluid oz per acre x the number of acres covered by the contents of the supply tank

(For example if the supply tank covers ten acres and the rate on the label for that crop is 2 fluid ounce per acre add $10 \times 2 = 20$ fluid ounces to the supply tank at the beginning of the last full cycle)

CHEMIGATION SYSTEMS CONNECTED TO PUBLIC WATER SYSTEMS

Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year

Chemigation systems connected to public water systems must contain a functional reduced pressure zone backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction As an option to the RPZ the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe

The pesticide injection pipeline must contain a functional automatic quick closing check valve to prevent the flow of fluid back toward the injection pump

The pesticide injection pipeline must contain a functional normally closed solenoid operated valve located on the intake side of the injection pump and connected to the system interlock to prevent

fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the pump motor stops or in cases where there is no water pump when the water pressure decreases to the point where pesticide distribution is adversely affected

Systems must use a metering pump such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with the pesticides and capable of being fitted with a system interlock

Do not apply when wind speed favors drift beyond the area intended for treatment

The pesticide supply tank should be agitated throughout the application of ASCEND ASCEND should be applied at the end of the water application

ASCEND should be applied at the end of the irrigation period in a sufficient amount of water to allow proper coverage of plant or crop but not to exceed 18 fluid ounces of ASCEND per acre per application

IN FURROW CHEMIGATION

1 Systems using a gravity flow pesticide dispensing system must meter the pesticide into the water at the head of the field and downstream of a hydraulic discontinuity such as a drop structure or weir box to decrease potential for water source contamination from backflow if water flow stops

2 Systems utilizing a pressurized water and pesticide injection system must meet the following requirements

a The system must contain a functional check valve vacuum relief valve and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow

b The pesticide injection pipeline must contain a functional automatic quick closing check valve to prevent the flow of fluid back toward the injection pump

c The pesticide injection pipeline must also contain a functional normally closed solenoid operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down

d The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops

e The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected

f Systems must use a metering pump such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock

Maintain agitation in the supply tank while adding the required amount of ASCEND and throughout the application ASCEND should be added to the supply tank at the end of water application (prior to last complete cycle in moving systems)

6/15

The correct amount of ASCEND to add is calculated as the rate in fluid oz per acre x the number of acres covered by the contents of the supply tank

(For example if the supply tank covers ten acres and the rate on the label for that crop is 2 fluid ounce per acre add $10 \times 2 = 20$ fluid ounces to the supply tank at the beginning of the last full cycle)

SPRINKLER CHEMIGATION

The system must contain a functional check valve vacuum relief valve and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow

The pesticide injection pipeline must contain a functional automatic quick closing check valve to prevent the flow of fluid back toward the injection pump

The pesticide injection pipeline must also contain a functional normally closed solenoid operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to a point where pesticide distribution is adversely affected

Systems must use a metering pump such as a positive displacement injection pump (e.g diaphragm pump) effectively designed and constructed of materials that are compatible with a system interlock

Do not apply when wind speed favors drift beyond the area intended for treatment

Maintain agitation in the supply tank while adding the required amount of ASCEND and throughout the application ASCEND should be added to the supply tank at the end of water application (prior to last complete cycle in moving systems)

The correct amount of ASCEND to add is calculated as the rate in fluid oz per acre x the number of acres covered by the contents of the supply tank

(For example if the supply tank covers ten acres and the rate on the label for that crop is 2 fluid ounce per acre add $10 \times 2 = 20$ fluid ounces to the supply tank at the beginning of the last full cycle)

ASCEND should be applied at the end of the irrigation period in a sufficient amount of water to allow proper coverage of plant or crop but not to exceed 18 fluid ounces of ASCEND per acre per application

IMPORTANT Read the entire Directions for Use and the Notice before using this product. If terms are not acceptable return the unopened product container at once

Apply ASCEND by ground or air. If applied by air use 3 to 5 gallons of water per acre. If applied by ground use 5 to 25 gallons of water per acre. For turfgrass apply ASCEND by ground using 0.2 to 0.5 gallon on water per 1 000 square feet

Test results have shown that this product can stimulate higher yields through a larger root mass earlier fruiting and increased fruit retention. ASCEND is a tool to increase plant efficiency

SPRAY PROGRAM FOR VEGETABLE CROPS

BEANS AND PEAS

- 1st Application Apply 3.2 fluid ounces per acre when the first trifoliate is unfolded
- 2nd Application Apply 3.2 fluid ounces per acre 2 weeks after the first application
- 3rd Application Apply 3.2 fluid ounces per acre at first bloom

ASPARAGUS BROCCOLI CABBAGE CELERY LETTUCE MINT AND SPINACH

- 1st Application Apply 3.2 fluid ounces per acre when the fifth leaf begins to unfold
- 2nd Application Apply 3.2 fluid ounces per acre 2 weeks after the first application
- 3rd Application Apply 3.2 fluid ounces per acre 2 weeks after the second application

For maximum benefit apply continuous applications of 0.8 - 1.2 fluid ounces per acre at 7-10 day intervals after the first application throughout the production season

CANTALOUPE CUCUMBERS MUSKMELON WATERMELON HONEYDEW OKRA AND SQUASH

- 1st Application Apply 3.2 fluid ounces per acre when the third leaf begins to unfold
- 2nd Application Apply 3.2 fluid ounces per acre 2 weeks after the first application
- 3rd Application Apply 3.2 fluid ounces per acre 2 weeks after the second application

For maximum yields make continuous applications of 2 fluid ounces per acre at 7-10 day intervals after the first application throughout the growing season

EGGPLANT PEPPER AND TOMATO

- 1st Application Apply 3.2 fluid ounces per acre when the plants have 3 true leaves
- 2nd Application Apply 3.2 fluid ounces per acre 2 weeks after the first application
- 3rd Application Apply 3.2 fluid ounces per acre 2 weeks after the second application

For maximum yields and quality make continuous applications of 0.8 fluid ounces per acre after the first application at 7-10 day intervals throughout the growing season

SWEET CORN AND POPCORN Apply one, two, or all of the following applications

Apply 4.5 to 6 fluid ounces per acre in furrow or alternatively 2 inches beside and 2 inches below seed or alternatively 3 inches below the seed with a strip till machine at planting

AND/OR

Apply 3.2 fluid ounces per acre when the plants are in the 4-6 leaf stage

AND/OR

Apply 3.2 fluid ounces per acre at the 8-10 leaf stage

NOTE If seed has been treated with ASCEND do not apply ASCEND as an in furrow, band, side dress or mark out application

WHITE OR RED POTATOES

Apply 4.5 to 6 fluid ounces per acre in a band, mark out, side dress or in furrow application before or after planting

NOTE If seed has been treated with ASCEND do not apply ASCEND as an in furrow, band, side dress or mark out application

For foliar applications apply according to *one* of the following schedules

To increase tuber size, number and promote better rooting

- 1st Application Apply 3.2 fluid ounces per acre at tuber initiation
 - 2nd Application Apply 3.2 fluid ounces per acre 2-3 weeks after the first application. The last application should be during tuber bulking
- OR

To enhance tuber size and uniformity

- 1st Application Apply 3.2 fluid ounces per acre at tuber initiation
- 2nd Application Apply 3.2 fluid ounces per acre at the onset of tuber bulking

CARROTS PARSLEY RADISHES AND TURNIPS

- 1st Application Apply 3.2 fluid ounces per acre when the plants have 3 true leaves
- 2nd Application Apply 3.2 fluid ounces per acre 2 weeks after the first application
- 3rd Application Apply 3.2 fluid ounces per acre 2 weeks after the second application

SWEET POTATOES AND YAMS

- 1st Application Apply 0.2 to 0.4 fluid ounces per acre on a band just wide enough to cover all the plants seven to fourteen days after transplanting
- 2nd Application Apply 0.5 fluid ounces per acre in a band as above at twenty eight days after transplanting
- 3rd Application Apply 0.1 fluid ounces per week along with a foliar fertilizer such as 15-5-5 at the rate of 32 fluid ounces or 1 quart per acre Continue this program on a weekly basis until the potatoes have desirable harvest size

NOTE If seed has been treated with ASCEND do not apply ASCEND as an in furrow band side dress or mark out application

FOLIAR SPRAY PROGRAM FOR FRUIT CROPS

BANANAS

- 1st Application Apply 3 to 6 fluid ounces per acre shortly prior to or at first bloom
- 2nd Application Apply 3 to 6 fluid ounces per acre two to three weeks after the first application

CITRUS (GRAPEFRUIT LEMON LIME AND ORANGES)

- 1st Application Apply 3 to 6 fluid ounces per acre at first bloom
 - 2nd Application Apply 3 to 6 fluid ounces per acre two to three weeks later
- If there is an extended bloom period make additional applications at 3 to 6 fluid ounces per acre

GRAPES

- 1st Application Apply 3.2 fluid ounces per acre shortly prior to or at bloom stage
- 2nd Application Apply 3.2 fluid ounces per acre 2 weeks after the first application

GUAVA AND PAPAYA

- 1st Application Apply 3.2 fluid ounces per acre shortly prior to or at first bloom stage
- 2nd Application Apply 3 to 6 fluid ounces per acre 2 to 3 weeks after the first application

POME (APPLE MAYHAW)

- 1st Application Apply 3 to 6 fluid ounces per acre shortly prior to or at first bloom
- 2nd Application Apply 3 to 6 fluid ounces per acre two to three weeks after the first application

STONE (PEACH, CHERRY APRICOT, NECTARINE)

- 1st Application Apply 3 to 6 fluid ounces per acre shortly prior to or at first bloom
- 2nd Application Apply 3 to 6 fluid ounces per acre two to three weeks after the first application

STRAWBERRIES

- 1st Application Apply 3.2 fluid ounces per acre shortly prior to or at first bloom stage
- 2nd Application Apply 3.2 fluid ounces per acre 2 weeks after the first application

SPRAY PROGRAM FOR FIELD CROPS

ALFALFA Established

- 1st Application Apply 3.2 fluid ounces per acre upon dormancy break Apply when sufficient re growth is present
- 2nd Application Apply 3.2 fluid ounces per acre after each cutting Apply when sufficient re growth is present

ALFALFA Newly Seeded

- 1st Application Apply 3.2 fluid ounces per acre when seedling alfalfa is in the 3rd to 4th trifoliate stage

CANOLA

- 1st Application Apply 3.2 oz per acre between the rosette stage and bolting
- 2nd Application Apply 3.2 oz per acre at 20 percent bloom
- 3rd Application Apply 3.2 oz per acre at early pod fill

COTTON Non Transgenic Varieties Apply ASCEND according to *one* of the following schedules

NOTE If seed has been treated with ASCEND do not apply ASCEND as an in furrow band side dress or mark out application

Schedule A

- 1st Application Apply 3 fluid ounces per 50 lbs of seed in the hopper box
OR
Apply 3.2 fluid ounces per acre in furrow or alternatively 2 inches beside and 2 inches below seed or alternatively 3 inches below the seed with a strip till machine at planting
- 2nd Application Apply 3 fluid ounces per acre at pinhead square This can be applied in a tank mix that contains 4 fl oz per acre of Mepex® Plant Regulator brand of mepiquat chloride
- 3rd Application Apply 4 fluid ounces per acre at early bloom

Schedule B

- 1st Application Apply 2 fluid ounces per acre on a band at the 3.7 leaf stage
- 2nd Application Apply 3 fluid ounces per acre at the pinhead square stage This can be applied in a tank mix that contains 4 fl oz per acre of Mepex® Plant Regulator brand of mepiquat chloride
- 3rd Application Apply 3 fluid ounces per acre at early bloom

COTTON Transgenic Varieties

(Cotton varieties that have been genetically manipulated to have insect resistance and/or herbicide resistance built in)

NOTE If seed has been treated with ASCEND do not apply ASCEND as an in furrow band side dress or mark out application

- 1st Application Use according to *one* of the following methods
Apply 3 fluid ounces per 50 lbs of seed in the hopper box
OR

10/15

Apply 3.2 fluid ounces per acre in furrow or alternatively 2 inches beside and 2 inches below seed or alternatively 3 inches below the seed with a strip till machine at planting

2nd Application Apply 4 fluid ounces per acre at pinhead square This can be applied in a tank mix that contains 4 fl oz per acre of Mepex® Plant Regulator brand of mepiquat chloride

3rd Application – Repeat the above application at first bloom If needed for vegetative growth control repeat the above application at mid bloom

Higher rates and/or late season applications may be warranted under high stress conditions where square and/or boll retention is needed During the bloom and post bloom period additional applications or higher rates can be applied but do not exceed a total of 24 fluid ounces per acre per season

FIELD CORN Apply one two or all of the following applications

Apply 4.5 to 6 fluid ounces per acre in furrow or alternatively 2 inches beside and 2 inches below seed or alternatively 3 inches below the seed with a strip till machine at planting

AND/OR

Apply 3.2 fluid ounces per acre at the 3.4 leaf stage

AND/OR

Apply 3.2 to 4 fluid ounces per acre between initial silking and brown silk

NOTE If seed has been treated with ASCEND do not apply ASCEND as an in furrow band side dress or mark out application

FLAX

1st Application Apply 3.2 fluid ounces per acre when the plant is 2.4 inches tall

2nd Application Apply an additional 3.2 fluid ounces per acre two to three weeks later

GRAIN SORGHUM

1st Application Apply 4.5 to 6 fluid ounces per acre in furrow or alternatively 2 inches beside and 2 inches below seed or alternatively 3 inches below the seed with a strip till machine at planting

2nd Application Apply 3.2 fluid ounces per acre at the 3.5 leaf stage

3rd Application Apply 3.2 fluid ounces per acre after the 8th but before the 12th leaf stage

NOTE If seed has been treated with ASCEND do not apply ASCEND as an in furrow band side dress or mark out application

PEANUTS

1st Application Apply 3.2 fluid ounces per acre at the 3.5 leaflet stage

2nd Application Apply 3.2 fluid ounces per acre at early flowering

3rd Application Apply 3.2 fluid ounces per acre at initial pegging

4th Application Apply 4.8 fluid ounces per acre during early pod fill

SOYBEANS Apply according to one of the following schedules

1) Apply 3.2 fluid ounces per acre at the 3.5 trifoliolate leaf stage Apply a second application of 3.2 fluid ounces between R1 and R3

2) If the first application is missed apply 6.4 fluid ounces per acre between R1 and R3

SUGAR BEETS

In Furrow or Mark Out

Apply 4.5 to 6 fluid ounces per acre in furrow or mark out

Foliar program

- 1st Application Apply 3.2 fluid ounces per acre between the 2nd and 10th true leaf stage
 2nd Application Apply 3.2 fluid ounces per acre 2-3 weeks after the first application

NOTE If seed has been treated with ASCEND do not apply ASCEND as an in furrow band side dress or mark out application

SUGARCANE

- 1st Application – Use *one* of the following methods
 Apply 2 fluid ounces per acre in the furrow at planting
 OR
 Apply 3.2 fluid ounces per acre at the 2-3 leaf stage
 2nd Application Apply 3.2 fluid ounces per acre one month after emergence
 Additional Applications Apply 3.2 fluid ounces per acre on monthly intervals throughout the production season for maximum benefit

SUNFLOWERS

- 1st Application Apply 3.2 fluid ounces per acre at 4 true leaves
 2nd Application Apply an additional 3.2 fluid ounces per acre two to three weeks later

WHEAT BARLEY OATS AND RYE Apply according to *one* of the following schedules

- 1) Apply 4.5 to 6 fluid ounces per acre in furrow at planting
 OR
- 2) If no at planting application apply 3.2 fluid ounces per acre prior to jointing Apply 3.2 fluid ounces at the flag leaf stage

NOTE If seed has been treated with ASCEND do not apply ASCEND as an in furrow band side dress or mark out application

FOLIAR SPRAY PROGRAM FOR RICE

ASCEND should be applied at 3.2 fluid ounces per acre as a foliar spray to the plant during either one of the following stages of development

Primary Recommendations 3 to 7 Leaf Stage This application must be made after the rice seedling has 3 fully emerged leaves and the 4th leaf is beginning to emerge but before the seedling has completed development of 7 leaves or 3 tillers. This period for application generally begins about 3-6 weeks after seeding and ends 5-9 weeks after seeding. The duration of this period depends on the variety and the growing conditions. This application may be made in conjunction with corresponding herbicide applications.

Alternate Recommendation Two Millimeter (mm) Panicle Growth Stage If the primary application is missed ASCEND can be applied to stimulate cell differentiation in the developing panicle. This application must be made when no more than 10% of the main culms are at the 2 mm panicle growth stage. The 2 mm panicle growth stage occurs immediately after internode elongation or joint movement has begun. ASCEND must be applied as soon as internode elongation is detected so the 2 mm panicle growth stage is not missed. It is better to apply slightly early than to apply late. **IMPORTANT** Timing of the application at 2 mm growth stage is critical. Check the entire field for stage of plant development. Large fields may require split applications on upper and lower ends of the field to ensure proper timing throughout the field.

12/15

TURFGRASS

On all turfgrass regardless of use use no more than 0.3 fluid ounces per 1 000 square feet per month

WARM SEASON TURF (Bermuda Bermuda hybrids, Zoysia Centipede St Augustine etc)

For lower traffic areas and where ASCEND is used as a maintenance program begin applications early in the growing season Apply at the rate of 0.1 to 0.15 fluid ounces per 1 000 square ft Maintenance application should be made on a two to three week schedule throughout the growing season Applications can be made with foliarly applied urea for added benefits

COOL SEASON TURF (Tall Fescue Rye Bentgrass Bluegrass etc) Apply 0.15 fluid ounces per 1 000 square feet in fall or when stand is established Repeat application in late winter when grasses begin to grow actively

APPLICATION WITH FOLIARLY APPLIED UREA Maximum benefit and color can be achieved when ASCEND applications are made with foliarly applied urea solutions To prepare urea solution dissolve 46% urea into spray solution at the rate of 1.0 lb per 5 000 square feet to be sprayed and apply with specified rate of ASCEND

SPECIFIC RATES OF APPLICATION

TEES AND GREENS Apply 0.1 to 0.15 fluid ounces per 1 000 square feet on a 2 week schedule throughout the growing season Begin in early spring after grasses have begun to grow Sunbelt and transition zones should continue spray program throughout playing season Courses north of the transition zone should continue applications through September

FAIRWAYS Begin applications in early spring as soon as grasses have begun to actively grow Apply 0.1 fluid ounce per 1 000 square feet and repeat on a monthly schedule as long as grass is growing

PRE TOURNAMENT QUICK GREEN UP Apply at the rate of 0.1 to 0.15 fluid ounces per 1 000 square feet in conjunction with urea solution 4 to 5 days prior to playing time Make application with a minimum spray volume of 0.5 gallon of water per 1 000 square feet

SPRING DORMANCY BREAK Apply 0.1 fluid ounce per 1 000 square feet in spring as soon as new growth (opening) is visible Raking of thatch prior to making this application is most desirable Application at this time generates rapid growth and often reduces incidence of spring die back on certain species of grass

FALL APPLICATION FOR WINTER HARDINESS Make 2 applications 7-10 days apart in late summer or early fall just prior to the cessation of normal active growth Apply 0.1 to 0.15 fluid ounces per 1 000 square feet Make application with a spray volume of 0.5 gallon of water per 1 000 square feet Applications at this time will greatly increase root mass and depth of roots Winter kill problems are often greatly reduced

COMMERCIAL TURF CEMETERIES, ATHLETIC FIELDS, GOLF COURSES AND OTHER FINE TURF AREAS Applications of 0.1 to 0.15 fluid ounces per 1 000 square feet made at any point during the growing season will produce desirable results Make applications during the very early growth stages and continue on a regular monthly schedule throughout the growing season Healthier and more beautiful turf can be realized in high traffic areas such as golf greens and tees by making regular applications every two weeks

13/15

SOD FARMS

Apply 4 to 8 fluid ounces per acre on a monthly basis during the growing season Two weeks prior to cutting sod make an application of 4 to 8 fluid ounces per acre

SPECIFIC RATES OF APPLICATION

After sod is cut a reestablishment program is necessary Start this program as soon as there is any greening over 30% of the area Spray with 4 to 8 fluid ounces per acre of ASCEND Repeat in 2 weeks and thereafter once per month throughout the growing season Make a final application of 4 to 8 fluid ounces per acre 2 weeks before dormancy

Start the monthly program again as soon as some green up has started in the spring

When species started from seed have reached 1 inch in height the monthly treatment may be started and followed in the same way as non seeded varieties

SPECIAL NOTE FOR ALL DIRECT SEEDED GRASSES

Acting through its unique combination of plant growth regulators ASCEND is a ready to use seed dressing that aids in enhancing germination and early season root and top growth

ASCEND can be used at the rate of 2 to 4 fluid ounces per 100 pounds of seed Sufficient water needs to be added to insure uniform coverage Improper coverage will minimize product performance

SEED TREATMENT

SPECIAL NOTE FOR ALL DIRECT SEEDED CROPS

Acting through its unique combination of plant growth regulators ASCEND is a ready to use seed dressing that aids in enhancing germination and early season root and top growth

Use ASCEND at the rate of 1 to 4 fl oz per 100 lbs of seed Use the higher rate when conditions favor poor germination such as cool soil temperatures or low germination seed Sufficient water needs to be added to insure uniform coverage Improper coverage will minimize product performance

Seed Type				
Alfalfa	Corn	Oats	Rye	Tomatoes
Barley	Cotton	Okra	Sorghum	Wheat
Cabbage	Cucumber	Onions	Soybeans	
Canola	Dry Beans	Peanuts	Spinach	
Carrots	Eggplant	Peas	Squash	
Cauliflower	Lettuce	Peppers	Sugarbeets	
Celery	Melons	Rice	Sunflowers	

NOTE If seed has been treated with ASCEND do not apply ASCEND as an in furrow band side dress or mark out application

RED OR WHITE POTATOES

Choose *one* of the following methods

Dip potato seed pieces in a solution of 1 part ASCEND to 375 parts water (0.34 fl oz /gal of water) for 30 to 60 seconds or spray seed pieces with the above solution so that seed pieces are covered and thoroughly wetted ASCEND can be used with a fungicide program

OR

Use 0.50 oz to 1.0 oz (volumetric measurement) which equals 8 grams to 16 grams on a dry basis of ASCEND per 100 lbs of cut seed pieces Treat seed pieces immediately after they have been

14/05

cut Apply so that the cut seed pieces are thoroughly covered ASCEND can be mixed with other seed treatments and carriers such as fir and alder bark to insure uniform coverage

NOTE If seed has been treated with ASCEND do not apply ASCEND as an in furrow band side dress or mark out application

SWEET POTATOES AND YAMS

Dip potato slips in a solution of 1 part ASCEND to 375 parts water (0.34 fl oz /gal of water) for 30 to 60 seconds ASCEND can be used with a fungicide program

NOTE If seed has been treated with ASCEND do not apply ASCEND as an in furrow band side dress or mark out application

MECHANICAL SEED TREATERS

Apply the appropriate amount of ASCEND to a premeasured amount of seed and mix thoroughly until all seed are uniformly coated Seed can be treated in this manner and stored until used for planting Do not use treated seed for food feed or oil purposes An approved dye must be added to distinguish ASCEND treated seed and prevent inadvertent use for food feed or oil purposes Seed treated with this product must be labeled in accordance with all applicable requirements of the Federal and State seed laws **DO NOT USE TREATED SEED FOR FOOD FEED OR OIL PURPOSES**

BROADCAST SEED APPLICATION

Partially fill broadcast spreader with a premeasured amount of seed Apply the appropriate amount of ASCEND diluted with water on the surface of the seed Mix with a stick or paddle until all seed are coated Repeat procedure until broadcast spreader is filled **DO NOT USE TREATED SEED FOR FOOD FEED OR OIL PURPOSES** Treat only those seeds needed for immediate use and planting Do not store excess treated seed beyond planting time

SPECIAL NOTE FOR ALL TRANSPLANTED CROPS

- Two methods are recommended for this program
 - A Dip or spray roots with a solution of 0.75 fluid ounces of ASCEND per gallon of water prior to transplanting
 - B Bedding seedlings may be sprayed or drenched in flats 12-24 hours before transplanting to reduce transplant shock with a solution of 0.75 fluid ounces of ASCEND per gallon of water
- Begin the foliar program two (2) weeks after transplanting A combination of the transplant and foliar spray program is most effective

STORAGE AND DISPOSAL

Do not contaminate water food or feed by storage or disposal

Pesticide Storage Protect from freezing Store out of direct sunlight

Pesticide Disposal Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility

Container Disposal Use label language appropriate for container size and type

Nonrefillable containers Do not reuse or refill this container Clean container promptly after emptying

Nonrefillable container equal to or less than 5 gallons Triple rinse as follows Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip Fill the container 1/4 full with water and recap Shake for 10 seconds Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal Drain for 10 seconds after the flow begins to drip Repeat this procedure two more times Then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill or by incineration or if allowed by state and local authorities by burning If burned stay out of smoke

15/15

Nonrefillable container greater than 5 gallons Triple rinse as follows Empty the remaining contents into application equipment or a mix tank Fill the container ¼ full with water Replace and tighten closures Tip container on its side and roll it back and forth ensuring at least one complete revolution for 30 seconds Stand the container on its end and tip it back and forth several times Tip the container over onto its other end and tip it back and forth several times Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal Repeat this procedure two more times Then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill or by incineration or if allowed by state and local authorities by burning If burned stay out of smoke

Refillable container Refill this container with pesticide only Do not reuse this container for any other purpose Cleaning the container before final disposal is the responsibility of the person disposing of the container Cleaning before refilling is the responsibility of the refiller To clean the container before final disposal empty the remaining contents from this container into application equipment or mix tank Fill the container about 10 percent full with water Agitate vigorously or recirculate water with the pump for 2 minutes Pour or pump rinsate into application equipment or rinsate collection system Repeat this rinsing procedure two more times Then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill or by incineration or if allowed by state and local authorities by burning If burned stay out of smoke

**FOR CHEMICAL EMERGENCY Spill leak, fire, exposure or accident call
CHEMTREC 1 800 424 9300**

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully TO THE EXTENT CONSISTENT WITH APPLICABLE LAW (1) THE GOODS DELIVERED TO YOU ARE FURNISHED AS IS BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES GUARANTEES OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER EITHER EXPRESS OR IMPLIED OR BY USAGE OF TRADE STATUTORY OR OTHERWISE WITH REGARD TO THE PRODUCT SOLD INCLUDING BUT NOT LIMITED TO MERCHANTABILITY FITNESS FOR A PARTICULAR PURPOSE USE OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE UNINTENDED CONSEQUENCES INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS OR THE MANNER OF USE OR APPLICATION INCLUDING WEATHER ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER MANUFACTURER AND SELLER AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL INCIDENTAL OR CONSEQUENTIAL DAMAGES OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD INCLUDING USE APPLICATION HANDLING AND DISPOSAL MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER IF ANY OR FOR ANY DAMAGES OR SUMS OF MONEY CLAIMS OR DEMANDS WHATSOEVER RESULTING FROM OR BY REASON OF OR ARISING OUT OF THE MISUSE OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER USER OR ITS CUSTOMERS BUYER'S OR USER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT

If you do not agree with or do not accept any of directions for use the warranty disclaimers or limitations of liability do not use the product and return it unopened to the Seller and the purchase price will be refunded

AgriSolutions is a trademark and ASCEND is a registered trademark of Winfield Solutions LLC
Mepex is a registered trademark of E I du Pont de Nemours and Co