

9779-357

11/25/2003

1/11



Please read instructions on reverse before completing form.

Form Approved, OMB No. 2070-0080, Approval expires 2-28-95



United States
Environmental Protection Agency
Washington, DC 20460

Registration
 Amendment
 Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number 9779-357	2. EPA Product Manager George Tompkins	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) ProSource One Magic Carpet Fertilizer w Barricade 0.38%	PM#	
5. Name and Address of Applicant (Include ZIP Code) Agriliance, LLC, c/o Alice Walker Consulting 481 Country Club Drive Senatobia, MS 38668 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____	NOTIFICATION NOV 25 2003
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.	
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.	

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Notification of alternate brand name, to wit, Faithful Farms Crabgrass Preventer Fertilizer with Barricade 0.38% per PR Notice 98-10. This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No		<input type="checkbox"/> Metal	<input type="checkbox"/> Plastic
				<input type="checkbox"/> Glass	<input type="checkbox"/> Paper
* Certification must be submitted	If "Yes" Unit Packaging wgt.	No. per container	If "Yes" Package wgt	Other (Specify) _____	
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions <input type="checkbox"/>	
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled			<input type="checkbox"/> Other _____		

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Alice Walker, Ph.D.	Title Agent	Telephone No. (Include Area Code) 662-502-5965
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received (Stamped)
2. Signature 	3. Title Agent	
4. Typed Name Alice Walker, Ph.D.	5. Date November 11, 2003	

2/11

NOTIFICATION
NOV 25 2003

FAITHFUL FARMS™ CRABGRASS PREVENTER FERTILIZER WITH BARRICADE® 0.38%

*For selective preemergence control of grass and broadleaf weeds in established turf grasses
(excluding golf course putting greens) and lawns and landscape ornamentals
(including established perennials & wildflower plantings)*

ACTIVE INGREDIENT:

Prodiamine, [N², N²-Di-n-propyl-2,4-dinitro-6-(trifluoromethyl)-m-phenylenediamine]..... 0.38%

OTHER INGREDIENTS: 99.62%

TOTAL: 100.00%

Barricade® is a registered trademark of Syngenta.

Guaranteed Fertilizer Analysis	
(will vary depending on grade used)	
Total Nitrogen (N).....	00%
0.0% Ammoniacal nitrogen	
0.0% Urea nitrogen	
Available Phosphate (P ₂ O ₅).....	00%
Soluble Potash (K ₂ O).....	00%
Derived from urea, monoammonium phosphate, and potassium chloride.	

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Flush thoroughly with water for several minutes. Get medical attention if irritation persists.

IF ON SKIN: Wash with soap and water. Rinse thoroughly. Get medical attention if irritation persists.

IF INHALED: Remove victim to fresh air.

EPA Reg. No. 9779-357

EPA Est. No. _____

Distributed by:
AGRILIANCE, LLC
P.O. BOX 64089
St. Paul, MN 55164-0089

Net Weight: _____

10/30/03D

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Causes moderate eye injury (irritation). Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Prolonged or frequently repeated skin contact while handling the material may cause allergic reactions in some individuals.

ENVIRONMENTAL HAZARDS

This product has low solubility in water. At the limits of solubility, this product is not toxic to fish. However, at concentrations above the level of water solubility, it may be toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

If material is spilled: Contain and sweep up material of spill for disposal or use per label instructions. Wash cleanup equipment to prevent contamination elsewhere. Excess quantities above normal application rates could result in damage to plants subject to spill. Water-in remainder, disc under, cover with layer of soil, or leave exposed to sunlight, as deemed appropriate to the situation.

Storage: Store in original container away from other fertilizer, feed, or food stuffs and separated from other pesticides.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal:

Paper Containers: Completely empty container into application equipment. Then dispose of empty container in a sanitary landfill or at an incineration facility, or, if allowed by state and local authorities, by burning locally. If burned, stay out of smoke.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

This product is a selective preemergence herbicide that provides residual control of many grass and broadleaf weeds in:

- Established turf grasses (excluding golf course putting greens) and lawns
- Landscape ornamentals
- Established perennials & wildflower plantings

This product controls susceptible weeds by inhibiting weed seed germination and root development. Most effective weed control will be obtained when this product is activated by at least 0.5 inches of rainfall or irrigation, or with shallow (1-2 inches) incorporation, prior to weed seed germination and within 14 days following application.

Not for use on plants being grown (1) for sale or other commercial use, (2) for commercial seed production, or (3) for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.

Do not graze or feed livestock forage cut from areas treated with this product.

4/11

Do not apply this product through any type of irrigation system.

Do not apply aerially.

Do not apply to golf course putting greens.

Failure to follow the directions for use and precautions on this label may result in poor weed control, crop injury, or illegal residues.

APPLICATION DIRECTIONS


Apply uniformly through suitable, calibrated application equipment.

ESTABLISHED TURF

This product is a selective preemergence herbicide that, when properly applied, will control certain grass and broadleaf weeds in established turf grasses and lawns. The maximum amount of this product that may be applied per year is given for each turf grass species in the "Maximum Annual Rates" section of this label.

Most effective weed control in turf grasses will be obtained when this product is activated by at least 0.5 inches of rainfall or irrigation prior to weed seed germination and within 14 days following application. See the map below for approximate crabgrass seed germination dates.

APPROXIMATE CRABGRASS SEED GERMINATION DATES



USE PRECAUTIONS

The following precautions apply to the use of this product in turf grasses and lawns:

- (1) Application of this product may thin emerged annual bluegrass and newly overseeded grasses.

(2) Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil.

(3) Do not cut (harvest) treated sod before 120 days after application. Do not apply to newly set sod until the following year.

(4) Application of this product to turf stressed by drought, low fertility, or pest damage may result in turf injury.

(5) Disturbing the herbicide barrier with cultural practices such as disking may result in reduced weed control.

(6) Do not apply this product to putting greens or areas where dichondra, colonial bentgrass, velvet bentgrass, or annual bluegrass (*Poa annua*) are desirable species.

RATES OF APPLICATION

This product may be applied as a single application or in sequential applications to control weeds germinating throughout the year. All applications must be made prior to germination of the target weeds. This product will not control established weeds.

Maximum use rate selection should be based on turf species. The length of time of residual weed control provided by this product is related to the rate applied.

MAXIMUM ANNUAL RATES

This product is recommended for use on the turf grass species listed in the following table. Do not exceed the maximum yearly rate as given in the following table:

Maximum Application Rate Per Calendar Year By Turf Grass Species ¹		
Turf Species	Lbs. Product/A	Lbs. ai/A
Bermudagrass ² Bahiagrass Centipedegrass Seashore Paspalum St. Augustinegrass Tall Fescue (including turf-type) Zoysia	400 ³	1.5
Buffalograss Kentucky Bluegrass Perennial Ryegrass	267	1.0
Creeping Red Fescue	200	0.75
Creeping Bentgrass	175	0.65

¹ These are the maximum rates per calendar year by species limitations.

² May be used on newly sprigged or plugged Bermudagrass at rates not to exceed 139 lbs./A or 3.1 lbs./1000 sq. ft. Newly sprigged or plugged Bermudagrass stolon rooting may be temporarily retarded. Suppression only of Foxtail, Goosegrass and Rescuegrass due to reduced product rates used in sprigging situations.

³ Do not apply more than 400 lbs. of product per calendar year.

- Use higher rates of this product to achieve higher levels of fertility and longer periods of weed control for each turf type, but do not exceed the "Maximum Annual Rates" table.

WEEDS CONTROLLED

When used in accordance with this label this product will provide control of the following weeds:

Barnyardgrass	Carpetweed
Bluegrass, Annual (<i>Poa annua</i>)	Chickweed, Common
Crabgrass (large, smooth)	Chickweed, Mouseear (from seed)
Crowfootgrass	Henbit
Cupgrass, Woolly	Knotweed
Foxtails, Annual	Kochia
Goosegrass ¹	Lambsquarter, Common
Itchgrass	Pigweed
Johnsongrass (from seed)	Purslane, Common
Junglerice	Pusley, Florida
Lovegrass	Shepherdspurse ²
Panicum (Texas, Fall, Browntop)	Speedwell, Persian
Rescuegrass ³	Spurge, Prostrate
Signalgrass, Broadleaf	Witchgrass
Sprangletop	Woodsorrel, Yellow (from seed)

¹ In many areas a single application of 175 to 400 lbs./A of this product will control goosegrass. However, under heavy goosegrass pressure and/or an extended growing season, most effective weed control may be obtained by making an initial application of 175 to 267 lbs./A followed after 60 to 90 days by a second application at doses that would not exceed those given in the "Maximum Annual Rates" table. Do not exceed the maximum rate for turf grass species listed in the "Maximum Annual Rates" table above.

² Applications for this weed should be made in late summer, fall, or winter prior to germination.

³ Suppression only.


Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rates recommended for each turf species. All applications must be made prior to germination of the weed seeds.

WHEN TO APPLY AFTER OVERSEEDING TURF

Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil.

WHEN TO OVERSEED AFTER APPLICATION

This product will inhibit the germination of turf species if overseeded too soon after application. Follow rates and intervals in the table below for best overseeding/reseeding results.



Rate (lb./A)		Months Before Overseeding		
Product	ai	North	Transition	South
175	.50	4	4	4
175	.66	5	4	4
175	.74	6	5	5
215	.80	---	6	6
267	1.00	---	7	7
305	1.14	---	---	9
345	1.30	---	---	10
400	1.50	---	---	12

LANDSCAPE ORNAMENTALS

(including Established Perennials and Wildflower Plantings)

This product may be applied for residual preemergence weed control in ornamentals.

USE RATES

Apply this product at 175 to 400 lbs./A in fall and/or spring. Use higher rates of application for longer control periods. Sequential or single applications are allowed so long as the total amount of product applied does not exceed the maximum annual application rate of 400 lbs./A.

APPLICATION TIMING AND INFORMATION

This product is a preemergence herbicide and will not control emerged weeds. Most effective weed control in ornamentals will be obtained when this product is activated by at least 0.5 inches of rainfall or irrigation, or with shallow (1 to 2 inches) mechanical incorporation, prior to weed seed germination and within 14 days following application.

Best weed control is obtained when this product is applied to soil free of clods, weeds, and debris such as leaves. Prior to application of this product, control existing vegetation by hand weeding, cultivation or the use of an appropriate postemergence herbicide.

ESTABLISHED LANDSCAPE ORNAMENTALS

This product may be applied to established ornamentals as a broadcast, over the top, or directed application. Irrigation or rainfall soon after application will wash residues off plant foliage and activate this product in the soil.

NEWLY TRANSPLANTED LANDSCAPE ORNAMENTALS

This product may be applied to newly transplanted ornamentals as a broadcast, over the top, or directed application. Delay application to allow soil to settle around new transplants and water thoroughly before applying this product. Apply this product after cuttings form roots and become established. Apply before budding/grafting or after buds/grafts have taken to avoid any inhibition of the tissue union.

TOLERANT ORNAMENTAL SPECIES

This product will not harm most trees, shrubs, vines, and flowers. The species listed below are tolerant to this product. Best results will be obtained when this product is uniformly applied to the soil surface; avoid applying to ornamental shoots which may defeat the effort to get uniform distribution on the soil surface. This product may be applied over the top of these species.

When plants are under stress (such as heat, drought, or frost damage), some cultivars of listed plants may be sensitive to this product.

Scientific name

Abies spp.
Abelia grandiflora
Acer palmatum
A. platanoides
Adiantum spp.
Adiantum chinensis
Agapanthus africanus
Agapanthus orientalis
Akebia quinata
Allium cernuum
Anemone hybrida
Aquilegia spp.
Arctostaphylos densiflora
Arctotheca calendula
Artemisia spp.
Aster spp.
Aster X Frikartii
Athrium Filiz-femina
Aucuba japonica
Begonia spp.
Berberis gladwynensis
B. julianae
B. mentorensis
B. thunbergii
B. verruculosa
Bergenia cordifolia
Boltonia asteroides
Bougainvillea spp.
Buddleia davidii
Buxus microphylla
Callistemon viminalis
Calluna vulgaris
Campanula carpatica
Campis X Tagliabuana

Carpobrotus edulis
Cassia artemisoides
Ceanothus rigidus
Ceratostigma plumbagonoides
Chamaecyparis pisifera
Chrysanthemum nipponicum
Cleyera japonica
Citrus spp.
Coreopsis spp.

Cornus florida
C. stolonifera
Cortaderia selloana
Cotoneaster apiculatus
C. buxifolius
C. dammeri
C. microphyllus
Crotagrus spp.
Crocosmia spp.
Cupressus sempervirens

Common name

Fir species** (Balsam, Fraser, Noble, etc.)
 Abelia: Sherwood
 Japanese Maple
 Norway Maple
 Yarrow: King Edward
 Kiwi*
 Lily-of-the-Nile (African Lily)

 Five-Leaf or Chocolate Vine
 Lady's Leek, Nodding Onion
 Japanese Anemone
 Aquilegia: Red and Gold
 Vine Hill Manzanita
 Cape Weed
 Wormwood: Silver Mound, Castle
 Aster: Bonny Blue, Purple Dome

 Lady Fern: Fern Lady
 Japanese Aucuba
 Fibrous Begonia: Hardy Grandis
 Barberry
 Wintergreen Barberry
 Mentor Barberry
 Japanese Barberry
 Warty Barberry

 Snowbank
 Bougainvillea
 Butterfly-Bush (Dwarf Blue): Royal Red
 Japanese Boxwood
 Weeping Bottlebrush
 Scotch Heather
 Tussock Bellflower, (White Clips)
 Trumpet Creeper, Trumpet Flower,
 Madame Galen
 Hottentot Fig (Ice Plant)
 Feathery Cassia
 Wild Lilac

 False Cypress

 Cleyera
 Citrus species*
 Coreopsis (Calliopsis): Early Sunrise,
 Moonbeam
 Flowering Dogwood
 American Dogwood
 Pampas Grass
 Cranberry Cotoneaster
 Cotoneaster
 Bearberry Cotoneaster
 Rockspray Cotoneaster
 Hawthorne
 Lucifer
 Italian Cypress

Delosperma alba
Delosperma spp.
Delphinium spp.
Dianthus deltooides
D. gratianopolitanus
Dodonea viscosa
Echinacea purpurea
Elaeagnus pungens
Euonymus fortunei
E. japonica
E. kiautschovica
Fatsia japonica
Forsythia intermedia
F. suspensa
F. viridissima
Gaillardia spp.
Gardenia jasminoides
Gaura spp.
Gentiana dahurica
Geranium cinereum
Gladiolus spp.
Gypsophila repens
Hedera helix
Hellanthemum spp.
Hemerocallis spp.



Hibiscus
Hibiscus spp.
Hibiscus rosa-sinensis
Hosta plantaginea
H. Sieboldiana
Houttuynia cordata var. *variegata*
Hydrangea macrophylla
Ilex cornuta
I. crenata
I. opaca
I. pernyi
I. vomitoria
Inula ensifolia
Iris ensata
Iris siberica
Iris spp.
Jasminum nudiflorum
Juniperus chinensis
J. conferta
J. horizontalis
Juglans spp.
Justicia brandegeana
Lagerstromia indica
Lagerstromia indica X Fauriei
Lantana montevidensis
Lavender spp.
Leontopodium alpinum
Ligustrum sinense
Ligustrum amurense
L. japonicum

White Trailing Ice Plant
 Cooperi Pink
 Larkspur: Blue Elf
 Dianthus: Maiden Pinks "Zing"
 Cheddar Pink
 Hop Bush
 Coneflower, Purple; Magnu
 Silverberry
 Wintercreeper
 Japanese Spindle Tree (Evergreen Euonymus)
 Spreading Euonymus
 Japanese Aralia
 Border Forsythia
 Weeping Forsythia
 Greenstem Forsythia
 Gaillardia, Blanket Flower: "Goblin"
 Gardenia, Cape-Jasmine, Radicans

Gentian
 Cranesbill
 Gladiolus species**
 Baby's Breath
 English Ivy
 Sunrose
 Daylily: Aztec Gold, Stella de Oro, Tender Love
 Rose of Sharon**
 Mallow: Disco Belle White
 Chinese Hibiscus
 Hosta, Plantain Lily (Fragrant)
 Hosta, "Searsucker"

Bigleaf Hydrangea
 Chinese Holly
 Japanese Holly: Helleri
 American Holly
 Holly
 Yaupon Holly: Schillings

Sword-Leaved Iris: Jodlesong
 Siberian Iris: Cabernet
 Iris species**
 Winter Jasmine
 Chinese Juniper: Nick's Compact, Parsonnii
 Shore Juniper: Blue Pacific
 Creeping Juniper
 Walnut*
 Shrimp Plant
 Crape Myrtle
 Crape Myrtle: Tuscarora
 Weeping Lantana
 Lavender: Munstead
 Edelweiss
 Chinese Privet: Variegata
 Amur Privet
 Japanese Privet



<i>L. lucidum</i>	Glossy Privet (Waxleaf)
<i>Lilium</i> spp.	Lily; Jazz
<i>Liriope muscari</i>	Big Blue Lily; Turf; Silver Mound; Evergreen Giant
<i>Liriope Muscari var. variegata</i>	Giant
<i>Liriope spicata</i>	Liriope, Variegated
<i>Lonicera Cardinalis</i>	Liriope, Creeping
<i>Lonicera japonica</i>	Cardinal Flower, Indian Pink
<i>L. tatarica</i>	Japanese Honeysuckle
<i>Loropetalum chinense</i>	Tatarian Honeysuckle
<i>Lethrum</i> spp.	Burgundy
<i>Magnolia</i> spp.	Loosestrife Modern Pink
<i>Malephora luteola</i>	Magnolia species**
<i>Maius</i> spp.	Ice Plant
<i>Miscanthus sinensis</i>	Crabapple*
<i>Nandina domestica</i>	Yaku Jima, Silberfeder
<i>Narcissus</i> spp.	Heavenly Bamboo
<i>Nerium oleander</i>	Narcissus species**
<i>Olea europaea</i>	Oleander
<i>Ophiopogon japonicus</i>	Olive*
<i>Osmanthus heterophyllus</i>	Mondo Grass
<i>Osteospermum fruticosum</i>	Osmanthus (False Holly): Gulf Tide
<i>Oxycandrum arboreum</i>	Trailing African Daisy
<i>Paeonia suffruticosa</i>	Sourwood
<i>Pennisetum setaceum</i>	Tree Peony
<i>Perovskia atriplicifolia</i>	Fountain Grass (Dwarf)
<i>Persea americana</i>	Avocado*
<i>Photinia fraseri</i>	Frasier's Photinia (Redtip)
<i>Piptotegia Virginiana</i>	Dragonhead, False; Vivid
<i>Picea</i> spp.	Spruce species**(Colorado Blue, Norway, etc.)
<i>Pieris japonica</i>	Lily-of-the-Valley Bush
<i>Pinus brutia</i>	Calabrian Pine
<i>P. canariensis</i>	Canary Island Pine
<i>P. eiliottii</i>	Slash Pine
<i>P. halepensis</i>	Aleppo Pine
<i>P. nigra</i>	Austrian Pine
<i>P. palustris</i>	Longleaf Pine
<i>P. radiata</i>	Monterey Pine
<i>P. thunbergiana</i>	Japanese Black Pine
<i>P. strobus</i>	Eastern White Pine
<i>P. sylvestris</i>	Scotch Pine
<i>P. taeda</i>	Loblolly Pine
<i>P. virginiana</i>	Virginia Pine
<i>Pistachio</i> spp.	Pistachio*
<i>Pittosporum rhombifolium</i>	Queensland Pittosporum
<i>P. tobira</i>	Japanese Pittosporum
<i>Podocarpus macrophyllus</i>	Japanese Yew
<i>Prunus laurocerasus</i>	English Laurel
<i>Prunus</i> spp.	Almond, Apricot, Nectarine, Peach, Plum and Prune*
<i>Pseudotsuga menziesii</i>	Douglas Fir**
<i>Pyraecantha coccinea</i>	Firethorn (Scarlet)
<i>P. fortuneana</i>	Firethorn
<i>P. koidzumii</i>	Firethorn
<i>Pyrus</i> spp.	Bradford Pear sp.

<i>Quercus shumardii</i>	Oak, Shumard's Red
<i>Quercus</i> spp.	Oak species
<i>Ranunculus indica</i>	Indian Hawthorne
<i>Ranunculus umbellata</i>	Yedda Hawthorn
<i>Rhododendron</i>	Coral Bells, Delaware Valley White, Flame
(including Azalea)	Creeper, Formosa, George L. Tabor, Girard
	Crimson, Hino-crimson, PJM, Roseum
	Elegans, Wakeiebisu, White Gumpo
<i>Rosmarinus officinalis</i>	Rosemary*
<i>Rudbeckia</i> spp.	Black-Eyed Susan: Goldstrum
<i>Santolina virens</i>	Santolina
<i>Saxifraga</i> spp.	Saxifrage: Purple Dome
<i>Scabiosa</i> spp.	Pincushion Flower
<i>Sedum album</i>	Stonecrop
<i>Sedum caudicola</i>	Stonecrop: Lidakense
<i>Sedum dasyphyllum</i>	Stonecrop
<i>Sedum spurium</i>	Stonecrop: Dragon's Blood
<i>Spiraea bumalda</i>	Spirea: Anthony Waterer
<i>Strygium paniculatum</i>	Australian Brushcherry
<i>Taxus cuspidata</i>	Japanese Yew
<i>T. media</i>	Yew
<i>Tenarium</i> spp.	Germander
<i>Thalictrum dipterocarpum</i>	Meadow Rue
<i>Thuja occidentalis</i>	American Arborvitae
<i>Trachelospermum asiaticum</i>	Star Jasmine
<i>Tsuga canadensis</i>	Canada Hemlock
<i>Tulipa</i> spp.	Tulip species
<i>Veronica</i> spp.	Veronica, Speedwell: Sunny Border
<i>Viburnum japonicum</i>	Japanese Viburnum
<i>V. odoratissimum</i>	Sweet Viburnum
<i>V. plicatum</i>	Japanese Snowball
<i>V. rigidum</i>	Canary Island Viburnum
<i>V. suspensum</i>	Arrowood Viburnum
<i>V. tinus</i>	Laurustinus
<i>V. trilobatum</i>	Cranberry Bush
<i>V. wrightii</i>	Leatherleaf Viburnum
<i>Vinca major</i>	Greater Periwinkle, Vinca
<i>Vinca minor</i>	Common Periwinkle, Vinca
<i>Vitis</i> spp.	Grape*
<i>Weigela florida</i>	Old Fashioned Weigela
<i>Yucca aloifolia</i>	Spanish Bayonet, Yucca
<i>Y. filamentosa</i>	Adam's Needle, Yucca

*Do not use on food-producing trees, vines, or plants.

**Not for use on container-grown plants.

Notice of Warranty: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, NOR IS ANY REPRESENTATIVE OF SELLER AUTHORIZED TO MAKE ANY SUCH WARRANTY OR MODIFY THESE TERMS. This warranty does not extend to the storage, handling or use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to Seller, and Buyer assumes the risk of any such storage, handling or use. Seller shall not be responsible for incidental or consequential damages, if any, resulting from a breach of warranty.