

9779-357

3/8/2010

10A/4



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Alice Walker, Ph.D.
Product Registrations
Winfield Solutions, LLC
3094 Country Club Road
Senatobia, MS 38668

MAR 8 2010

SUBJECT: Application for Pesticide Notification (PRN 98-10)
Request Alternate Brand Name "Prolific® Fertilizer with Barricade 0.38%"
EPA Reg. No. 9779-357
Application Dated January 26, 2010

Dear Registrant:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 dated 01/26/10 for the above product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action(s) requested fall within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions, please call me directly at 703-305-6249 or Owen F. Beeder of my staff at 703-308-8899.

Sincerely,

A handwritten signature in black ink, appearing to read "Linda Arrington".

Linda Arrington
Notifications & Minor Formulations Team Leader
Registration Division (7505P)
Office of Pesticide Programs

2014

Please read instructions on reverse before completing form.

Form Approved: OMB No. 2070-0060, Approval expires 2-28-95



United States
Environmental Protection Agency
Washington, DC 20460

<input type="checkbox"/>	Registration
<input type="checkbox"/>	Amendment
<input checked="" type="checkbox"/>	Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number 9779-357	2. EPA Product Manager Kathryn Montague	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Magic Carpet Fertilizer with Barricade 0.38%	PM# 23	
5. Name and Address of Applicant (Include ZIP Code) Winfield Solutions, LLC, c/o Alice Walker 3094 Country Club Road Senatobia, MS 38668 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

NOTIFICATION

MAR 8 - 2010

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Notification of alternate brand name, Prolific Fertilizer with Barricade 0.38%. This notification is consistent with the provisions of PR Notices 98-10 and 07-4 and EPA regulations at 40 CFR 152.46, 156.10, 156.140, 156.144, 156.146, and 156.156 and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notices 98-10 and 07-4 and 40 CFR 152.46, 156.10, 156.140, 156.144, 156.146, and 156.156 this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No		<input type="checkbox"/> Metal	<input checked="" type="checkbox"/> Plastic
* Certification must be submitted		If "Yes" Unit Packaging wgt. No. per container	If "Yes" Package wgt. No. per container	<input type="checkbox"/> Glass	<input checked="" type="checkbox"/> Paper
3. Location of Net Contents Information <input checked="" type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container 50 lbs.		5. Location of Label Directions <input checked="" type="checkbox"/> On label	
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph Paper glued Stenciled			<input checked="" type="checkbox"/> Other Flexography		

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Alice Walker, Ph.D.	Title Registration Specialist	Telephone No. (Includes Area Code) 662-562-5995
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received (Stamped)
2. Signature <i>Alice Walker</i>	3. Title Registration Specialist	
4. Typed Name Alice Walker, Ph.D.	5. Date 1-26-10	



30914

January 26, 2010

Document Processing Desk (Notif)
Office of Pesticide Programs (7504P)
U.S. Environmental Protection Agency
1200 Pennsylvania Ave., NW
Washington, DC 20460

**Re: Magic Carpet™ Fertilizer with Barricade 0.38%
EPA Reg. No. 9779-357**

This correspondence will constitute notification from Winfield Solutions, LLC to add the following alternate brand name to the above-referenced product registration:

Prolific® Fertilizer with Barricade 0.38%

To this end, please find enclosed one copy of new labeling along with the appropriate application form.


I believe this notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under section 12 and 14 of FIFRA.

Thank you for adding this notification to the registration records for this product.

Sincerely,

Alice Walker, Ph.D.
Registration Specialist

Enclosures



40x14

ProSource One

PROLIFIC® FERTILIZER WITH BARRICADE® 0.38%

ACTIVE INGREDIENT:

Prodiamine¹0.38%

OTHER INGREDIENTS:99.62%

TOTAL100.00%

¹ CAS No. 29091-21-2

This product is a preemergence herbicide that provides residual control of many grass and broadleaf weeds in:

NOTIFICATION

MAR 8 - 2010

- Established turf grasses (excluding golf course putting greens) and lawns
- Landscape ornamentals
- Established perennials & wildflower plantings

KEEP OUT OF REACH OF CHILDREN
CAUTION

FIRST AID

If Swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If in Eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes; then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on Skin or Clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If Inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-424-7452 for emergency medical treatment information.

EPA Reg. No. 9779-357

EPA Est. No. _____

Distributed by:
Winfield Solutions, LLC
P.O. Box 64589
St. Paul, MN 55164-0589

Net Weight: _____

1/0126/0

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Causes moderate eye injury (irritation). Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Prolonged or frequently repeated skin contact while handling the material may cause allergic reactions in some individuals.

ENVIRONMENTAL HAZARDS

This product has low solubility in water. At the limits of solubility, this product is not toxic to fish. However, at concentrations above the level of water solubility, it may be toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

If material is spilled: Contain and sweep up material of spill for disposal as per label instructions. Wash cleanup equipment to prevent contamination elsewhere. Excess quantities above normal application rates could result in damage to plants subject to spill. Water-in remainder, disc under, cover with layer of soil, or leave exposed to sunlight, as deemed appropriate to the situation.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE

Store in original container away from other fertilizer, feed, or foodstuffs and separated from other pesticides.

PESTICIDE DISPOSAL

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL

Paper and plastic bags: Nonrefillable Container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available or dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.


DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Not for use on plants being grown for sale or other commercial use, for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.

This product is a preemergence herbicide that provides residual control of many grass and broadleaf weeds in established turf grasses (excluding golf course putting greens) and lawns, landscape ornamentals, and established perennials & wildflower plantings.

This product controls susceptible weeds by inhibiting weed seed germination and root development. Most effective weed control will be obtained when this product is activated by at least 0.5 inches of rainfall or irrigation, or with shallow (1-2 inches) incorporation, prior to weed seed germination and within 14 days following application.

Failure to follow the directions for use and precautions on this label may result in poor weed control, crop injury, or illegal residues.



APPLICATION DIRECTIONS

This product must be applied prior to crabgrass seed germination and prior to germination of seed of the other weeds listed on this label. Apply uniformly through suitable, calibrated application equipment.


ESTABLISHED TURF

This product is a selective preemergence herbicide that, when properly applied, will control certain grass and broadleaf weeds in established turf grasses and lawns. The maximum amount of this product that may be applied per year is given for each turf grass species in **Table 1. Maximum Annual Rates** on this label.

Most effective weed control in turf grasses will be obtained when this product is activated by at least 0.5 inches of rainfall or irrigation prior to weed seed germination and within 14 days following application. See the map below for approximate crabgrass seed germination dates.

CRABGRASS GERMINATION DATES


Approximate Date



USE PRECAUTIONS

The following precautions apply to the use of this product in turf grasses and lawns:

1. Do not graze or feed livestock forage cut from areas treated with this product.
2. Do not use on food producing plants.
3. Do not apply this product through any type of irrigation system.
4. Do not apply aerially.
5. Do not apply to golf course putting greens.
6. Application of this product may thin emerged annual bluegrass and newly overseeded grasses.
7. Do not apply to over seeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if applied before seedling secondary roots are in the second inch of soil, not thatch plus soil.
8. Do not cut (harvest) treated sod before 120 days after application. Do not apply to newly set sod until the following year.



- 9. Application of this product to turf stressed by drought, low fertility, or pest damage may result in turf injury.
- 10. Disturbing the herbicide barrier with cultural practices, such as disking may result in reduced weed control.
- 11. Do not apply this product to putting greens or areas where dichondra, colonial bentgrass, velvet bentgrass, or annual bentgrass (*Poa annua*) are desirable species.
- 12. Keep this product off sidewalks, patios, driveways, etc. This product contains materials which will stain concrete and painted surfaces. If it comes in contact with a concrete surface, sweep off immediately before flushing with water. If concrete is wet when contact occurs, staining may be impossible to prevent. Be careful when using rotary spreaders which scatter product beyond lawn edge and onto concrete.

RATES OF APPLICATION

This product may be applied as a single application or in sequential applications to control weeds germinating throughout the year. All applications must be made prior to germination of the target weeds. This product will not control established weeds.

Do not exceed the Maximum Annual Rate per calendar year (See Table 1.).

Maximum use rate selection should be based on turf species. The length of time of residual weed control provided by this product is related to the rate applied.

MAXIMUM ANNUAL RATES

This product is recommended for use on the turf grass species listed in the following table. Do not exceed the maximum yearly rate as given in the following table:


Table 1 Maximum Application Rates ¹		
Turf Species	Lbs. Product/A	Lbs. ai/A
Bermudagrass ² Bahia grass Centipedegrass Seashore Paspalum St. Augustinegrass Tall Fescue (including turf-type) Zoysia	400 ³	1.5
Buffalograss Kentucky Bluegrass Perennial Ryegrass	267	1.0
Creeping Red Fescue	200	0.75
Creeping Bentgrass	175	0.65

¹ These are the maximum rates per calendar year by species limitations.

² May be used on newly sprigged or plugged Bermudagrass at rates not to exceed 139 lbs./A. or 3.1 lbs./1000 sq. ft. Newly sprigged or plugged Bermudagrass stolon rooting may be temporarily retarded. Suppression only of Foxtail, Goosegrass and Rescuegrass due to reduced product rates used in sprigging situations.

³ Do not apply more than 400 lbs. of product per calendar year.

- Use higher rates of this product to achieve higher levels of fertility and longer periods of weed control for each turf type, but do not exceed the maximum application rates specified in **Table 1. Maximum Application Rates.**



WEEDS CONTROLLED

When used in accordance with this label this product will provide control of the following weeds:

- | | |
|--|---------------------------------|
| Barnyardgrass | Carpetweed |
| Bluegrass, Annual (<i>Poa annua</i>) | Chickweed, Common |
| Crabgrass (large, smooth) | Chickweed, Mouseear (from seed) |
| Crowfootgrass | Henbit |
| Cupgrass, Woolly | Knotweed |
| Foxtails, Annual | Kochia |
| Goosegrass ¹ | Lambsquarter, Common |
| Itchgrass | Pigweed |
| Johnsongrass (from seed) | Purslane, Common |
| Junglerice | Pusley, Florida |
| Lovegrass | Shepherdspurse ² |
| Panicum (Texas, Fall, Browntop) | Speedwell, Persian |
| Rescuegrass ³ | Spurge, Prostrate |
| Signalgrass, Broadleaf | Witchgrass |
| Sprangletop | Woodsorrel, Yellow (from seed) |

¹ In many areas a single application of 175 to 400 lbs./A of this product will control goosegrass. However, under heavy goosegrass pressure and/or an extended growing season, most effective weed control may be obtained by making an initial application of 175 to 267 lbs./A followed after 60 to 90 days by a second application at doses that would not exceed those given in **Table 1. Maximum Application Rates.** Do not exceed the maximum rate for turf grass species listed in **Table 1. Maximum Application Rates.**

² Applications for this weed should be made in late summer, fall, or winter prior to germination.

³ Suppression only.

Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rates recommended for each turf species. All applications must be made prior to germination of the weed seeds.

WHEN TO APPLY AFTER OVERSEEDING TURF


Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil.

WHEN TO OVERSEED AFTER APPLICATION

This product will inhibit the germination of turf species if overseeded too soon after application. Follow rates and intervals in the table below for best overseeding/reseeding results.

TABLE 2. Months After Application Before Overseeding.

Rate Lbs. Product/A	North	Transition	South
132	4	4	4
175	5	4	4
195	6	5	5
213	---	6	6
267	---	7	7
305	---	---	9



345	---	---	10
400	---	---	12

LANDSCAPE ORNAMENTALS

(including Established Perennials and Wildflower Plantings)

This product may be applied for residual preemergence weed control in ornamentals.

USE RATES

Apply this product at 175 to 400 lbs./A in fall and/or spring. Use higher rates of application for longer control periods. Sequential or single applications are allowed so long as the total amount of product applied does not exceed the maximum annual application rate of 400 lbs./A.

APPLICATION TIMING AND INFORMATION

Apply uniformly to the surface of the soil in ornamental planting beds with a granular applicator. Calibrate granular applicator per manufacturer's instructions to assure appropriate dosage. To assure that the product is applied at the proper rates, apply at a rate of 4 lbs. per 1,000 sq. ft. or for flowerbeds, use 6.4 oz. per 100 sq. ft.

This product is a preemergence herbicide and will not control emerged weeds. Most effective weed control in ornamentals will be obtained when this product is activated by at least 0.5 inches of rainfall or irrigation, or with shallow (1 to 2 inches) mechanical incorporation, prior to weed seed germination and within 14 days following application.

Best weed control is obtained when this product is applied to soil free of clods, weeds, and debris such as leaves. Prior to application of this product, control existing vegetation by hand weeding, cultivation or the use of an appropriate postemergence herbicide.

ESTABLISHED LANDSCAPE ORNAMENTALS

This product may be applied to established ornamentals as a broadcast, over the top, or directed application. Irrigation or rainfall soon after application will wash residues off plant foliage and activate this product in the soil.

NEWLY TRANSPLANTED LANDSCAPE ORNAMENTALS

This product may be applied to newly transplanted ornamentals as a broadcast, over the top, or directed application. Delay application to allow soil to settle around new transplants and water thoroughly before applying this product. Apply this product after cuttings form roots and become established. Apply before budding/grafting or after buds/grafts have taken to avoid any inhibition of the tissue union.


TOLERANT ORNAMENTAL SPECIES

This product will not harm most trees, shrubs, vines, and flowers. The species listed below are tolerant to this product. Best results will be obtained when this product is uniformly applied to the soil surface; avoid applying to ornamental shoots which may defeat the effort to get uniform distribution on the soil surface. This product may be applied over the top of these species.

When plants are under stress (such as heat, drought, or frost damage), some cultivars of listed plants may be sensitive to this product.

- Scientific name**
Abies spp.
Abelia grandiflora
Acer palmatum
A. platanoides
Achillea spp.
Actinidia chinensis

- Common name**
 Fir species** (Balsam, Fraser, Noble, etc.)
 Abelia: Sherwood
 Japanese Maple
 Norway Maple
 Yarrow: King Edward
 Kiwi*



<i>Agapanthus africanus</i>	Lily-of-the-Nile (African Lily)	
<i>Agapanthus orientalis</i>		
<i>Akebia quintata</i>	Five-Leaf or Chocolate Vine	
<i>Allium cernuum</i>	Lady's Leek, Nodding Onion	
<i>Anemone hybrida</i>	Japanese Anemone	
<i>Aquilegia</i> spp.	Aquilegia: Red and Gold	
<i>Arctostaphylos densiflora</i>	Vine Hill Manzanita	
<i>Arctotheca calendula</i>	Cape Weed	
<i>Artemisia</i> spp.	Wormwood: Silver Mound, Castle	
<i>Aster</i> spp.	Aster: Bonny Blue, Purple Dome	
<i>Aster X Frikartii</i>		
<i>Athrium Filiz-femina</i>	Lady Fern: Fern Lady	
<i>Aucuba japonica</i>	Japanese Aucuba	
<i>Begonia</i> spp.	Fibrous Begonia: Hardy Grandis	
<i>Berberis gladwynensis</i>	Barberry	
<i>B. julianae</i>	Wintergreen Barberry	
<i>B. mentorensis</i>	Mentor Barberry	
<i>B. thunbergii</i>	Japanese Barberry	
<i>B. verruculosa</i>	Warty Barberry	
<i>Bergenia cordifolia</i>		
<i>Boltonia asteroides</i>	Snowbank	
<i>Bougainvillea</i> spp.	Bougainvillea	
<i>Buddleia davidii</i>	Butterfly-Bush (Dwarf Blue): Royal Red	
<i>Buxus microphylla</i>	Japanese Boxwood	
<i>Callistemon viminalis</i>	Weeping Bottlebrush	
<i>Calluna vulgaris</i>	Scotch Heather	
<i>Campanula carpatica</i>	Tussock Bellflower, (White Clips)	
<i>Campis X Tagliabuana</i>	Trumpet Creeper, Trumpet Flower, Madame Galen	
<i>Carpobrotus edulis</i>	Hottentot Fig (Ice Plant)	
<i>Cassia artemisoides</i>	Feathery Cassia	
<i>Ceanothus rigidus</i>	Wild Lilac	
<i>Ceratostigma plumbagonoides</i>		
<i>Chamaecyparis pisifera</i>	False Cypress	
<i>Chrysanthemum nipponicum</i>		
<i>Cleyera japonica</i>	Cleyera	
<i>Citrus</i> spp.	Citrus species*	
<i>Coreopsis</i> spp.	Coreopsis (Calliopsis): Early Sunrise, Moonbeam	
<i>Cornus florida</i>	Flowering Dogwood	
<i>C. stolonifera</i>	American Dogwood	
<i>Cortaderia selloana</i>	Pampas Grass	
<i>Cotoneaster apiculatus</i>	Cranberry Cotoneaster	
<i>C. buxifolius</i>	Cotoneaster	
<i>C. dammeri</i>	Bearberry Cotoneaster	
<i>C. microphyllus</i>	Rockspray Cotoneaster	
<i>Cretaegus</i> spp.	Hawthorne	o o o o
<i>Crocosmia</i> spp.	Lucifer	o o o o
<i>Cupressus sempervirens</i>	Italian Cypress	o o o o
<i>Delosperma alba</i>	White Trailing Ice Plant	o o o o o o
<i>Delosperma</i> spp.	Cooperi Pink	o o o o
<i>Delphinium</i> spp.	Larkspur: Blue Elf	o o o o o o
<i>Dianthus deltoides</i>	Dianthus: Maiden Pinks "Zing"	o o o o
<i>D. gratianopolitanus</i>	Cheddar Pink	o o o o
<i>Dodonea viscosa</i>	Hop Bush	o o o o
<i>Echinacea pupurea</i>	Coneflower, Purple; Magnus	o o o o o o
<i>Elaeagnus pungens</i>	Silverberry	o o o o

Euonymus fortunei
 E. japonica
 E. kiautschovica
 Fatsia japonica
 Forsythia intermedia
 F. suspensa
 F. viridissima
 Gaillardia spp.
 Gardenia jasminoides
 Gaura spp.
 Gentiana dahurica
 Geranium cinereum
 Gladiolus spp.
 Gypsophila repens
 Hedera helix
 Hellanthemum spp.
 Hemerocallis spp.

 Hibiscus
 Hibiscus spp.
 Hibiscus rosa-sinensis
 Hosta plantaginea
 H. Sieboldiana
 Houttuynia cordata var. variegata
 Hydrangea macrophylla
 Ilex cornuta
 I. crenata
 I. opaca
 I. pernyi
 I. vomitoria
 Inula ensifolia
 Iris ensata
 Iris siberica
 Iris spp.
 Jasminium nudiflorum
 Juniperus chinensis
 J. conferta
 J. horizontalis
 Juglans spp.
 Justicia brandegeana
 Lagerstromia indica
 Lagerstromia indica X Fauriei
 Lantana montevidensis
 Lavender spp.
 Leontopodium alpinum
 Ligustrum sinense
 Ligustrum amurense
 L. japonicum
 L. lucidum
 Lilium spp.
 Liriope muscari


 Liriope Muscari var. variegata
 Liriope spicata
 Lobelia Cardinalis
 Lonicera japonica
 L. tatarica

Wintercreeper
 Japanese Spindle Tree (Evergreen Euonymus)
 Spreading Euonymus
 Japanese Aralia
 Border Forsythia
 Weeping Forsythia
 Greenstem Forsythia
 Gaillardia, Blanket Flower: "Goblin"
 Gardenia, Cape-Jasmine, Radicans

 Gentian
 Cranesbill
 Gladiolus species**
 Baby's Breath
 English Ivy
 Sunrose
 Daylily: Aztec Gold, Stella de Oro, Tender Love
 Rose of Sharon**
 Mallow: Disco Belle White
 Chinese Hibiscus
 Hosta, Plantain Lily (Fragrant)
 Hosta, "Searsucker"

 Bigleaf Hydrangea
 Chinese Holly
 Japanese Holly: Helliery
 American Holly
 Holly
 Yaupon Holly: Schillings

 Sword-Leaved Iris: Jodlesong
 Siberian Iris: Cabernet
 Iris species**
 Winter Jasmine
 Chinese Juniper: Nick's Compact, Parsonnii
 Shore Juniper: Blue Pacific
 Creeping Juniper
 Walnut*
 Shrimp Plant
 Crape Myrtle
 Crape Myrtle: Tuscarora
 Weeping Lantana
 Lavender: Munstead
 Edelweiss
 Chinese Privet: Variegata
 Amur Privet
 Japanese Privet
 Glossy Privet (Waxleaf)
 Lily: Jazz
 Big Blue Lily: Turf: Silver Mound: Evergreen
 Giant
 Liriope, Variegated
 Liriope, Creeping
 Cardinal Flower, Indian Pink
 Japanese Honeysuckle
 Tatarian Honeysuckle



Santolina virens
Saxifraga spp.
Scabiosa spp.
Sedum album
Sedum cauticola
Sedum dasyphyllum
Sedum spurium
Spiraea Bumalda
Syzygium paniculatum
Taxus cuspidata
T. media
Teucrium spp.
Thalictrum dipterocarpum
Thuja occidentalis
Trachelospermum asiaticum
Tsuga canadensis
Tulipa spp.
Veronica spp.
Viburnum japonicum
V. odoratissimum
V. plicatum
V. rigidum
V. suspensum
V. tinus
V. trilobium
V. wrightii
Vinca major
Vinca minor
Vitis spp.
Weigela florida
Yucca aloifolia
Y. filamentosa

Santolina
 Saxifrage: Purple Dome
 Pincushion Flower
 Stonecrop
 Stonecrop: Lidakense
 Stonecrop
 Stonecrop: Dragon's Blood
 Spirea: Anthony Waterer
 Australian Brushcherry
 Japanese Yew
 Yew
 Germander
 Meadow Rue
 American Arborvitae
 Star Jasmine
 Canada Hemlock
 Tulip species
 Veronica, Speedwell: Sunny Border
 Japanese Viburnum
 Sweet Viburnum
 Japanese Snowball
 Canary Island Viburnum
 Arrowwood Viburnum
 Laurustinus
 Cranberry Bush
 Leatherleaf Viburnum
 Greater Periwinkle, Vinca
 Common Periwinkle, Vinca
 Grape*
 Old Fashioned Weigela
 Spanish Bayonet, Yucca
 Adam's Needle, Yucca

*Do not use on food-producing trees, vines, or plants.

**Not for use on container-grown plants.

WARRANTY DISCLAIMER

Winfield Solutions, LLC warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

The directions for use of this product must be followed carefully. EXCEPT AS PROVIDED ABOVE AND TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF

MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

Barricade® is a registered trademark of Syngenta.
Prolific® is a trademark of Winfield Solutions, LLC.

GUARANTEED FERTILIZER ANALYSIS: (To be attached to final printed labeling)

X-X-X *	
Total Nitrogen (N)X.XX%
Available Phosphate (P ₂ O ₅).....X.XX%
Soluble Potash (K ₂ O).....X.XX%
Chlorine not more than X.X%	
Derived from xxxxxxxxxxxxxxxx.	

* [Editor's Note: All information will vary depending on grade used)

