03/21/2005

1/13

Please read instructions on (everse before completin	g form.			Form App	roved	OMB No.	<u> 2070-0060</u>	O. Approval expires 2-28	-95
\$EPA	Environmental I	ted States Protection ton, DC 2046	•	: y		✓	Registra Amenda Other		OPP Identifier Number	
	Α	pplication	for Pe	sticio	le - Sect	ion	1			
1. Company/Product Number 9779-358	,				roduct Mana Miller	ger		\	posed Classification	
4. Company/Product (Name) Magic Carpet Fertilizer			1 .	M# !3						_
5. Name and Address of App	olicant (Include ZIP Code)	6.	Expe	dited Reve	oiw.	In accorda	nce with	FIFRA Section 3(c)(3)	
Agriliance, LLC, c/o A 481 Country Club Dri Senatobia, MS 38668	ve	lting .	to):	y product is				mposition and labeling	_
Check if this	is a new address		P	Produc	ct Name _					
	· · · · · · · · · · · · · · · · · · ·	-	Sectio	n - II		_				
Amendment - Explain	below.				Final printed			e to	NOTIFICATION	
Resubmission in resp	onse to Agency letter da	sted			"Me Too" A				MAR 2 1 2005	
Notification - Explain	below.				Other - Expl	ain be	low.			
Notification per PR Notice 98 provisions of PR Notice 98-14 formula of this product. I und notification is not consistent venforcement action and penal	0 and EPA regulations at lerstand that it is a violation with the terms of PR Notic	40 CFR 152.46 on of 18 U.S.C. e 98-10 and 40	6, and no oth Sec. 1001 t DCFR 152.4	her cha to willfu	nges have be ily make any	en ma false s	ade to the latestatement to	eling or the EPA. I furth	confidential statement of ner understand that if this	
	· · · · · · · · · · · · · · · · · · ·		Section	n - II			· · · · · · · · · · · · · · · · · · ·			
1. Material This Product Will	Be Packaged In:					_		· <u> </u>	<u></u>	_
Child-Resistant Packaging	Unit Packaging		Water Sol	uble Pa	sckaging		2. Type of	Container		
Yes	Yes		Ye.	8				Metal Plastic		
No	If "Yes"	No. per	If "Yes"	· -	No. per			Glass		
* Certification must be submitted	Unit Packaging wgt.	container	Package v	vgt	container			Other (S	pecify)	
3. Location of Net Contents	Information 4	. Size(s) Retai	l Container			5, Lo	cation of Lal	pel Directio	ns	
Label C	ontainer					<u> </u>	─ On labe	 	<u> </u>	
6. Manner in Which Label is	Affixed to Product	Lithogra Paper gl Stencile	ph ued d		Other			·		
			Section	n - I\	/	•				
1. Contact Point /Complete	items directly below for	identification	of individue	l to be	contacted, i	if nec	essary, to pi	ocess this	application.)	
Name Alice Walker, Ph.D.		1.	itle Agent					Telephone 662-562-5	n No. (Include Area Code 5995)
	ments I have made on th y knowlingliy false or mi aw.		ll attachme						6. Cata Application Recoved	
2. Signature		3.	Title				•	* * * *		
alien We	reku	A	Agent						****	
4. Typed Name Alice Walker, Ph.D.		5.	Date	Ech	ruon, 20	200)E			
AND THIRD I CILD.				. 60	ruary 28,	Z UU]		

ProSource One

Magic Carpet[™] Fertilizer with BARRICADE[®] .25

ACTIVE INGREDIENT:	
Prodiamine, [N ³ , N ³ -Di-n-propyl-2,4-dinitro-6-(trifluoromethyl)-	<u>m</u> -phenylenediamine]0.25%
OTHER INGREDIENTS:	
TOTAL	100.00%

Barricade[®] is a registered trademark of Syngenta. Magic CarpetTM is a trademark of Agriliance, LLC.

For preemergence control of grass and broadleaf weeds in established turf grasses (excluding golf course putting greens) and lawns and landscape ornamentals (including established perennials & wildflower plantings)

KEEP OUT OF REACH OF CHILDREN CAUTION

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Causes moderate eye injury (irritation). Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Prolonged or frequently repeated skin contact while handling the material may cause allergic reactions in some individuals.

EPA Reg. No. 9779-358	EPA Est. No
Distributed By:	NET WEIGHT
Agriliance, LLC St. Paul, MN 55164-0089	0/B28/5

FIRST AID

	Hold eye open and rinse slowly and gently with water for 15-20 minutes.
If in Eyes	• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
	Call a poison control center or doctor for treatment advice.
	Take off contaminated clothing.
If on Skin or	• Rinse skin immediately with plenty of water for 15-20 minutes.
Clothing	Call a poison control center or doctor for treatment advice.
	Move person to fresh air.
If Inhaled	• If person is not breathing, call 911 or an ambulance, then give artificial respiration,
	preferably by mouth-to-mouth, if possible.
	Call a poison control center or doctor for treatment advice.
Have the product c	ontainer or label with you when calling a poison control center or doctor, or going for
treatment. You ma	y also contact 1-877-424-7452 for emergency medical treatment information.

ENVIRONMENTAL HAZARDS

This product has low solubility in water. At the limits of solubility, this product is not toxic to fish. However, at concentrations above the level of water solubility, it may be toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

If material is spilled: Contain and sweep up material of spill for disposal or use per label instructions. Wash cleanup equipment to prevent contamination elsewhere. Excess quantities above normal application rates could result in damage to plants subject to spill. Water-in remainder; disc under, cover with layer of soil, or leave exposed to sunlight, as deemed appropriate to the situation.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Storage: Store in original container away from other fertilizer, feed, or foodstuffs and separated from other pesticides.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal:

Paper Containers: Completely empty container into application equipment. Then dispose of empty container in a sanitary landfill or at an incineration facility, or; if allowed by state and local authorities, by burning locally. If burned, stay out of smoke.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

This product is a selective preemergence herbicide that provides residual control of many grass and broadleaf weeds in:

- Established turf grasses (excluding golf course putting greens) and lawns
- Landscape ornamentals
- Established perennials & wildflower plantings

This product controls susceptible weeds by inhibiting weed seed germination and root development. Most effective weed control will be obtained when this product is activated by at least 0.5 inches of rainfall or irrigation, or with shallow (1-2 inches) incorporation, prior to weed seed germination and within 14 days following application.

Not for use on plants being grown (1) for sale or other commercial use, (2) for commercial seed production, or (3) for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.

Do not graze or feed livestock forage cut from areas treated with this product.

Do not apply this product through any type of irrigation system.

Do not apply aerially.

Do not apply to golf course putting greens.

Failure to follow the directions for use and precautions on this label may result in poor weed control, crop injury, or illegal residues.

APPLICATION DIRECTIONS

Apply uniformly through suitable, calibrated application equipment.


ESTABLISHED TURF

This product is a selective preemergence herbicide that, when properly applied, will control certain grass and broadleaf weeds in established turf grasses and lawns. The maximum amount of this product that may be applied per year is given for each turf grass species in the "Maximum Annual Rates" section of this label.

Most effective weed control in turf grasses will be obtained when this product is activated by at least 0.5 inches of rainfall or irrigation prior to weed seed germination and within 14 days following application. See the map below for approximate crabgrass seed germination dates.



APPROXIMATE CRABGRASS GERMINATION DATES



USE PRECAUTIONS

The following precautions apply to the use of this product in turf grasses and lawns:

- (1) Application of this product may thin emerged annual bluegrass and newly overseeded grasses.
- (2) Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil.
- (3) Do not cut (harvest) treated sod before 120 days after application. Do not apply to newly set sod until the following year.
- (4) Application of this product to turf stressed by drought, low fertility, or pest damage may result in turf injury.
- (5) Disturbing the herbicide barrier with cultural practices such as disking may result in reduced weed control.
- (6) Do not apply this product to putting greens or areas where dichondra, colonial bentgrass, velvet bentgrass, or annual bluegrass (*Poa annua*) are desirable species.
- (7) Keep this product off sidewalks, patios, driveways, etc. This product contains materials which will stain concrete and painted surfaces. If it comes in contact with a concrete surface, sweep off immediately before flushing with water. If concrete is wet when contact occurs, staining may be impossible to prevent. Be careful when using rotary spreaders which scatter product beyond lawn edge and onto concrete.

RATES OF APPLICATION

This product may be applied as a single application or in sequential applications to control weeds germinating throughout the year. All applications must be made prior to germination of the target weeds. This product will not control established weeds.

Maximum use rate selection should be based on turf species. The length of time of residual weed control provided by this product is related to the rate applied.

MAXIMUM ANNUAL RATES

This product is recommended for use on the turf grass species listed in the following table. Do not exceed the maximum yearly rate as given in the following table:

	Maximum Application Rate Per Calendar Year Of Magic Carpet Fertilizer with BARRICADE By Turf Grass Species			
Turf Species	Lbs. Product/A	Lbs. ai/A		
Bermudagrass ²	600 ³	1.5		
Bahiagrass				
Centipedegrass				
Seashore Paspalum				
St. Augustinegrass	İ			
Tall Fescue				
(including turf-type)				
Zoysia				
Buffalograss	400	1.0		
Kentucky Bluegrass				
Perennial Ryegrass				
Creeping Red Fescue	300	0.75		
Creeping Bentgrass	260	0.65		

¹ These are the maximum rates per calendar year by species limitations.

• Use higher rates of this product to achieve higher levels of fertility and longer periods of weed control for each turf type, but do not exceed the "Maximum Annual Rates" table.

WEEDS CONTROLLED

When used in accordance with this label this product will provide control of the following weeds:

Barnyardgrass Carpetweed Bluegrass, Annual (Poa annua) Chickweed, Common Crabgrass (large, smooth) Chickweed, Mouseear (from seed) Crowfootgrass Henbit Cupgrass, Woolly Knotweed Foxtails, Annual Kochia Goosegrass¹ Lambsquarter, Common Itchgrass Pigweed Johnsongrass (from seed) Purslane, Common Junglerice Pusley, Florida

² May be used on newly sprigged or plugged Bermudagrass at rates not to exceed 208 lbs./A. Newly sprigged or plugged Bermudagrass stolon rooting may be temporarily retarded. Suppression only of Foxtail, Goosegrass, and Rescuegrass due to reduced product rates used in sprigging situations.

³ Do not apply more than 600 lbs. of product per calendar year.

Lovegrass
Panicum (Texas, Fall, Browntop)
Rescuegrass³
Signalgrass, Broadleaf
Sprangletop

Shepherd's Purse²
Speedwell, Persian
Spurge, Prostrate
Witchgrass
Woodsorrel, Yellow (from seed)

In many areas a single application of 260 to 600 lbs./A of this product will control goosegrass. However, under heavy goosegrass pressure and/or an extended growing season, most effective weed control may be obtained by making an initial application of 260 to 400 lbs./A followed after 60 to 90 days by a second application at doses that would not exceed those given in the "Maximum Annual Rate" table. Do not exceed the maximum rate for turf grass species listed in the "Maximum Annual Rate" table above.

Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rates recommended for each turf species. All applications must be made prior to germination of the weed seeds.

WHEN TO APPLY AFTER OVERSEEDING TURF

Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil.

WHEN TO OVERSEED AFTER APPLICATION

This product will inhibit the germination of turf species if overseeded too soon after application. Follow rates and intervals in the table below for best overseeding/reseeding results.

Magic Carpet I BARRICADE		Months Before Overseeding			
Product	ai	North	Transition	South	
200	.50	4	4	4	
265	.66	5	4	4	
296	.74	6	5	5	
320	.80		6	6	
400	1.00		7	7	
456	1.14			9	
520	1.30	W-5		10	
600	1.50			12	

LANDSCAPE ORNAMENTALS

(including Established Perennials and Wildflower Plantings)

This product may be applied for residual preemergence weed control in ornamentals.

USE RATES

Apply this product at 265 to 600 lbs./A in fall and/or spring. Use higher rates of application for longer control periods. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rate of 600 lbs./A.

APPLICATION TIMING AND INFORMATION

This product is a preemergence herbicide and will not control emerged weeds. Most effective weed control in ornamentals will be obtained when this product is activated by at least 0.5 inches of rainfall or irrigation, or with

²Applications for this weed should be made in late summer, fall, or winter prior to germination.

³ Suppression only.

shallow (1 to 2 inches) mechanical incorporation, prior to weed seed germination and within 14 days following application.

Best weed control is obtained when this product is applied to soil free of clods, weeds, and debris such as leaves. Prior to application of this product, control existing vegetation by hand weeding, cultivation, or the use of an appropriate postemergence herbicide.

ESTABLISHED LANDSCAPE ORNAMENTALS

This product may be applied to established ornamentals as a broadcast, over the top, or directed application. Irrigation or rainfall soon after application will wash residues off plant foliage and activate this product in the soil.

NEWLY TRANSPLANTED LANDSCAPE ORNAMENTALS

This product may be applied to newly transplanted ornamentals as a broadcast, over the top, or directed application. Delay application to allow soil to settle around new transplants and water thoroughly before applying this product. Apply this product after cuttings form roots and become established. Apply before budding/grafting or after buds/grafts have taken to avoid any inhibition of the tissue union.

TOLERANT ORNAMENTAL SPECIES

This product will not harm most trees, shrubs, vines, and flowers. The species listed below are tolerant to this product. Best results will be obtained when this product is uniformly applied to the soil surface; avoid applying to ornamental shoots which may defeat the effort to get uniform distribution on the soil surface. This product may be applied over the top of these species.

When plants are under stress (such as heat, drought, or frost damage) some cultivars of listed plants may be sensitive to this product.

Scientific name

Abies spp.

Abelia grandiflora

Acer palmatum

A. platanoides

Achillea spp.

Actinidia chinensis

Agapanthus africanus

Agapanthus orientalis

Akebia guintata

Allium cernuum

Anemone hybrida

Aquilegia spp.

Arctostaphylos densiflora

Arctotheca calendula

Artemesia spp.

Aster spp.

Aster X Frikartii

Athrium Filiz-femina

Aucuba japonica

Begonia spp.

Berberis gladwynensis

B. julianae

B. mentorensis

B. thunbergii

B. verruculosa

Bergenia cordifolia

Boltonia asteroides

Bougainvillea spp.

Common name

Fir species** (Balsam, Fraser, Noble, etc.)

Abelia: Sherwood

Japanese Maple

Norway Maple

Yarrow: King Edward

Kiwi*

Lily-of-the-Nile (African Lily)

Five-Leaf or Chocolate Vine

Lady's Leek, Nodding Onion

Japanese Anemone

Aquilegia: Red and Gold

Vine Hill Manzanita

Cape Weed

Wormwood: Silver Mound, Castle

Aster: Bonny Blue, Purple Dome

Lady Fern: Fern Lady

Japanese Aucuba

Fibrous Begonia: Hardy Grandis

Barberry

Wintergreen Barberry

Mentor Barberry

Japanese Barberry

Warty Barberry

Snowbank

Bougainvillea

Buddleia davidii
Buxus microphylla
Callistemon viminalis
Calluna vulgaris
Campanula carpatica
Campis X Tagliabuana

Carpobrotus edulis Cassia artemisoides Ceanothus rigidus

Ceratostigma plumbagonoides

Chamaecyparis pisifera Chrysanthemum nipponicum

Cleyera japonica Citrus spp. Coreopsis spp.

Cornus florida C. stolonifera Cortaderia selloana Cotoneaster apiculatus

C. buxifolius C. dammeri C. microphyllus Cretaegus spp. Crocosmia spp.

Cupressus sempervirens

Delosperma alba
Delosperma spp.
Delphinium spp.
Dianthus deltoides
D. gratianopolitanus
Dodonea viscosa
Echinacea pupurea
Elaeagnus pungens
Euonymus fortunei
E. japonica
E. kiautschovica
Fatsia japonica

Forsythia intermedia F. suspensa F. viridissima Gaillardia spp. Gardenia jasminoides

Gaura spp.

Gentiana dahurica Geranium cinereum Gladiolus spp. Gypsophila repens Hedera helix Hellanthemum spp. Hemerocallis spp.

Hibiscus Hibiscus spp.

Hibiscus rosa-sinensis Hosta plantaginea Butterfly-Bush (Dwarf Blue): Royal Red

Japanese Boxwood Weeping Bottlebrush Scotch Heather

Tussock Bellflower, (White Clips) Trumpet Creeper, Trumpet Flower,

Madame Galen

Hottentot Fig (Ice Plant)

Feathery Cassia Wild Lilac

False Cypress

Cleyera

Citrus species*

Coreopsis (Calliopsis): Early Sunrise,

Moonbeam

Flowering Dogwood American Dogwood Pampas Grass

Cranberry Cotoneaster

Cotoneaster

Bearberry Cotoneaster Rockspray Cotoneaster

Hawthorne Lucifer Italian Cypress

White Trailing Ice Plant

Cooperi Pink Larkspur: Blue Elf

Dianthus: Maiden Pinks "Zing"

Cheddar Pink Hop Bush

Coneflower, Purple; Magnus

Silverberry Wintercreeper

Japanese Spindle Tree (Evergreen Euonymus)

Spreading Euonymus Japanese Aralia Border Forsythia Weeping Forsythia Greenstem Forsythia

Gaillardia, Blanket Flower: "Goblin" Gardenia, Cape-Jasmine, Radicans

Gentian Cranesbill

Gladiolus species**
Baby's Breath
English Ivy
Sunrose

Daylily: Aztec Gold, Stella de Oro, Tender

Love

Rose of Sharon**

Mallow: Disco Belle White

Chinese Hibiscus

Hosta, Plantain Lily (Fragrant)

H. Sieboldiana

Houttuynia cordata var. variegata

Hydrangea macrophylla

Ilex cornuta
I. crenata
I. opaca
I. pernyi
I. vomitoria
Inula ensifolia
Iris ensata
Iris siberica
Iris sop.

Jasminium nudiflorum Juniperus chinensis

J. conferta J. horizontalis Juglans spp.

Justicia brandegeana Lagerstromia indica

Lagerstromia indica X Fauriei

Lantana montevidensis

Lavender spp.

Leontopodium alpinum Ligustrum sinense Ligustrum amurense

L. japonicum L. lucidum Lilium spp. Liriope muscari

Liriope Muscari var. variegata

Liriope spicata Lobelia Cardinalis Lonicera japonica

L. tatarica

Loropetalum chinense

Lythrum spp. Magnolia spp. Maleophora luteola

Malus spp.

Miscanthus sinensis Nandina domestica Narcissus spp. Nerium oleander Olea europaea

Ophiopogon japonicus
Osmanthus heterphyllus
Osteospermum fruticosum
Oxydendrum arboreum
Paeonia suffruticosa
Pennisetum setaceumi
Perovskia atriplicifolia
Persea americana
Photinia fraseri

Picea spp.

Physotegia Viginiana

Hosta, "Searsucker"

Bigleaf Hydrangea Chinese Holly

Japanese Holly: Helleri American Holly

Holly

Yaupon Holly: Schillings

Sword-Leaved Iris: Jodlesong Siberian Iris: Cabernet

Iris species**
Winter Jasmine

Chinese Juniper: Nick's Compact, Parsonnii

Shore Juniper: Blue Pacific

Creeping Juniper Walnut* Shrimp Plant Crape Myrtle

Crape Myrtle: Tuscarora Weeping Lantana Lavender: Munstead

Edelweiss

Chinese Privet: Variegata

Amur Privet
Japanese Privet

Glossy Privet (Waxleaf)

Lily, Jazz

Big Blue Lily; Turf: Silver Mound: Evergreen

Giant

Liriope, Variegated Liriope, Creeping

Cardinal Flower, Indian Pink Japanese Honeysuckle Tatarian Honeysuckle

Burgundy

Loosestrife; Modern Pink Magnolia species**

Ice Plant
Crabapple*

Yaku Jima, Silberfeder Heavenly Bamboo Narcissus species**

Oleander Olive*

Mondo Grass

Osmanthus (False Holly): Gulf Tide

Trailing African Daisy

Sourwood Tree Peony

Fountain Grass (Dwarf)

Avocado*

Frasier's Photinia (Redtip) Dragonhead, False: Vivid

Spruce species**(Colorado Blue, Norway,

etc.)

Pieris japonica
Pinus brutia
P. canariensis
P. elliottii
P. halepensis
P. nigra
P. palustrus
P. radiata
P. thunbergiana
P. strobus
P. sylvestris
P. taeda
P. virginiana
Pistachio spp.

Pittosporum rhombifolium

P. tobira

Podocarpus macrophyllus Prunus laurocerasus

Prunus spp.

Pseudotsuga menziesii Pyracantha coccinea P. fortuneana

P. koidzumii Pyrus spp.

Quercus Shumardii Quercus spp. Raphiolepis indica Raphiolepsis umbellata

Rhododendron
(including Azalea)

Rosmarinus officinalis

Rudbeckia spp.
Santolina virens
Saxifraga spp.
Scabiosa spp.
Sedum album
Sedum cauticola
Sedum dasyphyllum
Sedum spurium
Spiraea Bumalda
Syzygium paniculatum
Taxus cuspidata

T. media Teucrium spp.

V. rigidum

Thalictrum dipterocarpum

Thuja occidentalis

Trachelospermum asiatum

Trucheiospermum asiai
Tsuga canadensis
Tulipa spp.
Veroniac spp.
Viburnum japonicum
V. odoratissimum
V. plicatum

Lily-of-the-Valley Bush

Calabrian Pine
Canary Island Pine
Slash Pine
Aleppo Pine
Austrian Pine
Longleaf Pine
Monterey Pine

Japanese Black Pine
Eastern White Pine
Scotch Pine
Loblolly Pine
Virginia Pine
Pistachio*

Queensland Pittosporum Japanese Pittosporum

Japanese Yew English Laurel

Almond, Apricot, Nectarine, Peach, Plum and Prune*

Douglas Fir**
Firethorn (Scarlet)
Firethorn

Firethorn Bradford Pear sp. Oak, Shumard's Red

Oak species Indian Hawthorne Yedda Hawthorn

Coral Bells, Delaware Valley White, Flame Creeper, Formosa, George L. Tabor, Girard Crimson, Hino-crimson, PJM, Roseum Elegans, Wakeiebisu, White Gumpo

Rosemary*

Black-Eyed Susan; Goldstrum

Santolina

Saxifrage: Purple Dome Pincushion Flower

Stonecrop

Stonecrop: Lidakense

Stonecrop

Stonecrop: Dragon's Blood Spirea: Anthony Waterer Australian Brushcherry

Japanese Yew

Yew Germander Meadow Rue

American Arborvitae Star Jasmine

Canada Hemlock Tulip species

Veronica, Speedwell: Sunny Border

Japanese Viburnum Sweet Viburnum Japanese Snowball Canary Island Viburnum V. suspensum
V. tinus
V. trilobium
V. wrightii
Vinca major
Vinca minor
Vitis spp.
Weigela florida
Yucca aloifolia
Y. filamentosa

Arrowood Viburnum
Laurustinus
Cranberry Bush
Leatherleaf Viburnum
Greater Periwinkle, Vinca
Common Periwinkle, Vinca
Grape*
Old Fashioned Weigela
Spanish Bayonet, Yucca
Adam's Needle, Yucca

Notice: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to Seller, and Buyer assumes the risk of any such use.

^{*}Do not use on food-producing trees, vines, or plants.

^{**}Not for use on container-grown plants.

CERTIFIED MAIL

February 28, 2005

Document Processing Desk (Notif) Office of Pesticide Programs (7504C) U.S. Environmental Protection Agency 1200 Pennsylvania Ave., NW Washington, DC 20460

Re: Magic Carpet Fertilizer with Barricade .25, EPA Reg. No. 9779-358

Notification under PRN 98-10 of Addition of Language Regarding Possible Staining of

Concrete

Attn: Sherada Hobgood

This correspondence will constitute notification from AGRILIANCE, LLC to add label language recommending that the product be kept off concrete to avoid staining. We have highlighted the new text on page 4 for your convenience. Following a telecon with Jim Stone on this date, we are submitting this label revision as a notification rather than an amendment.

I believe this notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under section 12 and 14 of FIFRA.

Thank you for adding this notification to the registration records for this product. We would appreciate a stamped "Notification" copy for our files.

Sincerely,

Alice Walker, Ph.D.

Regulatory Agent for

Walker

AGRILIANCE, LLC

Enclosures

cc: Mr. Gary Halvorson