



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

September 13, 2019

Diana Williams
Registration Specialist
Winfield Solutions LLC
P.O. Box 64589
St. Paul, MN 55164

Subject: Notification per PRN 98-10 – Add alternate brand names, optional claims, referral statement, and reformatting
Product Name: MAGIC CARPET FERTILIZER WITH BARRICADE .28%
EPA Registration Number: 9779-363
Application Date: April 26, 2019
Decision Number: 551484

Dear Ms. Williams:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 for the above referenced product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10.

The alternate brand names **“GROUNDWORK CRABGRASS PREVENTER FERTILIZER WITH .28% BARRICADE”** and **“GROUNDWORK CRABGRASS PREVENTER FERTILIZER WITH BARRICADE .28%”** have been added to the product record.

The label submitted with the application has been stamped “Notification” and will be placed in our records.

Should you wish to add/retain a reference to the company’s website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA’s Office of Enforcement and Compliance.

If you have any questions, please contact Curtis Hildebrandt at 703-347-8198 or by email at hildebrandt.curtis@epa.gov.

Sincerely,

A handwritten signature in cursive script that reads "Mindy Ondish".

Mindy Ondish
Product Manager 23
Herbicide Branch
Registration Division (7505P)
Office of Pesticide Programs

NOTIFICATION

9779-363

The applicant has certified that no changes, other than those reported to the Agency have been made to the labeling. The Agency acknowledges this notification by letter dated:

09/13/2019

**MAGIC CARPET™ FERTILIZER
WITH BARRICADE® .28%**

- [Alternate Brand Names: BRAZOS BEST FOUNDATION PLUS 0.28% BARRICADE
- BRAZOS BEST WINTERIZER PLUS 0.28% BARRICADE®
- FAITHFUL CRABGRASS PREVENTER FERTILIZER WITH BARRICADE® 0.28%
- FERTILIZER WITH BARRICADE® 0.28%
- GROUNDWORK CRABGRASS PRE-EMERGENT FERTILIZER
- GROUNDWORK CRABGRASS PREVENTER FERTILIZER WITH BARRICADE® 0.28%
- GROUNDWORK CRABGRASS PREVENTER FERTILIZER WITH .28% BARRICADE®
- GROUNDWORK CRABGRASS PREVENTER FERTILIZER WITH BARRICADE® .28%
- PROLIFIC FERTILIZER WITH BARRICADE® 0.28%
- PROSOURCE ONE PROFESSIONAL TURF SOLUTIONS WITH BARRICADE® 0.28%]

[Controls Crabgrass and Other Weeds Early Spring]

ACTIVE INGREDIENT:

Prodiamine, [N³, N³-Di-n-propyl-2,4-dinitro-6-(trifluoromethyl)-m-phenylenediamine] 0.28%

OTHER INGREDIENTS: 99.72%

TOTAL 100.00%

[Barricade® is a registered trademark of Syngenta.]

[Magic Carpet™ is a trademark of Winfield Solutions, LLC.]

[For selective preemergence control of grass and broadleaf weeds in established turf grasses (excluding golf courseputting greens) and lawns and landscape ornamentals (including established perennials & wildflower plantings)]

[PHOSPHOROUS FREE]

[FORMULATED FOR FIRST APPLICATION OF SEASON FOR CRABGRASS CONTROL]

[Control grass & broadleaf weeds in established lawns, landscape ornamentals (including established perennials and wildflower Plantings) and turf grass (excluding golf course putting greens)]

[Controls Crabgrass and Tough Broadleaf Weeds]

[Apply before crabgrass germinates]

[Pre-Emergent for Crabgrass]

[COVERS UP TO 12,500 SQUARE FEET]

KEEP OUT OF REACH OF CHILDREN

CAUTION

[See Back Panel for Additional Precautionary Statements]

FIRST AID

| | |
|----------------------|---|
| If in Eyes | <ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice. |
| If on Skin or | <ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. |

| | |
|---|---|
| Clothing | • Call a poison control center or doctor for treatment advice. |
| If Inhaled | • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for treatment advice. |
| Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-424-7452 for emergency medical treatment information. | |

EPA Reg. No. 9779-363
Distributed by:
Winfield Solutions, LLC
P.O. Box 64589, St. Paul, MN 55164-0589

EPA Est. No. _____
Net Weight: _____

~~1/0825/6~~ 1/0426/9

PRECAUTIONARY STATEMENTS [moved here from above First Aid]
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes moderate eye injury (irritation). Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Prolonged or frequently repeated skin contact while handling the material may cause allergic reactions in some individuals.

ENVIRONMENTAL HAZARDS

This product has low solubility in water. At the limits of solubility, this product is not toxic to fish. However, at concentrations above the level of water solubility, it may be toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

This product is a selective preemergence herbicide that provides residual control of many grass and broadleaf weeds in:

- Established turf grasses (excluding golf course putting greens) and lawns
- Landscape ornamentals
- Established perennials & wildflower plantings

This product controls susceptible weeds by inhibiting weed seed germination and root development. Most effective weed control will be obtained when this product is activated by at least 0.5 inches of rainfall or irrigation, or with shallow (1-2 inches) incorporation, prior to weed seed germination and within 14 days following application.

Not for use on plants being grown (1) for sale or other commercial use, (2) for commercial seed production, or (3) for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.

Do not graze or feed livestock forage cut from areas treated with this product.

Do not apply this product through any type of irrigation system.

Do not apply aerially.

Do not apply to golf course putting greens.

Failure to follow the directions for use and precautions on this label may result in poor weed control, crop injury, or illegal residues.

APPLICATION DIRECTIONS


Apply uniformly through suitable, calibrated application equipment.

ESTABLISHED TURF

This product is a selective preemergence herbicide that, when properly applied, will control certain grass and broadleaf weeds in established turf grasses and lawns. The maximum amount of this product that may be applied per year is given for each turf grass species in the "Maximum Annual Rates" section of this label.

Most effective weed control in turf grasses will be obtained when this product is activated by at least 0.5 inches of rainfall or irrigation prior to weed seed germination and within 14 days following application. See the map below for approximate crabgrass seed germination dates.

APPROXIMATE CRABGRASS SEED GERMINATION DATES



USE PRECAUTIONS

The following precautions apply to the use of this product in turf grasses and lawns:

(1) Application of this product may thin emerged annual bluegrass and newly overseeded grasses.

- (2) Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil.
- (3) Do not cut (harvest) treated sod before 120 days after application. Do not apply to newly set sod until the following year.
- (4) Application of this product to turf stressed by drought, low fertility, or pest damage may result in turf injury.
- (5) Disturbing the herbicide barrier with cultural practices such as disking may result in reduced weed control.
- (6) Do not apply this product to putting greens or areas where dichondra, colonial bentgrass, velvet bentgrass, or annual bluegrass (*Poa annua*) are desirable species.
- (7) Keep this product off sidewalks, patios, driveways, etc. This product contains materials which will stain concrete and painted surfaces. If it comes in contact with a concrete surface, sweep off immediately before flushing with water. If concrete is wet when contact occurs, staining may be impossible to prevent. Be careful when using rotary spreaders which scatter product beyond lawn edge and onto concrete.

RATES OF APPLICATION

This product may be applied as a single application or in sequential applications to control weeds germinating throughout the year. All applications must be made prior to germination of the target weeds. This product will not control established weeds.

Maximum use rate selection should be based on turf species. The length of time of residual weed control provided by this product is related to the rate applied.

MAXIMUM ANNUAL RATES

This product is recommended for use on the turf grass species listed in the following table. Do not exceed the maximum yearly rate as given in the following table:

| Maximum Application Rate Per Calendar Year By Turf Grass Species ¹ | | |
|---|----------------|-----------|
| Turf Species | Lbs. Product/A | Lbs. ai/A |
| Bermudagrass ² Bahagrass Centipedegrass Seashore Paspalum St. Augustinegrass Tall Fescue (including turf-type) Zoysia | 530 | 1.5 |
| Buffalograss Kentucky Bluegrass Perennial Ryegrass | 353 | 1.0 |
| Creeping Red Fescue | 265 | 0.75 |
| Creeping Bentgrass | 230 | 0.65 |

¹ These are the maximum rates per calendar year by species limitations.

² May be used on newly sprigged or plugged Bermudagrass at rates not to exceed 183 lbs./A or 4 lbs./1000 sq. ft. Newly sprigged or plugged Bermudagrass stolon rooting may be temporarily

retarded. Suppression only of Foxtail, Goosegrass and Rescuegrass due to reduced product rates used in sprigging situations.

Do not apply more than 530 lbs. of product per calendar year.

Use higher rates of this product to achieve higher levels of fertility and longer periods of weed control for each turf type, but do not exceed the "Maximum Annual Rates" table.

WEEDS CONTROLLED

When used in accordance with this label this product will provide control of the following weeds:

| | |
|--|---------------------------------|
| Barnyardgrass | Carpetweed |
| Bluegrass, Annual (<i>Poa annua</i>) | Chickweed, Common |
| Crabgrass (large, smooth) | Chickweed, Mouseear (from seed) |
| Crowfootgrass | Henbit |
| Cupgrass, Woolly | Knotweed |
| Foxtails, Annual | Kochia |
| Goosegrass ¹ | Lambsquarter, Common |
| Itchgrass | Pigweed |
| Johnsongrass (from seed) | Purslane, Common |
| Junglerice | Pusley, Florida |
| Lovegrass | Shepherdspurse ² |
| Panicum (Texas, Fall, Browntop) | Speedwell, Persian |
| Rescuegrass ³ | Spurge, Prostrate |
| Signalgrass, Broadleaf | Witchgrass |
| Sprangletop | Woodsorrel, Yellow (from seed) |

¹ In many areas a single application of 230 to 530 lbs./A of this product will control goosegrass. However, under heavy goosegrass pressure and/or an extended growing season, most effective weed control may be obtained by making an initial application of 230 to 353 lbs./A followed after 60 to 90 days by a second application at doses that would not exceed those given in the "Maximum Annual Rates" table. Do not exceed the maximum rate for turf grass species listed in the "Maximum Annual Rates" table above.

² Applications for this weed should be made in late summer, fall, or winter prior to germination.

³ Suppression only.

Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rates recommended for each turf species. All applications must be made prior to germination of the weed seeds.

WHEN TO APPLY AFTER OVERSEEDING TURF

Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil.

WHEN TO OVERSEED AFTER APPLICATION

This product will inhibit the germination of turf species if overseeded too soon after application. Follow rates and intervals in the table below for best overseeding/reseeding results.

| Magic Carpet Fertilizer with BARRICADE Rate (lb./A) | | Months Before Overseeding | | |
|--|-----------|----------------------------------|-------------------|--------------|
| Product | ai | North | Transition | South |
| 178 | .50 | 4 | 4 | 4 |
| 235 | .66 | 5 | 4 | 4 |
| 264 | .74 | 6 | 5 | 5 |
| 285 | .80 | --- | 6 | 6 |
| 357 | 1.00 | --- | 7 | 7 |
| 407 | 1.14 | --- | --- | 9 |
| 464 | 1.30 | --- | --- | 10 |
| 530 | 1.50 | --- | --- | 12 |

**LANDSCAPE ORNAMENTALS
(including Established Perennials and Wildflower Plantings)**

This product may be applied for residual preemergence weed control in ornamentals.

USE RATES

Apply this product at 235 to 530 lbs./A in fall and/or spring. Use higher rates of application for longer control periods. Sequential or single applications are allowed so long as the total amount of product applied does not exceed the maximum annual application rate of 530 lbs./A.

APPLICATION TIMING AND INFORMATION

This product is a preemergence herbicide and will not control emerged weeds. Most effective weed control in ornamentals will be obtained when this product is activated by at least 0.5 inches of rainfall or irrigation, or with shallow (1 to 2 inches) mechanical incorporation, prior to weed seed germination and within 14 days following application.

Best weed control is obtained when this product is applied to soil free of clods, weeds, and debris such as leaves. Prior to application of this product, control existing vegetation by hand weeding, cultivation or the use of an appropriate postemergence herbicide.

ESTABLISHED LANDSCAPE ORNAMENTALS

This product may be applied to established ornamentals as a broadcast, over the top, or directed application. Irrigation or rainfall soon after application will wash residues off plant foliage and activate this product in the soil.

NEWLY TRANSPLANTED LANDSCAPE ORNAMENTALS

This product may be applied to newly transplanted ornamentals as a broadcast, over the top, or directed application. Delay application to allow soil to settle around new transplants and water thoroughly before applying this product. Apply this product after cuttings form roots and become established. Apply before budding/grafting or after buds/grafts have taken to avoid any inhibition of the tissue union.

TOLERANT ORNAMENTAL SPECIES

This product will not harm most trees, shrubs, vines, and flowers. The species listed below are tolerant to this product. Best results will be obtained when this product is uniformly applied to the soil surface; avoid applying to ornamental shoots which may defeat the effort to get uniform distribution on the soil surface. This product may be applied over the top of these species.

When plants are under stress (such as heat, drought, or frost damage), some cultivars of listed plants may be sensitive to this product.

Table 1. Tolerant Ornamental Species – All States

| Scientific name | Common name |
|----------------------------------|--|
| <i>Abies</i> spp.** | Fir species** (Balsam, Fraser, Noble, etc.) |
| <i>Acer palmatum</i> | Japanese Maple |
| <i>A. platanoides</i> | Norway Maple |
| <i>Actinidia chinensis</i> * | Kiwi* |
| <i>Agapanthus africanus</i> | Lily-of-the-Nile (African Lily) |
| <i>Arctostaphylos densiflora</i> | Vine Hill Manzanita |
| <i>Arctotheca calendula</i> | Cape Weed |
| <i>Aucuba japonica</i> | Japanese Aucuba |
| <i>Berberis gladwynensis</i> | Barberry |
| <i>B. julianae</i> | Wintergreen Barberry |
| <i>B. mentorensis</i> | Mentor Barberry |
| <i>B. thunbergii</i> | Japanese Barberry |
| <i>B. verruculosa</i> | Warty Barberry |
| <i>Buxus microphylla</i> | Japanese Boxwood |
| <i>Callistemon viminalis</i> | Weeping Bottlebrush |
| <i>Calluna vulgaris</i> | Scotch Heather |
| <i>Carpobrotus edulis</i> | Hottentot Fig (Ice Plant) |
| <i>Cassia artemisoides</i> | Feathery Cassia |
| <i>Ceanothus rigidus</i> | Wild Lilac |
| <i>Chamaecyparis pisifera</i> | False Cypress |
| <i>Cleyera japonica</i> | Cleyera |
| <i>Citrus</i> spp.* | Citrus species* |
| <i>Cornus florida</i> | Flowering Dogwood |
| <i>C. stolonifera</i> | American Dogwood |
| <i>Cortaderia selloana</i> | Pampas Grass |
| <i>Cotoneaster apiculatus</i> | Cranberry Cotoneaster |
| <i>C. buxifolius</i> | Cotoneaster |
| <i>C. dammeri</i> | Bearberry Cotoneaster |
| <i>C. microphyllus</i> | Rockspray Cotoneaster |
| <i>Cretaeus</i> spp. | Hawthorne |
| <i>Cupressus sempervirens</i> | Italian Cypress |
| <i>Delosperma alba</i> | White Trailing Ice Plant |
| <i>Dodonea viscosa</i> | Hop Bush |
| <i>Elaeagnus pungens</i> | Silverberry |
| <i>Euonymus fortunei</i> | Wintercreeper |
| <i>E. japonica</i> | Japanese Spindle Tree (Evergreen Euonymus) |
| <i>E. kiautschovica</i> | Spreading Euonymus |
| <i>Fatsia japonica</i> | Japanese Aralia |
| <i>Forsythia intermedia</i> | Border Forsythia |
| <i>F. viridissima</i> | Greenstem Forsythia |
| <i>Gaillardia</i> spp. | Gaillardia, Blanket Flower: “Goblin” |
| <i>Gardenia jasminoides</i> | Gardenia, Cape-Jasmine |
| <i>Gladiolus</i> spp.** | Gladiolus species** |
| <i>Hedera helix</i> | English Ivy |
| <i>Hibiscus</i> ** | Rose of Sharon** |
| <i>Hibiscus rosa-sinensis</i> | Chinese Hibiscus |
| <i>Ilex cornuta</i> | Chinese Holly |
| <i>I. crenata</i> | Japanese Holly |
| <i>I. opaca</i> | American Holly |
| <i>I. pernyi</i> | Holly |
| <i>I. vomitoria</i> | Yaupon Holly |
| <i>Inula ensifolia</i> | |
| <i>Iris</i> spp.** | Iris species** |
| <i>Jasminium nudiflorum</i> | Winter Jasmine |

| | |
|--|---|
| <i>Juniperus chinensis</i> | Chinese Juniper |
| <i>J. conferta</i> | Shore Juniper: Blue Pacific |
| <i>J. horizontalis</i> | Creeping Juniper |
| <i>Juglans</i> spp.* | Walnut* |
| <i>Justicia brandegeana</i> | Shrimp Plant |
| <i>Lagerstromia indica</i> | Crape Myrtle |
| <i>Ligustrum amurense</i> | Amur Privet |
| <i>L. japonicum</i> | Japanese Privet |
| <i>L. lucidum</i> | Glossy Privet (Waxleaf) |
| <i>Liriope muscari</i> | Big Blue Lilyturf |
| <i>Lonicera japonica</i> | Japanese Honeysuckle |
| <i>L. tatarica</i> | Tatarian Honeysuckle |
| <i>Magnolia</i> spp.** | Magnolia species** |
| <i>Maleophora luteola</i> | Ice Plant |
| <i>Malus</i> spp.* | Crabapple* |
| <i>Nandina domestica</i> | Heavenly Bamboo |
| <i>Narcissus</i> spp.** | Narcissus species** |
| <i>Nerium oleander</i> | Oleander |
| <i>Olea europaea</i> * | Olive* |
| <i>Ophiopogon japonicus</i> | Mondo Grass |
| <i>Osteospermum fruticosum</i> | Trailing African Daisy |
| <i>Oxydendrum arboreum</i> | Sourwood |
| <i>Persea americana</i> * | Avocado* |
| <i>Photinia fraseri</i> | Frasier's Photinia (Redtip) |
| <i>Picea</i> spp.** | Spruce species** (Colorado Blue, Norway, etc.) |
| <i>Pieris japonica</i> | Lily-of-the-Valley Bush |
| <i>Pinus brutia</i> | Calabrian Pine |
| <i>P. canariensis</i> | Canary Island Pine |
| <i>P. elliotii</i> | Slash Pine |
| <i>P. halepensis</i> | Aleppo Pine |
| <i>P. nigra</i> | Austrian Black Pine |
| <i>P. palustris</i> | Longleaf Pine |
| <i>P. radiata</i> | Monterey Pine |
| <i>P. strobus</i> | Eastern White Pine |
| <i>P. sylvestris</i> | Scotch Pine |
| <i>P. taeda</i> | Loblolly Pine |
| <i>P. virginiana</i> | Virginia Pine |
| <i>Pistachio</i> spp.* | Pistachio* |
| <i>Pittosporum rhombifolium</i> | Queensland Pittosporum |
| <i>P. tobira</i> | Japanese Pittosporum |
| <i>Podocarpus macrophyllus</i> | Japanese Yew |
| <i>Prunus laurocerasus</i> | English Laurel |
| <i>Prunus</i> spp.* | Almond, Apricot, Nectarine, Peach, Plum and Prune* |
| <i>Pseudotsuga menziesii</i> ** | Douglas Fir** |
| <i>Pyracantha coccinea</i> | Firethorn (Scarlet) |
| <i>P. fortuneana</i> | Firethorn |
| <i>P. koidzumii</i> | Firethorn |
| <i>Pyrus</i> spp. | Bradford Pear sp. |
| <i>Quercus</i> spp. | Oak species |
| <i>Raphiolepis indica</i> | Indian Hawthorne |
| <i>Rhododendron</i> (including Azalea) | 'Coral Bells', 'Formosa', 'PJM', 'Roseum Elegans' |
| <i>Rosa banksiana</i> | Lady Bank's Rose |
| <i>Rosmarinus officinalis</i> * | Rosemary* |
| <i>Santolina virens</i> | Santolina |
| <i>Sedum album</i> | Stonecrop |
| <i>Syzygium paniculatum</i> | Japanese Boxcherry |

| | |
|----------------------------------|------------------------|
| <i>Taxus cuspidata</i> | Japanese Yew |
| <i>T. media</i> | Yew |
| <i>Thuja occidentalis</i> | American Arborvitae |
| <i>Trachelospermum asiaticum</i> | Star Jasmine |
| <i>Tsuga canadensis</i> | Canada Hemlock |
| <i>Tulipa</i> spp. | Tulip species |
| <i>Viburnum japonicum</i> | Japanese Viburnum |
| <i>V. odoratissimum</i> | Sweet Viburnum |
| <i>V. plicatum</i> | Japanese Snowball |
| <i>V. rigidum</i> | Canary Island Viburnum |
| <i>V. tinus</i> | Laurustinus |
| <i>V. trilobium</i> | Cranberry Bush |
| <i>V. wrightii</i> | Leatherleaf Viburnum |
| <i>Vinca major</i> | Vinca |
| <i>Vinca minor</i> | Dwarf Periwinkle |
| <i>Vitis</i> spp.* | Grape* |
| <i>Weigela florida</i> | Old Fashioned Weigela |
| <i>Yucca aloifolia</i> | Spanish Bayonet, Yucca |
| <i>Y. filamentosa</i> | Adam's Needle, Yucca |

*Do not use on food-producing trees, vines, or plants.

**Not for use on container-grown plants.

Table 2. Tolerant Ornamental Species/Varieties – All States Except CA

| Scientific Name | Common name |
|------------------------------------|--|
| <i>Abella grandiflora</i> | Abelia: Sherwood |
| <i>Achilles</i> spp. | Yarrow: King Edward |
| <i>Agapanthus orientalis</i> | |
| <i>Akebia quintara</i> | Five-Leaf or Chocolate Vine |
| <i>Allium cernuum</i> | Lady's Leek, Nodding Onion |
| <i>Anemone hybrida</i> | Japanese Anemone |
| <i>Aquilegia</i> spp. | Aquilegia: Red and Gold |
| <i>Artemisia</i> spp. | Wormwood: Silver Mound, Castle |
| <i>Aster</i> spp. | Aster: Bonny Blue, Purple Dome |
| <i>Aster X frikartii</i> | |
| <i>Athyrium filiz-femina</i> | Lady Fern: Fern Lady |
| <i>Begonia</i> spp. | Fibrous Begonia: Hardy Grandis |
| <i>Bergenia cordifolia</i> | |
| <i>Boltonia asteroides</i> | Snowbank |
| <i>Bougainvillea</i> spp. | Bougainvillea |
| <i>Buddleia davidii</i> | Butterfly-Bush (Dwarf Blue): Royal Red |
| <i>Caillstemon citrinus</i> | Crimson Bottlebrush |
| <i>Campanula carpatica</i> | Tussock Bellflower, (White Clips) |
| <i>Campis X Tagliabuana</i> | Trumpet Creeper, Trumpet Flower, Madame Galen |
| <i>Ceratostigma plumbagonoides</i> | |
| <i>Chrysanthemum nipponicum</i> | |
| <i>Coreopsis</i> spp. | Coreopsis (Calliopisi): Early Sunrise, Moonbeam |
| <i>Crocsmia</i> spp. | Lucifer |
| <i>Delosperma</i> spp. | Cooperi Pink |
| <i>Delphinium</i> spp. | Larkspur: Blue Elf |
| <i>Dianthus deltoides</i> | Dianthus: Maiden Pinks "Zing" |
| <i>D. gratuabioikutabys</i> | Cheddar Pink |
| <i>Echinacea pupurea</i> | Coneflower, Purple: Magnus |
| <i>Forsythia suspense</i> | Weeping Forsythia |
| <i>Gaillardia</i> spp. | Gaillardia, Blanket Flower: "Goblin" |

| | |
|---|---|
| <i>Gaura</i> spp. | Gentian |
| <i>Gentiana dahurica</i> | Cranesbill |
| <i>Geranium cinereum</i> | Baby's Breath |
| <i>Gypsophila repens</i> | Sunrose |
| <i>Hellanthemum</i> spp. | Daylily: Aztec Gold, Stella De Oro, |
| <i>Hemerocallis</i> spp. | Tender Love |
| <i>Heucherella</i> spp. | Caral Bell: Bridget Bloom |
| <i>Hibiscus</i> spp. | Mallow: Disco Belle White |
| <i>Hosta plantaginea</i> | Hosta, Plantain Lily (Fragrant) |
| <i>H. Sieboldiana</i> | Hasta, "Searsucker" |
| <i>Houttuynia cordata</i> var. <i>variegata</i> | |
| <i>Hydrangea macrophylla</i> | Bigleaf Hydrangea |
| <i>Inula ensifolia</i> | |
| <i>Iris ensata</i> | Sword-Leaved Iris: Jodlesong |
| <i>Iris siberica</i> | Siberian Iris: Cabernet |
| <i>Juniperus davurica</i> | Parsoni |
| <i>Lagerstromia indica</i> X <i>Fauriei</i> | Crape Myrtle: Tuscarora |
| <i>Lantana montevidensis</i> | Weeping Lantana |
| <i>Lavender</i> spp. | Lavender: Munstead |
| <i>Leontopodium alpinum</i> | Edelweiss |
| <i>Ligustrum sinense</i> | Chinese Privet: Variegata |
| <i>Lilium</i> spp. | Lil: Jazz |
| <i>Liriope spicata</i> | Liriope, Creeping |
| <i>Lobelia Cardinalis</i> | Cardinal Flower, Indian Pink |
| <i>Loropetalum chineense</i> | Burgundy |
| <i>Lythrum</i> spp. | Loosestrife: Modern Pink |
| <i>Miscanthus sinensis</i> | Yaku Jima, Silberfeder |
| <i>Oenothera missouriensis</i> | Evening Primrose |
| <i>Osmanthus heterophyllus</i> | Osmanthus (False Holly): Gulf Tide |
| <i>Paeonia suffruticosa</i> | Tree Peony |
| <i>Pennisetum setaceum</i> | Fountain Grass (Dwarf) |
| <i>Perovskia atriplicifolia</i> | |
| <i>Physotegia virginiana</i> | Dragonhead, False: Vivid |
| <i>Quercus Shumardii</i> | Oak, Shumard's Red |
| <i>Raphiolepis umbellata</i> | Yedda Hawthorn |
| <i>Rhododendron</i> (including Azalea) | 'Delaware Valley White', 'Flame Creeper', 'Girard Crimson', George L. Tabor', 'Wakeiebisu', 'White Gumpo' |
| <i>Rudbeckia</i> spp. | Black-Eyed Susan: Goldstrum |
| <i>Saxifraga</i> spp. | Saxifrage: Purple Dome |
| <i>Scabiosa</i> spp. | Pincushion Flower |
| <i>Sedum caudicola</i> | Stonecrop: Lidakense |
| <i>Sedum dasyphyllum</i> | Stonecrop |
| <i>Sedum spurium</i> | Stonecrop: Dragon's Blood |
| <i>Spiraea Bumalda</i> | Spirea: Anthony Waterer |
| <i>Syzygium paniculatum</i> | Australian Brushcherry |
| <i>Teucrium</i> spp. | Germander |
| <i>Thalictrum dipterocarpum</i> | Meadow Rue |
| <i>Veronica</i> spp. | Veronica, Speedwell: Sunny Border |
| <i>Viburnum suspensum</i> | Arrowood Viburnum |

STORAGE AND DISPOSAL [Moved here from above Directions for Use]

Do not contaminate water, food, or feed by storage or disposal.

Storage: Store in original container away from other fertilizer, feed, or food stuffs and separated from other pesticides.

If material is spilled: Contain and sweep up material of spill for disposal or use per label instructions. Wash cleanup equipment to prevent contamination elsewhere. Excess quantities above normal application rates could result in damage to plants subject to spill. Water-in remainder, disc under, cover with layer of soil, or leave exposed to sunlight, as deemed appropriate to the situation.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal:

Paper Containers: Nonrefillable Container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available or dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

| Guaranteed Fertilizer Analysis | |
|---|-----|
| [will vary depending on grade used] | |
| Total Nitrogen (N) | 00% |
| 0.0% Ammoniacal nitrogen | |
| 0.0% Urea nitrogen | |
| Available Phosphate (P ₂ O ₅) | 00% |
| Soluble Potash (K ₂ O) | 00% |
| Derived from urea, monoammonium phosphate and potassium chloride. | |

WARRANTY DISCLAIMER

Winfield Solutions, LLC warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

The directions for use of this product must be followed carefully. EXCEPT AS PROVIDED ABOVE AND TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.