

DIRECTIONS FOR USE

Pour recommended amount of this material on surface of water in nearly filled spray tank. Add balance of water to fill tank. Keep agitator running during filling, mixing, and spraying operations. Do not allow mixture to stand. Do not combine wettable powders with emulsifiable liquids in the same spray tank talks previous use of the materials being combined has proved them to be physically compatible.

Dosages are given in terms of pints per acre. Use sufficient water for thorough coverage.

BEANS: Bean Leafroller, Serpentine Leafminer, Apply at the rate of 2 pints per acre. Make 2 applications at 15 day intervals. Make no application to green or snap beans after pods start to form. DO NOT FEED TREATED FOF.AGE TO DAIRY ANIMALS OR TO ANIMALS BEING FINISHED FOR SLAUGHTER.

COLE CROPS: (Broccoli, Brussels Sprouts, Cabbage, Cauliflower): Armyworms, Imported Cabbage Worm use 2 pints per acre. Cabbage Looper, Climbing Cutworms, apply 1 - 1/2 pints per acre at 7 - 10 day intervals. DO NOT APPLY AFTER EDIBLE PARTS BEGIN TO FORM.

TOMATOES: Cabbage Looper, Climbing Cutworms, Flea Beetle, Hornworm, Tomatoe Fruitworm. Apply at the rate of 1½-4 pints per acre. DO NOT APPLY WITHIN 3 DAYS OF HARVEST.

COWPEAS: Cowpea Curculio – Use 1½ - 2 pints per acre. Do not apply to cowpeas to be used as green snaps after pods begin to form. DO NOT FEED TREATED FORAGE TO DAIRY ANIMALS OR TO ANIMALS BEING FINISHED FOR SLAUGHTER.

- CHILL	• WOODBURY
P	
AR 3	
one co	STOP READ THE LADEL GALLONS NET
2	TOXAPHENE 8-E
	Emulsifiable Liquid Insecticide
-	ACTIVE ING REDIENTS: Toxaphene (Technical Chlorinated Camphene containing 67%-69% chlorine) 71.6% Xylene-Range Aromatic Hydrocarbon Solvent
	TOTAL 100.0% Contains 8 lb. Toxaphene per gal.
	WARNING KEEP OUT OF REACH OF CHILDREN
	Read Complete Precautions on Right Side Panel
E	E.P.A. Reg. No. 9782-8 Manufactured by

Manufactured by

WOODBURY CHEMICAL COMPANY PRINCETON, FLORIDA 33030



TOXAPHENE 8-E Emulsifiable Liquid Insecticide

ACTIVE INGREDIENTS:	
Toxaphene (Trehnical Chlorinated Camphene containing 67%-69% cl	hiorine) 71.6%
Xylene-Range Argmatic Hydrocarbon Solvent	· · · · · 25.4%
INERT INGREDIENTS:	· · · · . <u>3.0%</u>
TOTAL	100.0%

Contains 8 lb. Toxaphene per gal.

WARNING **KEEP OUT OF REACH OF CHILDREN**

Read Complete Precautions on Right Side Panel E.P.A. Reg. No. 9782-8 E.P.A. Est. No. 9782-F1-1

Manufactured by

WOODBURY CHEMICAL COMPANY PRINCETON, FLORIDA 33030

Do not use this product on feed or forage to be fed to dairy animals or to animals being finished for slaughter.

ENVIRONMENTAL CAUTIONS

This product is toxic to fish, birds and other wildlife. Birds and other wildlife in treated areas may be killed. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

MAY BE ABSORBED THROUGH SKIN - MAY BE INJURIOUS TO EYES AND SKIN - COMBUSTIBLE

Do not get in eyes or on skin. Wash thoroughly after handling. Wash contaminated clothing before re-use. Flush contaminated eyes with plenty of water and get prompt medical attention. Do not breathe vapors and spray mist. Keep away from food, feedstuffs and domestic water supplies. Keep container closed. Do not use, pour, spill or store near heat or open flame. Handle concentrate in a ventilated area.

NOTICE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of MERCHANTABILITY or FITNESS FOR A PARTICU-LAR PURPOSE, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.



GENERAL USE PRECAUTIONS

Observe use limitations given on the label for specific crops. In order that pesticidal residues on food and forage crops will not exceed tolerances established by the Federal Food and Drug Administration, use only at recommended rates and intervals. Do not apply or allow to drift onto adjoining food, fiber or pasture crops, THE GROWER IS RESPONSIBLE for residues on his crops as well as for damages by drift from his property to that of others. Protective information may be obtained from your Cooperative Agricultural Extension Service.

CONTAINER DISPOSAL

Do not reuse empty container. Destroy it by perforating or crushing. Bury or discard in a safe place away from water supplies.

WARNING **KEEP OUT OF REACH OF CHILDREN** MAY BE FATAL IF SWALLOWED – COMBUSTIBLE

