



PESTS INDOORS. General Information: Use DURSBAN 2E Insecticide as a spot treatment to control cockroaches, ants, silverfish, spiders, and brown dog ticks in and around residential, industrial, institutional and commercial buildings including homes, apartment buildings, hospitals, stores, manufacturing plants, and warehouses. DURSBAN 2E may also be used as a crack and crevice treatment to control certain of the above pests in and around both food and nonfood areas of food handling establishments (i.e., any area or place in which food is held, processed, prepared and/or served, e.g., restaurants, canneries, bottling plants, grocery stores, bakeries, etc.). Dilute DURSBAN 2E with water or with a suitable base oil to make a 0.25% or 0.5% chlorpyrifos spray as shown in the following SPRAY DILUTION CHART. Apply according to directions given for pests listed. Repeat treatment as needed.

SPRAY DILUTION CHART

| Gallons of Spray Mixture Desired | Fluid Ozs. of DURSBAN 2E to Use | | Gallons of Spray Mixture Desired | Fluid Ozs. of DURSBAN 2E to Use | |
|----------------------------------|---------------------------------|--------------------|----------------------------------|---------------------------------|-------------|
| | Water-Base spray | Oil-base spray: ** | | 0.25% spray | 0.50% spray |
| 1 | 1-1/8 | 2-2/3 | 1 | 1 | 2 |
| 3 | 4 | 8 | 3 | 3 | 6 |
| 5 | 6-2/3 | 13-1/3 | 5 | 5 | 10 |
| 10 | 13-1/3 | 26-2/3 | 10 | 10 | 20 |

*Note: 1 fluid ounce equals 2 tablespoonfuls or 6 teaspoonfuls. **Calculated on basis of deodorized kerosene.

Cockroaches (including strains resistant to certain other insecticides), **Ants**, **Crickets**, **Firebrats**, **Silverfish**, **Spiders**: Use a 0.25% spray to control light infestations and a 0.5% spray to quickly reduce heavy infestations. For spot treatment, apply as a coarse spray or with a paint brush to localized areas where these pests are found or normally occur, including dark corners of rooms and closets, cracks and crevices in walls; along and behind baseboards; beneath and behind sinks, stoves, refrigerators, and cabinets, around plumbing and other utility installations. For ants apply to ant trails, also around doors and windows, and wherever else these pests may find entrance. Note: A period of 4 to 7 days is normally required for maximum effect on cockroaches.

FOOD HANDLING ESTABLISHMENTS

Places other than private residences in which exposed food is held, processed, prepared or served.

FOOD AREAS - APPLICATION LIMITED TO CRACK AND CREVICE TREATMENT ONLY

Includes areas for receiving, storage, packing (canning, bottling, wrapping, boxing) preparing, edible waste storage and enclosed processing systems (mills, dairies, edible oils, syrups). Serving areas (when food is exposed and facility is in operation) also would be considered food area.

Using a 0.50% spray apply in small amounts directly into cracks and crevices using equipment capable of delivering a pin stream of insecticide in points between different elements of construction, between equipment and floors, openings leading to voids and hollow spaces in walls, equipment legs and bases where cockroaches, ants and silverfish hide. Do not use this product in conduits, motor housings, and electrical switch boxes.

Care should be taken to avoid depositing the product onto exposed surfaces or introducing the material into the air. Avoid contamination of food or food processing surfaces.

APPLICATIONS OF THIS PRODUCT IN THE FOOD AREAS OF FOOD HANDLING ESTABLISHMENTS OTHER THAN AS A CRACK AND CREVICE TREATMENT ARE NOT PERMITTED

SERVING AREAS - Facilities where prepared foods are served, such as dining rooms, but excluding areas where foods may be prepared or held. Using a 0.50% spray apply as a spot treatment to selective surfaces such as baseboards, under elements of construction into cracks and crevices. Avoid treating surfaces likely to be contacted by food. (Do not apply when facility is in operation or foods are exposed.)

NON-FOOD AREAS - Includes garbage rooms, lavatories, food drains, sewer entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage (after canning or bottling).

Using a 0.50% spray apply as a coarse spray or with a paint brush to baseboard areas, around water pipes, surfaces behind and beneath sinks, lockers, tables, pallets and similar areas where insects hide or through which they may enter.

Brown Dog Ticks: Thoroughly apply a 0.5% spray as a spot treatment to infested areas such as pet beds and resting quarters, nearby cracks and crevices, along and behind baseboards, window and door frames and localized areas of floor and floor coverings where these pests may be present. Old bedding of pets should be removed and replaced with clean, fresh bedding after treatment. Do not treat pets with this product.

(Pests Indoors): Do not allow children or pets to contact treated surfaces until spray has dried. Remove pet and silverfish bowl (tanks) before spraying. Do not allow spray to contact food, feedstuffs, and water supplies. Thoroughly wash dishes and food handling utensils and surfaces with soap and water if they become contaminated by application of this product. Do not use DURSBAN 2E for spot treatment in poultry houses, other animal buildings, or in edible products areas of food processing plants, restaurants, or other areas where food is commercially prepared and processed. Do not use in serving areas while food is exposed.

ACCEPTED
MAR 21 1975
Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide EPA Reg. No. 9782-20



DURSBAN* 2E INSECTICIDE

To Be Applied Only By or Under the Supervision of Pest Control Operators and Other Trained Personnel Responsible for Insect Control Programs

ACTIVE INGREDIENTS:

| | |
|--|--------------|
| Chlorpyrifos [O, O-diethyl O-(3,5,6-trichloro-2-pyridyl) phosphorothioate] | 22.4% |
| Aromatic petroleum derivative solvent | 42.1% |
| INERT INGREDIENTS: | 35.5% |
| Contains 2 pounds of chlorpyrifos per gallon | Total 100.0% |

*DURSBAN® - Trademark of the Dow Chemical Company
U. S. Patent No. 3,244,586

WARNING

KEEP OUT OF REACH OF CHILDREN

Read complete precautions on right side panel

E.P.A. Reg. No. 9782-20 Manufactured by E.P.A. Est. No. 9782-FL-1

_____ GALLONS NET

WOODBURY CHEMICAL COMPANY
PRINCETON, FLORIDA 33030





1975
Federal Insecticide,
Fungicide, and Rodenticide Act
E.P.A. Reg. No. 9782-20



**WOODBURY
CHEMICAL COMPANY**

_____ GALLONS NET

DURSBAN* 2E INSECTICIDE

To Be Applied Only By or Under the Supervision of Pest Control Operators and Other Trained Personnel Responsible for Insect Control Programs

ACTIVE INGREDIENTS:

| | |
|--|--------------|
| Chlorpyrifos [O, O-diethyl O-(3,5,6-trichloro-2-pyridyl) phosphorothioate] | 22.4% |
| Aromatic petroleum derivative solvent | 42.1% |
| INERT INGREDIENTS: | 35.5% |
| Contains 2 pounds of chlorpyrifos per gallon | Total 100.0% |

*DURSBAN® - Trademark of the Dow Chemical Company
U. S. Patent No. 3,244,586

WARNING

KEEP OUT OF REACH OF CHILDREN

Read complete precautions on right side panel

E.P.A. Reg. No. 9782-20 Manufactured by E.P.A. Est. No. 9782-FL-1

USE DIRECTIONS
PESTS OUT-OF-DOORS: General Information: Use DURSBAN 2E for control of certain pests of lawns. Consult the following directions for dosage and application details to control the pests listed. Pests of Lawns and other Ornamental and Recreational Turf Grass Areas. Use DURSBAN 2E insecticide to control ants, brown dog ticks, chinch bugs, crickets, cutworms, earwigs, grasshoppers and sod webworms (lawn moths). Mix DURSBAN 2E with water according to directions given in the following table and apply as a coarse, low pressure spray using suitable application equipment. For best results, the lawn should be moist at time of treatment, and care should be taken to obtain complete and uniform coverage of pest infested areas. Spray when pests first appear; retreat when needed.

DOSAGE AND DILUTIONS FOR USE ON TURF

| Area to be Treated | Amount of DURSBAN 2E | Dilute with Water To Give | Area to be Treated | Amount of DURSBAN 2E | Dilute with Water to Give |
|--------------------|----------------------|---------------------------|--------------------|----------------------|---------------------------|
| 500 sq. ft. | 3/4 fl. oz. | 15 gallons | 10,700 sq. ft. | 1 pint | 325 gallons |
| 1,000 sq. ft. | 1-1/2 fl. oz. | 30 gallons | One Acre | 2 quarts | 1,300 gallons |
| 5,000 sq. ft. | 7-1/2 fl. oz. | 150 gallons | | | |

To control certain white grubs (larvae of European chafer and Japanese beetle): Use two to four times the amounts of DURSBAN 2E in the TURF table, i.e., 3 to 6 fl. oz. per 1,000 sq. ft. or 1 to 2 gallons per acre. Apply in enough water to obtain uniform coverage of the area treated followed immediately by thorough watering to wash the insecticide into the turf. For best results treat when grubs are young and actively feeding near the soil surface, usually during late July and August. Consult your local Agricultural Extension Service specialist for proper time to treat in your area.

Note: To help prevent infestation of buildings by ants, crickets, earwigs and sowbugs (pillbugs), treat a band of soil 6 to 10 feet wide around and adjacent to the building, also the building foundation to a height of 2 to 3 feet, where pests are active and may find entrance. Use the same dosage and dilution recommended for control of these pests on turf as shown in the above table. Apply as a coarse spray to thoroughly and uniformly wet the band area.

Note: Mosquitoes coming to rest on areas treated for control of turf pests will be killed for varying periods of time after treatment, depending on exposure of treated areas to weathering conditions.

Warning (Pests Out-of-Doors): Keep children and pets off treated areas until spray has dried. Keep out of lawn and garden fishpools and other bodies of water. Fish may be killed if their waters are contaminated with this product. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated.

ENVIRONMENTAL CAUTIONS

This product is toxic to fish, birds and other wildlife. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

GENERAL USE PRECAUTIONS


This product is highly toxic to bees exposed to direct treatment or residues on plants. Protective information may be obtained from your Cooperative Agricultural Extension Service. Promptly crush empty containers and bury them and wastes in locations away from domestic water supplies.

WARNING

KEEP OUT OF REACH OF CHILDREN
MAY BE FATAL IF SWALLOWED - MAY BE ABSORBED THROUGH SKIN
MAY BE INJURIOUS TO EYES AND SKIN - COMBUSTIBLE

Do not get in eyes or on skin. Wash thoroughly after handling. Wash contaminated clothing before re use. Flush contaminated eyes with plenty of water and get prompt medical attention. Avoid breathing vapors and spray mist. Avoid spraying food crops. Keep away from food, feedstuffs and domestic water supplies. Keep container closed. Do not use, pour, spill or store near heat or open flame. Handle concentrate in a ventilated area.

Note to Physician: Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. Atropine is an antidote.
NOTICE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of MERCHANTABILITY or FITNESS FOR A PARTICULAR PURPOSE, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.



WOODBURY CHEMICAL COMPANY
PRINCETON, FLORIDA 33030