



Net Contents \_\_\_\_\_ Pound(s)

Carbaryl RQ  
ORM-A NA2757

SEVIN® 5 DUST

I N S E C T I C I D E

ACCEPTED  
with COMMENTS  
in EPA Letter Dated:

APR 27 1984

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.

9782-62

READY TO USE DUST  
FOR CONTROL OF NUMEROUS  
INSECTS AS LISTED ON

- ORNAMENTAL SHRUBS & FLOWERS - SMALL FRUITS
- DOGS - CATS - POULTRY
- VEGETABLES - TOBACCO - RECREATIONS AREAS
- PEST CONTROL IN & AROUND BUILDINGS

ACTIVE INGREDIENT:	
Carbaryl(1-naphthyl N-methylcarbamate)	5.0%
INERT INGREDIENTS:	95.0%
	Total 100.0%

® Sevin - Reg.T.M. of Union Carbide Corporation for Carbaryl

KEEP OUT OF REACH OF CHILDREN  
CAUTION

See Side Panel For Additional Precautionary Statements

EPA Reg. No. 9782-62

EPA Est. No. 9782-FL-1

STATEMENT OF PRACTICAL TREATMENT

If Swallowed - Call a physician or Poison Control Center immediately. Induce vomiting by giving victim 1 or 2 glasses of water and touching back of throat with finger. Never give anything by mouth to an unconscious person.

If In Eyes - Flush eyes with plenty of water. Get medical treatment.

If On Skin - Remove contaminated clothing and wash affected area with soap and water. Get medical attention.

NOTE TO PHYSICIAN - Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine is antidotal. Do not use 2-PAM, opiates, or cholinesterase inhibiting drugs.

WOODBURY CHEMICAL COMPANY  
Princeton, Florida 33032

BEST DOCUMENT AVAILABLE

"PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que le etiqueta haya sido explicado amplimente".

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**SMALL ORNAMENTAL SHRUBS AND FLOWERS** - This product may be used on herbaceous annual, biennial and perennial flowering plants and small woody shrubs such as Arborvitae (evergreen), Azalea, Carnations, Cyranthemums, Gladioli, Zinnia, Roses, Birch, Dogwood, Elm, Hydrangea, Hybiscus, Junipers, Lilac, Maple, Oak, and Pine. Do not apply to Boston Ivy Or Virginia Creeper.

**Insects Contolled: Flowering Plants** - Blister Beetles, Flea Beetles, Boxelder Bugs, Japanese Beetle, June Beetles, Lace Bugs, Leafhoppers, Leaf-rollers, Mealybugs, Plant Bugs, Psyllids, Rose Aphids, Thrips (Exposed).

**Woody Small Shrubs** - Apple aphids, Bagworms, Birch Leaf-miner, Boxelder Bugs, Boxwood Leafminer, Cankerworms, Eastern Spruce Gall Aphid, Elm Leaf Aphid, Elm Leaf Beetles, Gypsy Moth, Japanese Beetle, June Beetles, Lace Bugs, Leafhoppers, Leafrollers, Mealybugs, Mimosa Webworms, Oak Leafminers, Orange Tortrix, Periodical Cicada, Puss Caterpillar, Rose Aphid, Rose Slug, Sawflies (exposed), Scale Insects, Tent Caterpillars, Thrips (Exposed), and Willow Leaf Beetles.

**How To Apply:** Apply to plants using a hand or power duster to thoroughly cover the upper and lower portions of leaves, stems and flowers. Apply in such a manner as to walk away from the dusted area. Do not walk into airborne dust. Apply approximately 1 pound of dust per 1,000 sq. ft. of area or 40 pounds of dust per acre. Treat when insects are first seen or their damage is eminent and repeat as needed for control.

**SMALL FRUITS** - May be used on Blackberries, Raspberries, Dewberries, Loganberries, Boysenberries, Grapes and Strawberries when insect damage is first seen or is eminent. For control of Japanese Beetle, Leafrollers, and European Raspberry Aphid, and Leafhoppers use 1 lb. dust per 1,000 sq. ft. of plants at 7 to 10 day intervals as needed for control. Do not apply within 7 days of harvest on the above except Grapes & Strawberries. See Below.

For Grapes apply to foliage in spring before first brood of leafrollers larvae emerge. Apply using 1 pound per 1,000 sq. ft. to thoroughly dust all plant parts. Will control, European Fruit Lecanium, Grape Leaf-foller, Grape Leafhoppers, and Western Grape Leaf Skeletonizer, Grape Berry Moth Larvae, Japanese Beetle, June Beetles, Orange Tortrix, Omnivorous Leafroller, and Redbanded Leafrollers. Grapes may be harvested same day.

For Strawberries thoroughly apply 1 lb per 1,000 sq. ft. of plantings to control Meadow Spittlebug, Strawberry Leafroller, and Strawberry Weevil. Apply when insects are seen or their damage is a problem. Do not apply within 1 day harvest.

**TOBACCO** - Use in Plant Beds for control of Tobacco Flea Beetle using approx.  $\frac{3}{4}$  to 1 pound per 1,000 sq. ft. Apply when insects are first seen and repeat until plants are moved to field.

For Use on Field Plants for control of Budworms, Flea Beetles, Japanese Beetle, June Beetles, and Tobacco Hornworm apply at the rate of 20-40 pounds per acre to thoroughly cover plant parts. Begin treatments when worms are small. Avoid excessive dust in the "Buds" as plant injury may result. If late applications are needed allow 3 days before priming or cutting.

THIS DOCUMENT AVAILABLE

## DIRECTIONS FOR USE(Cont.)

9782-67

PEST CONTROL IN & AROUND BUILDINGS - For use in and around buildings such as Homes, Apartments, Warehouses, Barns, and Municipal & Recreational Areas. DO NOT USE IN FOOD AREAS of food handling establishments, restaurants, or other areas where food is commercially prepared or processed. Do not use in service areas while food is exposed. Avoid application to surfaces where visible dust residues are objectionable. Apply with a Hand or Power Duster.

How To Apply as a Crack & Crevice Treatment Only: For control of Ants, Cockroaches, Crickets, Millipedes and Silverfish use a "Squeeze" or "Puff" Duster to apply dust thoroughly to all cracks and crevices, baseboards, corners, walls, door sills, and window frames, quarters of household pets, ant trails, outside perimeters of dwellings and other areas where insects may rest or hide such as under cabinets, refrigerators, stoves, sinks, and around bookcases or in ATTIC areas or other storage areas where insects may breed.

For Fleas and Brown Dog Ticks thoroughly dust all cracks and crevices around pet living quarters, under carpets, pets' bedding and resting areas.

FOR USE ON DOGS & CATS - For Fleas and Brown Dog Ticks dust liberally over the entire pet, avoiding the eyes, and thoroughly rub into the hair or fur. Comb out the dead ticks and fleas a few hours after treatment. Repeat at weekly intervals if needed. Avoid treatment of kittens or puppies less than 4 months of age. Do not allow powder to enter pets' eyes or mouth.

For Pet Sleeping Quarters dust liberally in and around sleeping quarters, kennels, dog runs, and dog houses using a hand "Squeeze" or "Puff" duster. Clean and replace pet bedding. Repeat as needed for control.

POULTRY INSECT CONTROL - Bird Treatment: Northern Fowl Mite and Lice - Use 1 pound per 100 birds. Apply thoroughly with a "Squeeze" duster or a shaker can. Repeat in 4 weeks if necessary.

Premise Treatment: For Chicken Mites, Fleas, and Bedbugs use 1 lb. per 40 sq.ft. of litter or floor space and to roosts with a "duster". Force dust into cracks and crevices. Repeat as necessary. For heavy infestations birds should always be directly treated. Do not contaminate nests, eggs, and feeding and watering troughs. Do not treat premises nor birds within 7 days of slaughter.

This treatment may also be used on Ducks, Geese, Gamebirds, Pigeons and Turkeys. Do not apply to Gamebirds within 7 days of slaughter.

VEGETABLE INSECT CONTROL - Dust vegetable plants to thoroughly cover upper and lower portions of leaves, stems, flowers and fruit. Apply when insects or their damage appears. Treat in morning or evening when it is cool. Apply with a "Squeeze" or power duster. Repeat as necessary, generally at 7 day intervals for best insect control. Observe waiting period between treatment and harvest.

BEST DOCUMENT AVAILABLE

9782-62

DIRECTIONS FOR USE(Cont.)  
DUSTING RA.

VEGETABLE PLANTS	INSECTS	1,000 Sq.Ft.	Acre	(DAYS BEFORE HARVEST) COMMENTS
Asparagus	Asparagus Beetles	1/2-1 lb.	20-40 lbs.	Do not treat more than once every 3 days. ( 1 DAY ).
Beans(Green, Lima, Snap, Navy, Cowpeas, Black-eyed).	Blister Beetle, Mexican Bean Beetle, Alfalfa Caterpillar, Bean Leaf Beetle, Cucumber Beetle, Flea Beetles, Leafhopper, Thrips, Velvet Caterpillar, Western Bean Cutworm, Armyworm, Corn Earworm, Stink Bug, Tarnished Plant Bug, Webworms.	3/4-1 lb.	30-40 lbs.	Harvest same day as treatment
Cabbage, Broccoli, Brussel Sprouts, Cauliflower, Kohlrabi	Flea Beetles, Harlequin Bug, Armyworm, Corn Earworm, Fall Armyworm, Imported Cabbage Worm.	3/4-1 lb.	30-40 lbs.	Wait 3 days before harvest. Dust base of plant at soil level.
Cucumbers, Melons, Squash, Pumpkins	Pickleworm, Melonworm, Cucumber Beetles, Flea Beetles, Leafhoppers, Squash Bugs,	1/3 lb.	15-20 lbs.	Avoid excessive application Do not apply to wet foliage.
Carrots, Celery, Parsnips, Parsley, Radish	Flea Beetles, Leafhoppers, Harlequin Bug, Spittlebugs, Lygus Bug, Armyworm, Corn Earworm, Tarnished Plant Bug, Stink Bug,	3/4-1 lb.	30-40 lbs.	Wait 3 days before harvest of Parsnips & Radishes; 14 days for Celery & Parsley. Same day harvest for Carrots. Treat on a 5 to 7 day schedule.
Dandelions, Lettuce, Endive, Salsify	Flea Beetles, Leafhopper, Harlequin Bug, Spittlebug, Armyworm, Corn Earworm, Fall Armyworm, Imported Cabbage Worm, Stink Bugs, Tarnished Plant Bug.	3/4-1 lb.	30-40 lbs.	Head Lettuce & Salsify 3 DAYS. 14 DAYS on Dandelion, Endive, Escarole, Leaf Lettuce, Salsify tops.
Tomato, Potato, Eggplant, Peppers	Colorado Potato Beetle, Flea Beetles, Leafhoppers, European Corn borer, Fall Armyworm, Tomato Fruit Worm, Tomato Horn Worm, & Pinworm.	3/4-1 lb.	30-40 lbs.	Harvest same day as treatment. Use at 5-7 day intervals.

BEST DOCUMENT AVAILABLE

9782-62

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION:** Harmful if inhaled or swallowed. Do not breathe dust. Do not take internally. Skin contact may be harmful. Avoid contact. Wash hands thoroughly after handling and before eating or smoking. Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons. Do not enter treated areas without protective clothing until dusts have settled.

ENVIRONMENTAL HAZARDS

This product is highly toxic to bees exposed to direct treatment or residues on weeds or crops. Protective information may be obtained from your cooperative Agricultural Extension Service. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply directly to water. Do not apply when weather conditions favor drift of dust from area being treated.

\*\*

STORAGE & DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.  
**STORAGE-** Store in original container in a cool dry place out of reach of children. **PESTICIDE DISPOSAL** - Securely wrap original container in several layers of newspaper and discard in trash.  
**CONTAINER DISPOSAL** - Do not reuse container (bag). Discard bag in trash.

**NOTICE** - Buyer assumes all risk and liability of use, storage and/or handling of this material not in accordance with the terms of this label.

\*\* The above Storage and Disposal Statement has been revised to conform to EPA letter dated August 5, 1983 by J. S. Ellenberger PM-12.

BEST DOCUMENT AVAILABLE

U.S. ENVIRONMENTAL PROTECTION AGENCY  
OFFICE OF PESTICIDES PROGRAMS  
REGISTRATION DIVISION (WH-567)  
WASHINGTON, D.C. 20460

EPA REGISTRATION NO.

DATE OF ISSUANCE

000000

4/27/84

TERM OF ISSUANCE

INDIVIDUAL REGISTRATION

NAME OF PESTICIDE PRODUCT

NOTICE OF PESTICIDE:  REGISTRATION

REREISTRATION

(Under the Federal Insecticide, Fungicide,  
and Rodenticide Act, as amended)

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

W. H. ...  
P.O. Box ...  
... ..

BEST DOCUMENT AVAILABLE

NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(B) provided that you:

1. Submit and/or date all data requirements for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrations of similar products to be submitted.

2. Advise the Agency of any changes in your product to be made before you release the product for sale.

3. Submit a copy of your label to the Agency for review before you release the product for sale. You must submit a copy of your label to the Agency for review before you release the product for sale.

4. If you are a manufacturer, submit a copy of your label to the Agency for review before you release the product for sale. You must submit a copy of your label to the Agency for review before you release the product for sale.

A copy of the label is being returned to you with this registration.

ATTACHMENT IS APPLICABLE

SIGNATURE OF APPROVING OFFICIAL

DATE