



SEVIN[®] 5% BAIT

**FOR THE CONTROL OF CUTWORMS, ARMYWORMS, FIELD CRICKETS,
DARKLING BEETLES, GRASSHOPPERS, SOWBUGS AND ANTS.**

ALSO FOR CONTROL OF MOLE CRICKETS

ACTIVE INGREDIENT:

Carbaryl (1-Naphthylmethylcarbamate) 5.0%

INERT INGREDIENTS: 95.0%

Total 100.0%

*Sevin[®] is the Registered Trademark of Union Carbide for Carbaryl Insecticide.

FOR VEGETABLES, FLOWERS, FRUIT, LAWNS & ORNAMENTAL PLANTS

KEEP OUT OF REACH OF CHILDREN AND DOMESTIC ANIMALS

CAUTION

STATEMENTS OF PRACTICAL TREATMENTS

IF SWALLOWED: Call a physician or Poison Control Center immediately. Induce vomiting by giving victim 1 or 2 glasses of water and touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person. IF IN EYES: Flush eyes with plenty of water. Get medical attention. IF ON SKIN: Remove contaminated clothing. Wash affected area with soap and water.

NOTE TO PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal. Do not use 2-PAM, opiates, or cholinesterase inhibiting drugs.

See Side/Back Panel For Additional Precautionary Statements

EPA Reg. No. 9782

EPA Est. No. 9782-FL-1

NET CONTENTS

POUNDS ()

ACCEPTED

MAY 8 1985
Under Federal Insecticide,
Fungicide and Rodenticide Act,
this pesticide is registered under
EPA Reg. No.

WOODBURY CHEMICAL COMPANY
PRINCETON, FLORIDA 33032

SEVIN
5%
BAIT

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

HARMFUL IF INHALED OR SWALLOWED. Avoid breathing of Dust. Do not take internally. Avoid contact with skin or eyes. Wear regular long sleeved work clothing. Change to clean clothing daily. Wash hands and face before eating. Wash thoroughly after handling.

NOTE TO PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal. Do not use 2-PAM, opiates or cholinesterase inhibiting drugs.

AVOID CONTAMINATION OF FOOD, FEED, WATER SUPPLIES, STREAMS, AND PONDS DURING APPLICATION OR WHEN CLEANING EQUIPMENT. Keep pets off treated areas.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

IMPORTANT PRE-HARVEST AND GRAZING USE INFORMATION

Tolerances established under the Federal Food, Drug, and Cosmetic Act permit the sale of crops bearing probable Sevin residues when SEVIN is used in accordance with label directions. If SEVIN insecticide is used in accordance with label directions, these crops including bean vines, carrot tops, pea vines, sugar beet tops, corn forage, and cotton forage may be grazed or harvested for use as feed for dairy and meat animals without resulting in residues in milk, or meat in excess of established tolerances.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. **STORAGE:** Store product in a cool, dry, locked place out of reach of children and domestic animals. **DISPOSAL:** Do not reuse empty container. Wrap and put in trash.

Apply granules as a broadcast application over the soil with ground equipment. Treatments may be repeated as necessary. Dosages from 1/2 lb. to 1 lb. per 1,000 sq. ft. will control Ants, Cutworms, Armyworms, Crickets, Darkling Beetles, Grasshoppers and Sowbugs. "Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl labels."

CROP	DOSAGE RATE 1,000 SQ. FT.	DAYS BETWEEN FINAL APPLICATION AND HARVEST
Cucumbers, Squash, Melons, Peanuts	1/2 lb	None
Alfalfa, Peas, & Cotton	1/2 lb	3 Days - 7 Days
Beans, Carrots, Corn, Eggplant, Onions, Peppers, Potatoes, Sweet Corn, Tomatoes	1 lb	None
Asparagus, Strawberries	1 lb	1 Day
Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Head Lettuce, Garden Beets, (roots), Horseradish, Parsnips, Rutabagas, and Turnips (roots)	1 lb	3 Days
Sugar Beets, Collards, Endive, Garden Beets (tops), Kale, Leaf Lettuce, Parsley, Spinach, Swiss Chard, Turnips (tops)	1 lb	14 Days
Tobacco	1/2 lb	None
Fruits - Apples, Apricots, Cherries, Citrus, Grapes, Nectarines, Peaches, Pears, Plums Ornamentals - Annual plants, herbaceous plants and shrubs, trees, woody plants	1 lb	Apples - 1 Day Apricots - 3 Days Cherries - 1 Day Citrus - 5 Days Nectarines - 3 Days Peaches - 1 Day Pears - 1 Day Plums - 1 Day

LAWN, TURF AND ORNAMENTAL PLANTS

Treatments may be repeated as necessary to control Cutworms, Armyworms, Field Crickets, Mole Crickets, Darkling Beetles, Grasshoppers, and Sowbugs.

LAWNS AND TURF - Treat entire area evenly at a dosage of 3/4 to 1 lb. per 1000 Sq.Ft.. Apply with a rotary spreader to obtain uniform coverage, giving particular attention to edges of sidewalks and along porch and patio foundations. For best results cut lawn, remove clippings and water thoroughly before treatment.
ORNAMENTAL PLANTINGS - Apply evenly to soil around plants, shrubs and trees at a rate of 3/4 to 1 lb./1000 Sq.Ft. For best results, areas to be treated should be damp.

NOTICE - Buyer assumes all risk and liability of use, storage and/or handling of this material not in accordance with the terms of this label.