


1983 8/1



AUTHORIZED FOR USE  
IN FEDERALLY  
INSPECTED MEAT AND  
POULTRY PLANTS



ACCEPTED

AUG 18 1986

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide

NET CONTENTS 5 GALLONS  
EPA Reg. No. 9782-CC

# FOOD PLANT FOGGING INSECTICIDE

APPLY WITH EQUIPMENT DESIGNED AND ADJUSTED TO DELIVER  
A TRUE AEROSOL MIST - Also Apply as a Coarse Wet Spray

For Control of Stored Food Insects, Roaches, Ants, Fleas, Clothes Moths,  
Flies, Gnats and Other Flying or Crawling Insects as Listed

**ACTIVE INGREDIENTS:**

Pyrethrins .....	0.5%
Piperonyl Butoxide, Technical .....	5.0%

<b>INERT INGREDIENTS:</b> .....	<b>94.5%</b>
	Total 100.0%

Equivalent to 4.0% (butylcarbityl) (6-propylpiperonyl) ether and 1.0% related compounds.

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION** SEE SIDE/BACK PANEL FOR  
ADDITIONAL PRECAUTIONARY STATEMENTS

**STATEMENTS OF PREVENTIVE TREATMENT**

IF SWALLOWED: Call a physician or Poison Control Center immediately. Do not induce vomiting unless directed to do so by a physician. Aspiration is a potential hazard. This product contains a petroleum Distillate.

IF IN EYES: Flush eyes with plenty of water. Contact a physician if irritation persists.

IF ON SKIN: Wash affected area with soap and water.

IF INHALED: Remove victim to fresh air. Apply artificial respiration if indicated.

EPA REG. NO. 8782-66

OCT. 85 EPA EST. NO. 9742-FL-1





### DIRECTIONS FOR USE

**Space Applications:** For rapid control of flies, fruit flies, gnats, mosquitoes, skipper flies, wasps, hornets, Angoumois grain moths, Mediterranean flour moths, Indian meal moths, and tobacco moths

Close windows and doors and shut off ventilating systems. Apply with a good machine adjusted to deliver a fine mist or fog. In Official Meat and Poultry Establishments use a fogger or vaporizer adjusted to deliver an aerosol spray (no droplets over 50 microns in diameter and 80% less than 30 microns). Apply at the rate of 1 ounce per 1000 cubic feet of space. Direct the spray toward the ceiling and upper corners of the area and behind obstructions. Keep the area closed for at least 15 minutes. Vacate the treated area and ventilate before reoccupying. Repeat treatment as necessary. In other than Official Meat and Poultry Establishments a combination of spot surface treatment and fogging is recommended.

**In Barns, Poultry Houses and Storage Rooms:** To control house flies, stable flies and horn flies in Barns, Poultry Houses, Warehouses, Storage Bins and Storage Rooms dilute 1 part of material with 4 parts of oil. Apply as a fine mist over the livestock in stanchions or holding pens. Use a fogger or mist generator to direct spray above the animals and to the center of the area where animals are housed. Apply 1 ounce per 1000 cubic feet uniformly throughout the entire area sprayed. Do not ventilate the area for at least 10 minutes after spray has been applied. If used in Dairy Barns, aerate the barn thoroughly before milking. Repeat daily in the morning and afternoons, or as required.

**Surface Spraying:** To control roaches, ants, silverfish, spiders, crickets, clover mites, cheese mites, granary weevils, rice weevils, confused flour beetles, rust red flour beetles, sawtoothed grain beetles, spider beetles, cigarette beetles, Drugstore beetles, mealworm, grain mites, and caddises. Use a good sprayer adjusted to deliver a coarse wet spray. Direct the spray into hiding places, cracks and crevices, under pallets, around containers of stored foods, around the base of machinery, behind shelves and drawers. For silverfish, spray bookcases. For ants, spray trails, nests and points of entry. If surface application only is to be used, spray floors, walls, and other surfaces at a rate of 1 gallon to 750 square feet of surface.

To control carpet beetles. Spray edges of carpeting and under carpeting and rugs. Make a localized application to floors and baseboards. Spray directly into cracks, closets, and infested areas of shelving. Repeat treatment as necessary.

To control bedbugs. Spray mattresses lightly, particularly around tufts and seams. Take beds apart and spray in all joints. Treat baseboards, moldings, and floors. Repeat treatment as needed.

To control fleas and brown dog ticks. Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, moldings, window and door frames, and localized areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Concurrent treatment of animals with an approved insecticide is recommended. Repeat treatment as needed.

To control clothes moths. Remove any infested articles from storage, brush thoroughly and air for several hours in sunlight, if possible. Apply spray liberally to empty chests, closets, bureaus, and other storages, directing the spray into cracks, joints, and crevices. After airing, infested articles may be treated lightly keeping nozzle at least three feet from fabric to avoid staining. Store clothes in moth tight containers or repeat treatments monthly.

**OUTDOOR APPLICATIONS: Adult Mosquito Control:** Apply as a fine mist to thoroughly treat the entire area. For best results, treat when the wind velocity does not exceed 5 miles per hour. Treat shrubbery or vegetation where mosquitoes may rest, low marshy areas, stagnant pools or ponds and shorelines. Apply at a rate of 10 fluid ounces per acre (0.2 fluid ounce per 1000 square feet).

Using truck-mounted conventional misting equipment traveling at 5 m p h to treat a 300 swath, calibrate equipment to deliver 0.23 fluid ounce per minute.



**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION:** Harmful if swallowed or absorbed through the skin. Avoid breathing vapors or spray mist. Avoid contact with skin and eyes.

Do not apply directly to food. In commercial food handling facilities, cover or remove any food and food processing equipment before application. Do not apply while food processing is under way. After space spraying in meat packing plants, bakeries and all other food processing plants wash all equipment (benches, shelving, etc.) with a suitable effective detergent cleaning compound followed by a potable water rinse after the spraying where exposed food will be handled. All food processing surfaces and utensils should be covered during treatment or thoroughly washed before use.

Remove pets and birds, cover fish aquariums before spraying.

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish. Do not apply to any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes.

**PHYSICAL OR CHEMICAL HAZARDS**

Do not use, pour, spill or store near heat or open flame.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**STORAGE AND DISPOSAL**

**Do not contaminate water, food or feed by storage or disposal.**

**STORAGE:** Store product in original container in a cool, dry, locked place out of reach of children.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by other procedures approved by State or local authorities.

**NOTICE:** Buyer assumes all risk and liability of use, storage and/or handling of this material not in accordance with the terms of this label.