


DO NOT REUSE EMPTY DRUM. RETURN TO DRUM RECONDITIONER OR DESTROY BY PERFORATING OR CRUSHING AND BURYING IN A SAFE PLACE.

CAUTION:
This product is toxic to fish. Keep out of any body of water. Apply this product only as specified on this label.

5 GALS.

INSECTICIDE, MALATHION EMULSIFIABLE CONCENTRATE



Contains 60 grams of technical malathion per 100 ml.
(Approximately 5 lbs. technical malathion per gallon.)

WARNING
KEEP OUT OF REACH
OF CHILDREN

ACTIVE INGREDIENTS:

Malathion (0,0-dimethyl dithiophosphat of diethyl mercaptosuccinate) ... 57%
Aromatic petroleum derivative ... 38%

INERT INGREDIENT: ... 5%

CAUTION! FLAMMABLE!

Harmful if swallowed or absorbed through the skin.
Avoid prolonged breathing of spray mist.
Avoid repeated contact with eyes or skin.
Wash thoroughly after using.
Avoid contamination of foodstuffs or mess gear.
Keep out of reach of children.
Do NOT apply as a complete coverage residual spray to interior surfaces.
Wear coveralls, gloves and use respirator or mask.

NOTE TO PHYSICIAN: Atopine is antidotal.

WARNING!

Spray should not be applied in conjunction with sulfur sprays, or if sulfur residue is present.

DIRECTIONS FOR USE

CONTROL OF RESISTANT HOUSE FLIES

Use to control house fly populations that have developed resistance to DDT and other chlorinated hydrocarbon insecticides.

Sugar-Malathion Bait

Add 1/2 pound (1/2 pint) of 57% malathion emulsifiable concentrate and 3 teaspoonfuls of lamp black to 14 1/2 pounds of granulated sugar. Mix thoroughly with a paddle or by shaking or rolling in a container with a tightly fitting lid.

Cornmeal-Malathion Bait

This bait is satisfactory on damp surfaces where sugar bait is unsuitable. Stir 12 1/4 pounds of cornmeal in a suitable container and slowly add 1/2 pound of 57% malathion emulsifiable concentrate and 1/4 lb. vegetable oil to produce a uniform mix. Then add 3 teaspoonfuls of lamp black and 2 pounds of confectioners sugar. The oil improves the bait but is not absolutely essential and may be omitted if desired.

Application of Baits

Place the bait in a shaker-top can or jar. Scatter the bait thinly in places where flies congregate around mess halls and around garbage racks. **DO NOT EXPOSE BAIT IN DISHES OR OTHER CONTAINERS.** Apply in mess halls at a rate of 1 oz. to 300 to 400 sq. ft. on floors, window sills, ledges, porches, and other locations where the baits will not contaminate food, utensils, or surfaces used in the preparation of serving of food. Heavier applications may be required in exterior locations. Apply baits daily for a week to bring house fly population under control. Less frequent applications will do thereafter.

Malathion Sprays

As a contact and residual spray for the control of house flies resistant to DDT and other chlorinated hydrocarbon insecticides, use 1/2 lb. (1/2 pint) of 57% malathion emulsifiable concentrate in water to make 3 gallons. Treat garbage cans, racks, dumps, or barns, but do not apply in buildings that are occupied by people or where food is prepared or served. Use about 1 gal./1000 sq. ft. on painted or non-absorbent surfaces and about 2 gals./1000 sq. ft. on other surfaces.

Control of Resistant Cockroaches and Resistant Fleas

Add 1 lb. (1 pint) of 57% malathion to 3 gals. water and mix thoroughly. This makes a finished spray containing 2 1/2% actual malathion. Apply as a coarse wet spray only to actual and potential cockroach harborages. To control fleas, apply at the rate of 1 to 2 gals./1000 sq. ft. on floor or ground surfaces in infested kennels, yards, rat harborages, and similar infested locations. Avoid contamination of animal food and water containers. Do not apply in buildings where food is prepared or served.

Sold by

YUKON SERVICE CO.
NEW YORK, N. Y.

U.S.D.A. Reg. No. 9791-5