

MuniChem

D E A D S U R E

KILLS RATS AND MICE

Contains WARFARIN*

CAUTION:
KEEP OUT OF REACH
OF CHILDREN
SEE OTHER PRECAUTIONS
LISTED ELSEWHERE

Ingredients:	
Active: Warfarin	
3-(a-acetylbenzyl)-4-	
hydroxycoumarin*.....	0.025%
Inert:.....	99.975%
	<hr/>
	100.000%

USDA Reg. No.

*Warfarin is a patented product discovered in the laboratories of Dr. Karl Paul Link, Biochemistry Department, University of Wisconsin, Wisconsin Alumni Research Foundation Compound #42.

IMPORTANT: Baiting should be continued as long as evidence of feeding is observed. Place baits in locations where rats and mice feed, water or travel. Bait placements along walls, in corners and in concealed locations where rats and mice can feed unobserved are desirable. Where a continuous source of infestation is present from nearby dumps or fields, permanent bait stations should be used and bait placed in them as often as necessary. Contaminated or spoiled baits should be replaced promptly.

CAUTION: This rat killer contains as its active ingredient an anti-coagulant chemical, therefore, it must be protected from or placed where humans, livestock, poultry or pets will not eat them. In case it is eaten, give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Call a physician immediately.

Note to Physicians: When a human has been known to have ingested any of this rat killer, blood transfusions combined with intravenous injections and oral doses of Vitamin K are indicated as in the case of hemorrhage caused by overdoses of Dicoumerol.

DIRECTIONS

RATS: Place a small amount, four ounces or more on a piece of paper in locations frequented by rats or where rats have been known to feed. Keep out of reach of dogs, cats, poultry, livestock, etc. by using bait boxes, cages or other suitable means. An uninterrupted supply of bait should be maintained for at least 10 days.

