

PM31 9807-5 1/2

US ENVIRONMENTAL PROTECTION AGENCY OFFICE OF PESTICIDES PROGRAMS REGISTRATION DIVISION (TS-767) WASHINGTON, DC 20460  <b>NOTICE OF PESTICIDE:</b> <input checked="" type="checkbox"/> REGISTRATION <input type="checkbox"/> REREISTRATION (Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended)	EPA REGISTRATION 9807-5	DATE OF ISSUANCE
	TERM OF ISSUANCE	<b>OCT 16 1985</b>
	NAME OF PESTICIDE PRODUCT  Sanitox	

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

\_\_\_\_\_  
 \_\_\_\_\_  
 Texas Department of Correction  
 Soap and Detergent Plant  
 1 Circle Drive  
 Sugarland, TX 77478

**NOTE:** Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration/registration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data.
2. Make the labeling change listed below before you release the product for shipment:
  - a. Add the phrase "EPA Registration No. 9807-5."

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

*John H. Lee*  
 John H. Lee

Product Manager (31)  
 Disinfectants Branch  
 Registration Division (TS-767C)

Enclosure

ATTACHMENT IS APPLICABLE

SIGNATURE OF APPROVING OFFICIAL	DATE
---------------------------------	------

**Sanitizing of Food Processing Equipment and Other Hard Surfaces in Food Contact Locations.** For sanitizing food processing equipment, dairy equipment, food utensils, dishes, silverware, glasses, sink tops, countertops, refrigerated storage and display equipment and other hard non-porous surfaces. No Potable water rinse is required.

Wash and rinse all articles thoroughly, then apply a solution of 1 oz. SANITEX in 4 gallons of water (150ppm active). Surfaces should remain wet for at least one minute followed by adequate draining and air drying. Fresh solution should be prepared daily or when use solution becomes visibly dirty. For mechanical application, use solution may not be reused for sanitizing applications.

Apply to sink tops, countertops, refrigerated storage and display equipment and other stationary hard surfaces by cloth or brush or mechanical spray device. No Potable water rinse is required.

Dishes, silverware, glasses, cooking utensils and other similar size food processing equipment can be sanitized by immersion in a 1 oz./4 gallon solution of SANITEX. No Potable water rinse is required.

At 1 oz./4 gallons SANITEX fulfills the criteria of Appendix F of the Grade "A" Pasteurized Milk Ordinances 1978 Recommendations of the U.S. Public Health Services in waters up to 800 ppm of hardness calculated as CaCO<sub>3</sub> when evaluated by the AOAC Germicidal and Detergent Sanitizer Method against Escherichia coli and Staphylococcus aureus.

The udders, flanks and teats of dairy cows can be sanitized by washing with a solution of 1 oz. SANITEX in 4 gallons of warm water. No Potable water rinse is required. Use a fresh towel for each cow. Avoid contamination of sanitizing solution by dirt and soil. Do not dip used towel back into sanitizing solution. When solution becomes visibly dirty, discard and provide fresh solution.

### Precautionary Statements

**Hazard to Humans and Domestic Animals**

## DANGER

Keep Out of Reach of Children.

Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin, or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful if swallowed. Do not breathe spray mist. Avoid contamination of food.

### Storage and Disposal

- Do not contaminate water, food, or feed by storage disposal.
- Do not store on side.
- Avoid creasing or impacting of side walls.

#### PESTICIDE DISPOSAL

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray or mixture of insecticide is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

#### CONTAINER DISPOSAL

Do Not Reuse empty polyethylene container. Wrap container and put in trash.

Texas Specification No.  
435-56-02

## SANITEX

Sanitizing-Disinfectant  
Fungicide-Virucide-Deodorizer  
with Organic Soil Tolerance  
for Hospital, Institutional  
Industrial, School, Dairy and Other Farm Use

### Active Ingredients

Octyl decyl dimethyl ammonium chloride	2.250%
Didecyl dimethyl ammonium chloride	1.125%
Dioctyl dimethyl ammonium chloride	1.125%
Alkyl (C <sub>14</sub> , 50%; C <sub>12</sub> , 40%; C <sub>16</sub> , 10%) dimethyl benzyl ammonium chloride	3.000%

### Inert Ingredients

92.500%  
100.000%

KEEP OUT OF REACH OF CHILDREN

## DANGER

### STATEMENT OF PRACTICAL TREATMENT

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution; or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression, and convulsion may be needed.

SEE LEFT PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA Registration No.

EPA Establishment No.

Net Contents One Gallon

MANUFACTURED BY:

TEXAS DEPARTMENT OF CORRECTIONS  
ONE CIRCLE DRIVE, SUGAR LAND, TX., 77478

For Sale For Use And Storage By  
Maintenance Personnel Only

DIRECT

It is a violation of product in a market labeling.

Spot SANITEX device. When applied must be sprayed until surfaces remain wet for 10 minutes daily or when the use is required.

Disinfection in Hospital Care Institutions. For disinfecting bathing areas, lavatories, toilets and other hard non-porous surfaces.

Add 3.5 oz. SANITEX to 4 gallons of water. SANITEX is a virucide. Staphylococcus aureus in the presence of the AOAC Use-Dilution Test.

Virucidal Performance. SANITEX was evaluated and found to be effective against Types 1 and 2. Virucidal on inanimate hard non-porous surfaces.

Disinfection in Institutions. SANITEX is a virucide.

Fungicidal Performance. SANITEX Trichophyton mentagrophytes used on surfaces in areas of showers and bath areas and other hard non-porous surfaces.

Disinfection of Barber Shop Combs, brushes, razors, etc. Immerse in a 0.5 oz./gallon solution.

Disinfection of Poultry Kennels. Poultry brooders and other animal quarters should be disinfected after thorough cleaning with 2 oz. SANITEX per gallon of water.

Disinfection of Poultry. SANITEX should be immersed in poultry, other animals and premises. This includes other feeding and watering troughs, droppings from floors, etc.

After disinfection, ventilate areas. Do not house poultry until treatment has been completed. All treated equipment must be rinsed with potable water.

ACCEPTED

IN COMMENTS

in EPA Letter Dated:

OCT 16 1985

Under 91807-TX-01 Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 9807-5