

Sanitizing of Food Processing Equipment and Other Hard Surfaces in Food Contact Locations. For sanitizing food processing equipment, dairy equipment, food utensils, dishes, silverware, glasses, sink tops, countertops, refrigerated storage and display equipment and other hard non-porous surfaces. No Potable water rinse is required.

Wash and rinse all articles thoroughly, then apply a solution of 1 oz. SANITEX in 4 gallons of water (150ppm active). Surfaces should remain wet for at least one minute followed by adequate draining and air drying. Fresh solution should be prepared daily or when use solution becomes visibly dirty. For mechanical application, use solution may not be reused for sanitizing applications.

Apply to sink tops, countertops, refrigerated storage and display equipment and other stationary hard surfaces by cloth or brush or mechanical spray device. No Potable water rinse is required.

Dishes, silverware, glasses, cooking utensils and other similar size food processing equipment can be sanitized by immersion in a 1 oz./4 gallon solution of SANITEX. No Potable water rinse is required.

1/4 oz. SANITEX fulfills the criteria of Appendix F of the Grade "A" Pasteurized Milk Ordinances 1978. Recommendations of the U.S. Public Health Services in waters up to 800 ppm of hardness calculated as CaCO₃ when evaluated by the AOAC Germicidal and Detergent Sanitizer Method against Escherichia coli and Staphylococcus aureus.

The udders, flanks and teats of dairy cows can be sanitized by washing with a solution of 1 oz. SANITEX in 4 gallons of warm water. No Potable water rinse is required. Use a fresh towel for each cow. Avoid contamination of sanitizing solution by dirt and soil. Do not dip used towel back into sanitizing solution. When solution becomes visibly dirty, discard and provide fresh solution.

Precautionary Statements

Hazard to Humans and Domestic Animals

DANGER

Keep Out of Reach of Children. Corrosive. Causes eye damage to skin irritation. Do not get in eyes, on skin, or on clothing. Wear goggles or face shield and rubber gloves when handling. Fatal if swallowed. Do not breathe spray mist. Avoid contamination of food.

Storage and Disposal

- Do not contaminate water, food, or feed by storage disposal.
- Do not store on side.
- Avoid creasing or impacting of side walls.

PESTICIDE DISPOSAL

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray or mixture of insecticide is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL

Do Not reuse empty polyethylene container.

Drop container and mix in trash.

SANITEX

Sanitizer-Disinfectant

Fungicide-Virucide-Deodorizer
with Organic Soil Tolerance
for Hospital, Institutional

Industrial, School, Dairy and Other Farm Use

Active Ingredients

Octyl decyl dimethyl ammonium chloride	2.250%
Didecyl dimethyl ammonium chloride	1.125%
Diocetyl dimethyl ammonium chloride	1.125%
Alkyl (C ₁₄ , 50%, C ₁₂ , 40%, C ₁₆ , 10%) dimethyl benzyl ammonium chloride	3.000%

Inert Ingredients

ACCEPTED
JUN 16 1989
Under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, for the pesticide registration number 92,500-000 EPA Reg. No. 7001-2-100-0000

KEEP OUT OF REACH OF CHILDREN
DANGER

STATEMENT OF PRACTICAL TREATMENT

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression, and convulsion may be needed.

SEE LEFT PANEL FOR ADDITIONAL
PRECAUTIONARY STATEMENTS

EPA Registration No.

EPA Establishment No.

Net Contents

09807-Tx-01

Manufactured By
Texas Correctional Industries
Sugar Land, Texas 77478

For Sale For Use And Storage By
Maintenance Personnel Only

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Apply with a cloth, mop or mechanical spray device. When applied with a mechanical spray device, surface must be sprayed until thoroughly wetted. Treated surfaces must remain wet for 10 minutes. Fresh solution should be prepared daily or when the use solution becomes visibly dirty.

Disinfection in Hospitals, Nursing Homes and Other Health Care Institutions. For disinfecting floors, walls, countertops, bathing areas, lavatories, bedframes, tables, chairs, garbage pails and other hard non-porous surfaces.

Add 35 oz. SANITEX to 45 gallons water. Apply to previously cleaned hard surfaces. At this use level SANITEX is effective against Pseudomonas aeruginosa, Staphylococcus aureus and Salmonella choleraesuis in the presence of 5% blood serum when evaluated by the AOAC Use-Dilution test.

Virucidal Performance. At a 35 oz./45 gallon use level SANITEX was evaluated in the presence of 5% serum and found to be effective against the following viruses: Herpes Simplex Types 1 and 2, Vaccinia, and Influenza A₂ (Hong Kong) on inanimate hard non-porous environmental surfaces.

Disinfection in Institutions, Industry and Schools. Add 2 oz. of SANITEX to 45 gallons of water.

Fungicidal Performance. At 0.5 oz. to 2.25 gallons of water use level SANITEX is an effective fungicide against Trichophyton mentagrophytes (the athletes foot fungus) when used on surfaces in areas such as locker rooms, dressing rooms, shower and bath areas and exercise facilities, utilizing the AOAC Fungicidal Test.

Disinfection of Barber Tools. Pre-cleaned barber tools (such as combs, brushes, razors and scissors) can be disinfected by immersing in a 0.5 oz./gallon solution of SANITEX.

Disinfection of Poultry Equipment, Animal Quarters and Kennels. Poultry brooders, watering founts, feeding equipment and other animal quarters (such as stalls and kennel areas) can be disinfected after thorough cleaning by applying a solution of 2 oz. SANITEX to 45 gallons of water. Small utensils should be immersed in this solution. Prior to disinfection, poultry, other animals and their feeds must be removed from the premises. This includes emptying all troughs, racks and other feeding and watering appliances. Remove all litter and droppings from floors, walls and other surfaces occupied or traversed by poultry or other animals.

After disinfection, ventilate buildings, coops and other closed spaces. Do not house poultry, or other animals or employ equipment until treatment has been absorbed, set or dried.

All treated equipment that will contact feed or drinking water must be rinsed with potable water before reuse.

Texas Registration No.

BEST AVAILABLE COPY