

CAF-FOG

[9838-2] 1 of 2

THIS PRODUCT MAY NOT BE APPLIED BY UNTRAINED OPERATORS ONLY

Non-Staining: May be dispensed from electric or hand sprayer or in vaporizing or fog generator....
 for control of Insect Pests in institutions, hotels, restaurants, school cafeterias, homes, hospitals, clubs, food processing plants and similar situations where an effective insecticide is required, when used as directed....
 Kills even Resistant Insect Pests.

CAF-FOG KILLS BY CONTACT

ACTIVE INGREDIENTS 100%

- Petroleum Distillate
- * Technical Piperonyl Butoxide
- Pyrethrins

* Consists of (Butylcarbitol) (6-propylphenyl) ether and related compounds.

| | | | | |
|-----------|----------------|----------------------|---------|---------|
| Roaches | Gnats | Drug Store Beetle | Worms | Scabies |
| Ants | Sand Flies | Cigarette Beetle | Borrels | Scabies |
| Crickets | Skippers | Angoumois Grain Moth | Spiders | Scabies |
| Silkworms | Flying Moths | Flies | Spiders | Scabies |
| Spiders | Carpet Beetles | Mosquitoes | Scabies | Scabies |

Control of these insects is limited to egg and stages.

PRECAUTIONS: For best effective results use in a good sprayer. Follow instructions on label. Remove all food during treatment. Remove or cover all food products so as to protect them from damage before back into operation.

PLANT TREATMENT: For control of flies, mosquitoes, gnats, ants, beetles, roaches, etc., mix 1 part CAF-FOG to 10 parts water. Fill a spray bottle with mixture. Spray all foliage, walls and ceiling surfaces of building thoroughly.

DRINK: Harmful if swallowed, injected or absorbed through skin. Do not drink. Rinse mouth with prepared water after drinking and do not eat or drink until at least one hour after application. Do not apply to unpreserved food or to food which will be eaten raw.

MAY 9 1968

| | |
|-------------------|---------|
| SEARCHED | INDEXED |
| SERIALIZED | FILED |
| 9838-2 | |
| TO AGENT COMMENTS | |

CRAWLING INSECTS:

[9838-2] 20f2

For control of roaches, waterbugs, silverfish, crickets, spiders, and brown dog ticks, thorough and repeated spraying of the infested area, directing the spray into cracks and other hiding places, is recommended. Emphasis should be placed on hitting the insects directly, and on repeated applications.

For controlling ants follow above directions for control of roaches, but similar emphasis should be placed on the treatment of ant trails and areas around door sills and window frames where the pests enter the premises. Repeated applications are required in all cases. For the control of fleas in premises follow directions for coverings, and wall cracks, as well as sleeping quarters of animals.

For control of carpet beetles spray thorough application to infested areas of carpets, localized areas of floors and baseboards, and into cracks and under carpets, where these pests may be found. Localized treatments of areas on floors, baseboards and shelves of closets may also be indicated. Repeat application as often as necessary to maintain effective control.


NOTICE: The Manufacturer and Seller warrants that this product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer does so at risk of damage.

DO NOT USE OR STORE NEAR FIRE OR OPEN FLAME

DO NOT REUSE CONTAINERS. DESTROY WHEN EMPTY.

CAUTION: Keep Out of Reach of Children.

UNIVERSITY DRUG CO., INC.
Not Concentrated



SOLD BY:

Continental Pesticide Co., Inc.
St. Louis-Kan., City 6, Mo.
Post 8725
St. Louis (Offices), Mo. 63105