

STERI-365

For use in killing weeds and grasses to keep areas free of vegetation and for treating soil prior to topping with asphalt or pavement. STERI-365 kills Johnson grass, Quack grass, Bermuda grass, Poison Ivy, Poison Oak, Bindweed, Canada Thistle, Sand Burr and many other grasses and weeds. Toxic to all types of vegetation. This material is recommended for vegetation control on non-cropped areas such as, fence-rows, ditch banks, around buildings, along roadsides, power lines, right-of-ways, and other similar areas, where there is no danger to valuable plants, or where unproductive soil is not objectionable.

DIRECTIONS

GENERAL SPRAYING: Spray to wet plants thoroughly, using power or knapsack sprayer and apply chemical uniformly over infested area. Applications should be made while weeds are small. Avoid application in hot, dry weather, as the killing action of STERI-365 is through both roots and tops of plants, and therefore moisture is necessary after application to take the chemical into the soil to kill deep-rooted perennials. Fall applications may be made in areas of 20 inches or less rainfall. Retreatment may be necessary under some conditions to keep area free of plant growth. By preventing weed growth early in the year, fire hazard from dead and dry vegetation later on, are avoided.

ANNUAL AND PERENNIAL WEEDS -- To control annuals and knock down top growth of perennials, use 1 gallon in 4 gallons of water and spray to wet plants thoroughly.

PERENNIAL WEEDS AND GRASSES -- Use 5 to 6 pints per 100 Sq. Ft. and apply in sufficient water for uniform coverage. For control of weeds with creeping rootstalks, treat soil 6 Ft. beyond visible vegetation to get creeping roots.

HOW TO USE: A gravity flow spray system or a low pressure sprayer giving uniform coverage is satisfactory. State or local agricultural authorities for specific recommendations as to dosage and time of application and for the control of other weeds.

Follow directions carefully. Consult state agricultural experiment station or extension service weed specialist for specific timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this products which are beyond the control of the seller. Buyer assumes all risks of use, storage, or handling of this material not in strict accordance with directions given herewith.

NOTE: This material is non-selective and affects all types of vegetation. Extreme care should be exercised in its use to prevent killing lawns, shrubs, trees, and other valuable plants. This includes the treatment of areas which may be underlaid by tree roots.

Due to the varied conditions of weather, soil, types of vegetation, location and time of applications, the manufacturer does not guarantee complete kill of vegetation or accept responsibility for its use or mis-use.

NET CONTENTS

STERI-365

9838-5

STERI-365

NON-SELECTIVE, VEGETATION KILLER

ACTIVE INGREDIENTS:

Sodium Chlorate (NaClO3).....18.5%

Sodium Metaborate (Na2B2O4)*10.1%

INERT INGREDIENTS.....71.4%

*Equivalent to 1.67% boron expressed as elemental boron.

USDA No.

CAUTION: KEEP OUT OF REACH OF CHILDREN

CAUTION: Harmful if Swallowed. Avoid getting in eyes; avoid breathing spray mist. Wash clothing and equipment after application. Do not leave containers where children and animals may take the chemical internally. Avoid contamination of, and storage near feed and food products. Do not use on or near turf, valuable trees or other plants. Avoid contamination of irrigation or domestic waters. The chemical should be used only where unproductive soil is not objectionable.

KEEP LID TIGHTLY CLOSED
DO NOT Store with Foodstuffs

CONTINENTAL RESEARCH CO

P. O. BOX 8428


ST. LOUIS, MISSOURI, 63113

STERI-365

NET CONTENTS

ATED

969



STERI-365

NON-SELECTIVE, VEGETATION KILLER

INGREDIENTS:

m Chlorate (NaClO3).....18.5%
 m Metaborate (Na2B2O4)*10.1%
 INGREDIENTS.....71.4%
 equivalent to 1.67% boron expressed as
 elemental boron.

USDA No.

: KEEP OUT OF REACH OF CHILDREN

: Harmful if Swallowed! Avoid getting in eyes;
 nothing spray mist. Wash clothing and equipment
 ication. Do not leave containers where children
 ls may take the chemical internally. Avoid contami-
 and storage near feed and food products. Do not apply
 - turf, valuable trees or other plants. Avoid contami-
 rrigation or domestic waters. The chemical should be
 where unproductive soil is not objectionable.

KEEP LID TIGHTLY CLOSED
 DO NOT Store with Foodstuffs

CONTINENTAL RESEARCH CORP.

P. O. BOX 8428

ST. LOUIS, MISSOURI, 63132

For use in killing weeds and grasses to keep areas free of vegetation and
 for treating soil prior to topping with asphalt or pavement. STERI-365 kills
 Johnson grass, Quack grass, Bermuda grass, Poison Ivy, Poison Oak, Bindweed,
 Canada Thistle, Sand Burr and many other grasses and weeds. Toxic to all
 types of vegetation. This material is recommended for vegetation control on
non-cropped areas such as, fence-rows, ditch banks, around buildings, along
roadsides, power lines, right-of-ways, and other similar areas, where there is
 no danger to valuable plants, or where unproductive soil is not objectionable.

DIRECTIONS

GENERAL SPRAYING: Spray to wet plants thoroughly, using power or knapsack sprayer
 and apply chemical uniformly over infested area. Applications should be made while
 weeds are small. Avoid application in hot, dry weather, as the killing action of
 STERI-365 is through both roots and tops of plants, and therefore moisture is
 necessary after application to take the chemical into the soil to kill deep-rooted
 perennials. Fall applications may be made in areas of 20 inches or less rainfall.
 Retreatment may be necessary under some conditions to keep area free of plant
 growth. By preventing weed growth early in the year, fire hazard from dead and
 dry vegetation later on, are avoided.

ANNUAL AND PERENNIAL WEEDS -- To control annuals and knock down top
 growth of perennials, use 1 gallon in 4 gallons of water and spray to wet plants
 thoroughly.

PERENNIAL WEEDS AND GRASSES -- Use 5 to 6 pints per 100 Sq. Ft. and apply
 in sufficient water for uniform coverage. For control of weeds with creeping
 rootstalks, treat soil 6 Ft. beyond visible vegetation to get creeping roots.

HOW TO USE: A gravity flow spray system or a low pressure sprayer giving
 uniform coverage is satisfactory. State or local agricultural authorities for
 specific recommendations as to dosage and time of application and for the control
 of other weeds.

Follow directions carefully. Consult state agricultural experiment station or
 extension service weed specialist for specific timing and method of application,
 weather and crop conditions, mixtures with other chemicals not specifically
 recommended, and other influencing factors in the use of this products which are
 beyond the control of the seller. Buyer assumes all risks of use, storage, or
 handling of this material not in strict accordance with directions given herewith.

NOTE: This material is non-selective and affects all types of vegetation.
 Extreme care should be exercised in its use to prevent killing lawns, shrubs,
 trees, and other valuable plants. This includes the treatment of areas which
 may be underlaid by tree roots.

Due to the varied conditions of weather, soil, types of vegetation, location and
 time of applications, the manufacturer does not guarantee complete kill of
 vegetation or accept responsibility for its use or mis-use.