

BAC-TROL XS

E.P.A. REGISTRATION No. 9839-5

BAC-TROL XS when used as directed will provide residual bacteriostatic and residual self-sanitizing properties to fabric when exposed to high relative humidity or wet contamination.

BAC-TROL XS will inhibit the growth of ammonia producing organisms (B. ammoniagenes) in diapers which act on urine to produce ammonia.

BAC-TROL XS when used as directed will assist in reducing the hazards of cross-infection from laundry.

Rinse empty container thoroughly with water and return to drum reconditioner or destroy by perforating or crushing and bury in safe place.

ACTIVE INGREDIENTS:

| | |
|---|----------------|
| n-alkyl (60% C ₁₄ , 30% C ₁₆ , 5% C ₁₂ , 5% C ₁₈ ,) | |
| dimethyl benzyl ammonium chlorides | 22.5% |
| n-alkyl (68% C ₁₂ , 32% C ₁₄) dimethyl | |
| ethylbenzyl ammonium chlorides | 22.5% |
| Ethyl Alcohol | 11.25% |
| INERT INGREDIENTS:..... | 43.75% |
| | 100.00% |

DANGER

KEEP OUT OF REACH OF CHILDREN. See Right Side Panel for additional precautions and antidote.

Manufactured By

TROST CHEMICAL COMPANY

7013 Gainesborough Drive
Knoxville, Tennessee 37919

NET CONTENTS GALLONS

USE DIRECTIONS

Add BAC-TROL XS in the final rinse cycle at ¾ of an ounce per each 100 lbs of dry laundry. Add fabric softener five minutes after addition of BAC-TROL XS.

DANGER

Keep Out Of Reach Of Children. Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful or fatal if swallowed. Avoid contamination of food.

FIRST AID

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN

Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

[Handwritten signature and date]
7/1/72