

**SANSOFT**  
**SANITIZING-FABRIC**  
**CONDITIONER CONCENTRATE**  
**E.P.A. REGISTRATION No. 9839-6**

Fabric treated as directed will develop residual bacteriostatic activity and residual self-sanitizing activity against Gram positive (S. Aureus) and Gram Negative (E. Coli) organisms, under conditions of high relative humidity or wet contamination.

SANSOFT inhibits the growth of ammonia producing bacteria (B. ammoniagenes) which act on urine in the diapers to produce ammonia.

SANSOFT when used as directed will assist in reducing the hazards of cross-infection from treated fabrics.

SANSOFT softens clothes, thus eliminating chafing, "static cling" out of nylons and other synthetic fabrics. Improves ironing, drying, conditioning, softness, handling and feel.

Rinse empty container with water and discard it. Keep away from children and livestock.

RESIDUAL BACTERIOSTAT AND RESIDUAL SELF-SANITIZER UNDER CONDITIONS OF HIGH RELATIVE HUMIDITY OR WET CONTAMINATION

**ACTIVE INGREDIENTS:**

n-Alkyl (60% C <sub>14</sub> , 30% C <sub>16</sub> , 5% C <sub>12</sub> , 5% C <sub>18</sub> ) dimethyl benzyl ammonium chlorides .....	8.0%
n-Alkyl (68% C <sub>12</sub> , 32% C <sub>14</sub> ) dimethyl ethylbenzyl ammonium chlorides .....	8.0%

<b>INERT INGREDIENTS:</b> .....	84.0%
	100.0%

**DANGER**

**KEEP OUT OF REACH OF CHILDREN.** See Right Side Panel for additional precautions and antidote.

Manufactured By

**TROST CHEMICAL COMPANY**

7013 Gainesborough Drive  
 Knoxville, Tennessee 37919

NET CONTENTS ..... LBS.

**USE DIRECTIONS:**

FOR COMMERCIAL & INSTITUTIONAL LAUNDRING add SANSOFT in final rinse cycle at 2.5 oz. per each 100 lbs. of dry laundry. Run this rinse for 5 minutes at temperatures between 80 degrees and 120 degrees F.

**DANGER**

Keep Out Of Reach Of Children. Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful or fatal if swallowed. Avoid contamination of food.

**FIRST AID**

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

**NOTE TO PHYSICIAN**

Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

9839-6