BACSOFT when used as directed will provide residual bacteriostatic activity against Staphylococcus Aureus and Brevibacterium ammoniagenes under conditions of high relative humidity or wet contamination.

BACSOFT inhibits the growth of ammonia producing bacteria (B. ammoniagenes) which act on urine in diapers to produce ammonia.

BACSOFT when used as directed will assist in reducing the hazards of cross-infection from treated fabrics.

BACSOFT softens clothes, thus eliminating chafing, "Static Cling" out of nylons and other synthetic fabrics. Improves ironing, drying, conditioning, softness, handling and feel.

Rinse empty container with water and discard it. Keep away from children and livestock.

BACSOFT

BACTERIOSTAT-FABRIC CONDITIONER CONCENTRATE

E.P.A. REGISTRATION No. 9839-7

Residual Bacteriostat under conditions of high relative humidity or wet contamination.

ACTIVE INGREDIENTS:

INIERT INICREDIENTS.	02.00/
ethylbenzyl ammonium chlorides	4.0%
n-Alkyl (68% C _{1.2} , 32% C _{1.4}) dimethyl	
dimethyl benzyl ammonium chlorides	4.0%
n-Alkyl (60% C ₁₄ , 30% C ₁₆ , 5% C ₁₂ , 5% C ₁₈)	

DANGER

KEEP OUT OF REACH OF CHILDREN. See Right Side Panel for additional precautions and antidote.

Manufactured By

TROST CHEMICAL COMPANY

7013 Gainesborough Drive Kr.oxville, Tennessee 37919

NET CONTENTS LBS.

USE DIRECTIONS

FOR COMMERCIAL & INSTITUTIONAL LAUN-DERING add BACSOFT in final rinse cycle at 2.5 oz. per each 100 lbs. of dry laundry. Run this rinse for 5 minutes at temperatures between 80 degrees and 120 degrees F.

DANGER

Keep Out Of Reach Of Children. Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful or fatal if swallowed. Avoid contamination of food.

FIRST AID

100.0%

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN

Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

