

DIRECCIONES DE USO Sostenga el atomizador verticalmente contra la pared de la cocina o la estancia. Apriete la válvula hacia abajo y rocie el cuarto indicado.

PARA EL CONTROL DE INSECTOS CASEROS: Para control de moscas, mosquitos, pequeños pollos, voladoras, abejas o avispas, cierra las puertas y ventanas. Remueve los animales caseros, pájaros, y cubre los acuarios de peces, antes de rociar. Manténgase el cuarto cerrado por al menos 10 minutos después de rociar. No se quede en el área tratada y ventile las áreas después de que el tratamiento haya sido completado.

Para matar cucarachas y peces lo dorado. Rocíe directamente en las grietas y rendijas y otras sitios apartados. Dirigir el rociado directamente a los insectos cuando aparezcan. Una cocina de tamaño regular requiere que se rocíe a rededor de 1 a 2 pies lineales por segundo. Para matar hormigas. Rociar los insectos directamente y los senderos y otros lugares donde las hormigas se congregan.

AVISO

DARÑINO SI SE INGIERE: EVITE EL CONTACTO CON PIEL Y OJOS. EVITE LA INHALACION DEL PRODUCTO ROCIADO. NO ROCAR CERCA O A LLAMA VIVA. EVITAR CONTAMINACION DE ALIMENTOS PARA ANIMALES O COMESTIBLES.

SI SE UAN EN PLANTAS PROCESADORAS DE ALIMENTOS O AREAS DONDE SE PREPARAN ALIMENTOS COMERCIALMENTE LAS ALIMENTOS TIENEN QUE SER REMOVIDAS O CUBIERTAS DURANTE EL TRATAMIENTO. TODAS LAS SUPERFICIES DONDE SE PRODUZCAN ALIMENTOS TIENEN QUE SER CUBIERTAS DURANTE EL TRATAMIENTO O LAVADAS CONCERNIENDAMENTE ANTES DE VOLVER A USARSE. APLICAR SOLAMENTE CUANDO EL ESTABELO MIENTO NO ESTE EN OPERACION.

Este producto es tóxico a los pescados. Manténgase fuera de los alcances de los niños. Al aplicar este producto, mantenerlo lejos de la boca y los ojos.

Contenidos bajo presión. No aplique sobre o almacene cerca de calor intenso. Si se expone a temperaturas superiores a 120°F, puede explotar. Nunca tire el atomizador al fuego o incinerarlo.

Net Wt. 13 Oz.

RITE-OFF

INSTITUTIONAL INSECT SPRAY

Contains A Pleasant Room Freshener TOO!

TAMBIEN CONTIENE UN
REFRESCANTE DE CUARTO

Spray

ACTIVE INGREDIENTS

Pyrethrins	0.2%
Piperonyl Butoxide Technical	1.0%
Petroleum Distillate	0.8%
INERT INGREDIENTS	98.0%

*Equivalent to 0.8% of (butylbenzyl) propoxy (ipiperonyl) ether and 0.2% of related compounds.

WARNING

KEEP OUT OF REACH OF CHILDREN.

Contents under pressure.

See other cautions on Back Panel.

AVISO:

MANTENGASE FUERA DEL ALCANCE DE LOS NIÑOS

CONTENIDO BAJO PRESIÓN

VEANSE OTRAS AVISO AL COSTADO

EPA Reg. No. 9852-31

EPA EST. No. 9852-N.Y.-1

DIRECCIONES FOR USE
Hold container upright, close to wall. Press valve down and spray. Hold 12 inches from wall.

**FOR THE CONTROL OF
HOUSEHOLD INSECTS**
To control houseflies, mosquitoes, small flying moths, gnats, or wasps; shut doors and windows and turn off lights. Remove nets, birds, and insects in rooms before spraying. Keep them closed for at least 10 minutes after spraying. Do not remain in treated areas and ventilate the area after treatment is completed.

To kill roaches & silverfish; hold, direct, into cracks, crevices and other small places. Direct spray at insects when they appear. An average size kitchen requires spraying 10 linear feet per second. To kill ants; Spray directly on insects and ant trails and other places where ants congregate.

PHYSICAL HAZARD

Harmful if swallowed. Do not spray near or toward an open flame. Avoid contamination of feed and foodstuffs. Avoid contact with skin and eyes. Avoid inhalation of spray mist. If used in food processing plants or commercial food preparation areas, foods should be removed or covered during treatment. All food processing surfaces should be cleaned during treatment or thoroughly cleaned before using. Apply only where facility is not in operation. This product is toxic to fish. Keep out of drain streams or ponds. Apply this product only as specified on the label. Avoid skin contact. Avoid inhalation of mist.

CHEMICAL HAZARDS

Contents under pressure. Do not puncture. Do not use or store near heat, open flame. Exposure to temperatures above 120°F may cause bursting. Never throw empty containers or rinsing water.

RITE OFF, INC.
1545 5th Industrial Ct
Bay Shore, NY 11706

4852-31

FDA Reg. No 41824-1, A Est. No 41824-1001
A Concentrated PYRENONE Micro Emulsion

Clear

-Stable

-Leaves No Oily or Objectionable Residues

Effective, Fast Acting Insecticide

For Use in Homes, Institutions, Restaurants

Food Processing Plants, Dairies, Livestock and Outdoors for Mosquito Control

Apply with Conventional Mechanical or Compressed Air Equipment

Active Ingredients:

Pyrethrins.....	0.1%
*Piperonyl Butoxide, Technical..	1.0%
Petroleum Distillate.....	0.4%
INERT INGREDIENTS.....	8.5%
	100.0%

*Equivalent to 0.3% (butylcarbitol)(6-propylpiperonyl) ether and 0.2% related compounds.

PYRENONE Registered trademark of Fairfield American Corporation

KEEP OUT OF REACH OF CHILDREN

CAUTION

See side panel for additional precautionary statements

DO NOT TRANSPORT OR STORE UNDER 32°.

NET CONTENTS ONE GALLON

BEST DOCUMENT AVAILABLE

AUGUST 1 PICTURES

INSTRUCTIONS FOR USE

INDOOR APPLICATION

The use of this product in food processing establishments should be confined to time periods when the plant is not in operation. Food should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using.

Vacate treated area and ventilate before reoccupying.

For maximum effectiveness, a combination of spot surface treatment and fogging is recommended.

SURFACE SPRAYING: To control roaches, ants, silverfish, spiders, crickets, clover mites, cheese mites, granary weevils, rice weevils, confused flour beetles, sawtoothed grain beetles, spider beetles, cigarette beetles, drugstore beetles, meal worms, grain mites, and cadelles: Use a good sprayer adjusted to deliver a coarse wet spray. Direct the spray into hiding places, cracks and crevices, under pallets, around containers of stored foods, around the base of machinery, behind shelves and drawers. For silverfish, spray bookcases. For ants, spray trails, nests and points of entry. If surface application only is to be used, spray floors, walls, and other surfaces applying at a rate of 1 gallon to 750 square feet of surface.

To control carpet beetles: Spray edges of carpeting and under carpeting and rugs. Make localized application to floor and baseboards. Spray directly into cracks, closets and infested areas of shelving. Repeat treatment as needed.

To control bedbugs: Spray mattresses lightly, particularly tufts and seams. Take beds apart and spray in all joints. Repeat treatment as needed.

To control clean animal white flies: Thoroughly spray infested areas, etc., resting quarters, nearby cracks and crevices, along and behind baseboards, partitions, window and door frames, and in calico areas or floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Concurrent treatment of animals with an approved Insecticide is recommended. Repeat treatment as needed.

To control clothes moths: Remove any infested articles from storage, brush thoroughly and air for several hours in sun light, if possible. Apply spray liberally to empty chests, closets, bureaus, and other storages, directing spray into cracks, joints, and crevices. After airing, infested articles may be treated lightly keeping nozzle at least three feet from fabric to avoid staining.

SPACE SPRAYING: Close doors and windows and shut off ventilating systems. Apply with a good machine adjusted to deliver a fine mist or fog. For rapid control of flies, fruit flies, gnats, mosquitoes, skipper flies, wasps, hornets, Angoumols, grain moths, Mediterranean flour moths, Indian meal moths and tobacco moths: direct spray at an upward angle, distributing it uniformly throughout the entire area at a rate of 3-6 ounces per 1000 cubic feet of space.

FOGGING: To kill exposed or accessible stages of the above named crawling insects, use a rate of one gallon to 20,000 to 40,000 cubic feet of space. Contact insects directly whenever possible.

ON ANIMALS: To protect cattle and horses from horn flies, house flies, mosquitoes and gnats, apply a light mist sufficient to wet the surface of the hair. To control stable flies, horse flies, etc.

lies, apply at a rate of 1/2 ounces per adult cow or horse sufficient to wet the hair thoroughly. Repeat treatment daily or at intervals necessary to give continued protection.

To control Blood sucking Lice apply to the infested areas of the animal using a stiff brush to get spray to the base of the hair. Repeat every 2 or 3 weeks if required.

OUTDOOR ADULT MOSQUITO CONTROL: Apply as a fine mist to thoroughly treat the entire area. For best results, treat when the wind velocity does not exceed 5 miles per hour. Treat shrubbery or vegetation where mosquitoes may rest, low marshy places, stagnant pools or ponds and shore lines. Apply at a rate of 1½ gallons per acre (4½ ounces per 1000 square feet).

Using truck-mounted conventional misting equipment such as Rotomist models 51C, 11 or 100 FT, traveling at 6 m.p.h. to treat a 300 foot swath, calibrate equipment to deliver 8.5 gallons per minute.

RECAUTIONARY STATEMENT

HAZARD TO HUMAN AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed or absorbed through skin. Avoid breathing vapors or spray mist. Avoid contact with skin and eyes. In case of contact, immediately flush eyes or skin with plenty of water. Obtain medical attention if irritation persists.

Do not apply directly to food. In commercial food handling facilities, cover or remove any food and food processing equipment during application. Do not apply while food processing is in progress. After space spraying in food processing plants, wash all equipment with potable water, benches, shelving, etc., where exposed food will be handled. In the home, all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use.