

DIRECTIONS FOR:

To Use: Hold nozzle 12" away from you and press actuator button spraying as directed.

To Treat Animals: To protect cattle from attacks of stable flies, horse flies, and ticks, spray horses, mares, and foals. For horses, mosquitoes and ticks, spray around the body. Be careful to spray back, withers and between forelegs. To protect from face flies, spray the face and head, but do not spray into the eyes. Repeat treatment as necessary when flies are troublesome.

In Animal Quarters & Milk Rooms: For rapid control of House flies, Mosquitoes, Gnats, Wasps and Small Flying Moths, close windows and doors and spray at the rate of one second per 1000 cubic feet of space. For best results, keep area closed for 10 minutes after spraying. Do not re-ventilate area and ventilate after treatment is completed.

When using this product in food processing plants, foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. When using this product in these areas, apply only when facility is not in operation.

Remove pets, birds and cover fish aquariums before spraying.

Flies, Mosquitoes, Gnats, Wasps and Small Flying Moths: Close all doors and windows and spray room at the rate of 1 second per 1000 cubic feet. Do not spray to all parts of the room, especially windows and other light sources which attract these insects. Keep nozzle in constant motion. Keep room closed for 10 to 15 minutes after treatment. Repeat treatment as necessary.

Beaches, Water Bugs, Silverfish, Crickets, Spiders, Centipedes, and Sow Bugs: Spray into hiding places, such as cracks and crevices, behind sinks, washbasins and toilet tanks, and floors, around drains and plumbing, with this spray whenever possible. Repeat treatment as necessary.

Ants: Spray trails, nests, and points of entry. Spray on ants where possible. Repeat as necessary.

Indian Meal Moth, Anguonous Grain Moth: Close doors and windows, direct spray into all parts of room, especially around and stored food product containers, and around and under bins, trays, 7 to 10 seconds for each 1000 cubic feet of space. Keep area closed for 15 minutes following treatment. Repeat application weekly or as needed. Infested stored food products should be fumigated or treated by other effective methods using an approved product intended for this purpose.

Exposed areas of Confused Flour Beetle, Saw Toothed Grain Beetle, Red Flour Beetle, Face Weevil, Granary Weevil, Grain Mites: Spray on exposed areas, such as cracks and holes, and behind a wall of paper or cloth to reach into other areas where the insects tend to hide. Repeat spray at second, fourth, sixth, eighth and tenth days. Keep room closed for 15 minutes to 1 hour after treatment. Repeat treatment as necessary.

Carpenter Beets: Spray directly on holes, floors, baseboards, shelves and cabinets. Repeat treatment as necessary.

Bees: Spray in the area where they are nesting. Treat by spraying into the hole and allowing the bees to fly out. Repeat treatment as necessary.

Flea: Spray the pet, ears, neck, and between the legs, when pets are kept. Repeat as necessary. Use only a recommended pet spray on pets.

Manufactured in U.S.A. by
RITE OFF CORPORATION
163 Dupont Street, Plainview, N.Y. 11803

Cameraman's Note

Poor Copy

RITE-OFF

DAIRY AND INDUSTRIAL AEROSOL

EPA Est. No. 9852 N.Y. 1

EPA Reg. No. 9852-42

Contains PYRENONE™

Industrial Strength — Low Dosage Required

For Use on Beef and Dairy Cattle,
in Animal Quarters, and Milk Rooms.

KILLS AND REPELS Stable Flies, Horse Flies, Face Flies, Deer Flies, House Flies, Horn Flies, Mosquitoes and Gnats

For use in Schools, Hospitals, Institutions, Dairies, Supermarkets, Hotels, Restaurants, Druggists, Grocers, Drug Stores, Food Plants, and Animal Quarters.

ACTIVE INGREDIENTS

Pyrenone	6.5
Pyrethrin, natural, Technical*	4.0
Piperonyl Butoxide	15.5
INERT INGREDIENTS	30.0

*Equivalent to 1.7% of active ingredient. 10 percent pyrethrin ether and 1.5% of active ingredient.

PYRENONE is a registered trademark of E. I. du Pont de Nemours and Company.

CAUTION

KEEP OUT OF REACH OF CHILDREN

CAUTION: Do not apply to food, feed, or food products. Any exposed surfaces on food, feed, or food products should be thoroughly cleaned before use. Do not use in areas where fish or reptiles may be exposed to the spray.

WARNING: Contents under pressure. Do not puncture. Do not use or store near heat or open flame. Exposure to temperatures above 130 F. may cause bursting. Never throw container into fire or incinerator.

NET WT. 1 lb. 2 oz.

ACCEPTED

MAY 21 1975

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under 4957-42