

## CAUTION

Harmful if swallowed. Avoid prolonged breathing of dust or spray mist. Avoid prolonged or repeated contact with skin; wash thoroughly after using. Avoid contamination of feed and foodstuffs. Keep out of reach of children.

## NOTICE

Seller's guaranty shall be limited to the terms of the label and subject thereto. The buyer assumes any risk of damage to persons or property arising out of use or handling, and accepts the product on these conditions.

### ANIMALS

**BEEF CATTLE: LICE** — 1 gallon to 100 gallons of water or 6½ oz. per 5 gallons of water.

**TICKS** — 1 - 2 gallons per 100 gallons of water or 6½ - 13 oz. per 5 gallons of water. Repeat application only as needed at 2 week intervals.

**HORN FLY** — 1 - 1½ gallons to 100 gallons of water or 6½ - 13 oz. per 5 gallons.

Apply complete coverage spray. One treatment may be sufficient. Repeat application only as need at 2 week intervals. DO NOT use on animals less than 1 month old. Avoid contamination of feed or food.

**DAIRY CATTLE (Non-Lactating)** use 1 - 1½ gallons per 100 gallons of water for dip and spray. DO NOT treat during lactation or spray within 2 weeks of freshening. DO NOT dip dairy animals within 2 months of freshening. DO NOT apply to calves under one month of age. DO NOT apply to lactating dairy cattle.

**SWINE: LICE** — 1 gallon to 100 gallons of water or 6½ oz. per 5 gallons of water. Apply as a complete spray to animals, pens and litter. One treatment may be sufficient. Repeat application only as needed.

### DOGS & CATS DIRECT APPLICATION Sprays

For control of fleas apply spray of 1 oz. of Malathion 57% Emulsifiable Liquid per 1 gal. of water. Wet the animal thoroughly using a hand sprayer.

If the animal has free run, treatments may have to be made every second or third week. To avoid unnecessary applications, animal should be closely examined for fleas before retreating. If the animal is confined to an area free of fleas, one treatment may be sufficient.

### ANIMAL QUARTERS Sprays

For control of fleas in kennels, pens, yards, lawns and under houses. Remove manure or debris before spraying.

Use spray of 5 ozs. of 57% Emulsifiable Liquid per 1 gal. of water at rate of 1 gal per 1000 sq. ft. Use garden sprayer.

### ORNAMENTALS

For Oyster Shell Scale and Lace Bugs, use 1 pint. For Spider Mites, Aphids, White Flies, Mealybugs, Thrips, Japanese Beetle Adults, European Pine Shoot Moth, Four-Lined Plant Bug, Tarnished Plant Bug, Rose Leafhopper, Potato Leafhopper, Scurfy Scale and Euonymus Scale, use 1½ pints. For Bagworm, Tent Caterpillar, Boxwood Leaf Miner, Birch Leaf Miner, Oak Kermes, Florida Red Scale, Fletcher Scale, Juniper, Azalea and Magnolia Scales, use 2 pints. For Black Scale crawler, Soft Brown and Monterey Scales, use 2½ pints. Apply thorough full coverage sprays as needed. Note: DO NOT apply to petunias, certain ferns including Boston, Maidenhair, Pteris, and some species of Crassulaceae since plant injury may occur.

All above rates per 100 gallons of water.


# M

ACTIVE ING

Malathi  
dieth  
Aromat  
INERT INGR

# CAI

U.S. REG. NO. 9859-E



**HOUSEHOLD INSECTS**

For the control of ants, clover mites, crickets, earwigs, pantry pests, roaches, scorpions, silverfish and spiders in and about homes, and food-processing plants, use Malathion 57% Emulsifiable Liquid at the rate of **6½ fluid ounces per gallon** of water. Apply by means of a coarse spray or paint brush to window sills, baseboards, under sinks, drainboards, stoves, cracks and crevices and to other places frequented by insects.

For control of bedbugs in dwellings use Malathion 57% Emulsifiable Liquid at a rate of **1 to 2 fluid ounces per gallon** of deodorized water. Apply lightly to all mattress surfaces in sufficient quantity to "mist" the fabric and generously to beds and woodwork with special care taken to wet all possible hiding places. Remove fish, birds and pets before treating.

Avoid contamination of food, utensils, milk, milk equipment and water.

Water emulsions may stain finished surfaces such as wallpaper, furniture, baseboards, etc.

Repeat applications as necessary.

\*Pantry pests such as the exposed stages of: saw-toothed grain beetle, flour beetle, rice weevil, cigarette beetle, drug store beetle and Indian meal moth.

This product is toxic to fish.

Keep out of lakes, streams and ponds.

# MALATHION EM-5

**An Emulsifiable Concentrate containing  
5 lbs. of Actual Malathion per Gallon**

ACTIVE INGREDIENTS:	BY WEIGHT
Malathion (0,0-dimethyl dithiophosphate of diethyl mercaptosuccinate)	57.0%
Aromatic petroleum derivative solvent	32.0%
INERT INGREDIENTS:	11.0%
TOTAL	100.0%

**CAUTION: KEEP OUT OF REACH OF CHILDREN**

SEE SIDE PANEL FOR OTHER CAUTIONS.