

8-30-72
 UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND RODENTICIDE ACT
 FOR ECONOMIC PEST CONTROL REGISTERED
 UNDER NO. 10163-40

PROKIL Cryolite 75 Dust

ACTIVE INGREDIENTS	%by Wt.
SODIUM FLUOALUMINATE*	72.0
INERT INGREDIENTS	28.0
	100.0
*Flourine expressed as elemental	40.6

CAUTION
 KEEP OUT OF REACH OF CHILDREN
 Avoid Inhalation of Dust

THIS PRODUCT HIGHLY TOXIC TO BEES exposed to treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

NET WEIGHT 50 LBS.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS, WARNINGS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS. USES, TIMING AND DOSAGE MAY VARY AS A CONSEQUENCE OF LOCAL WEATHER OR CONDITIONS. WE RECOMMEND REFERENCE TO STATE AGRICULTURAL EXTENSION SERVICE CONCERNING SPECIFIC USES.

KEEP PESTICIDE IN ORIGINAL CONTAINER!

DO NOT REUSE CONTAINER. DESTROY WHEN EMPTY.

Use at rate of 17 to 40 pounds per acre depending on plant size and insect population. Repeat application as necessary. Remove residues remaining at harvest by washing, brushing or other effective means. Do not apply under conditions involving possible drift to food, forage or other planting that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

6-72-113A 1M

EPA Reg. No. 10163-40 AA

BEANS: Mexican Bean Beetles, Bean Leafroller--Repeat at two week intervals.

CABBAGE, CAULIFLOWER, BROCCOLI: Flea Beetles, Cabbage Caterpillars — DO NOT APPLY to Broccoli after the edible parts start to form. DO NOT APPLY to Cabbage after the heads start to form. DO NOT APPLY to Cauliflower after the seedling stage.

CARROTS: Vegetable Weevil.

CUCUMBERS: Striped Cucumber Beetle, Twelve-Spotted Cucumber Beetle, Flea Beetle — Dust both sides of foliage at first signs of Beetles.

EGGPLANT: Flea Beetle — Dust thoroughly.

LETTUCE: Climbing Cutworms, Alfalfa Looper, Corn Earworm — Apply when worms first appear. DO NOT APPLY after the seedling stage.

MELONS: Cucumber Beetle — Dust both sides of foliage. Repeat if necessary.

PEPPERS: Pepper Weevil — Dust thoroughly at first set of fruit. Repeat every 5 days until about 6 applications have been made.

STRAWBERRIES: Leaf Beetle — Dust both sides of leaves.

TOMATOES: Flea Beetle — Apply when plants are first set out and repeat as necessary. If applied after fruit starts to set, remove residue at harvest by washing, bushing or other effective means.

NOTICE ON CONDITIONS OF SALE

Our recommendations for use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice. The buyer must assume all responsibility, including injury or damage, resulting from its misuse or such, or in combination with other materials.

CALIFORNIA (Plant)
 P. O. Box 406
 340 East Main Street
 Calipatria, Calif. 92233
 Telephone: 348-5150

The Dune Company

AGRICULTURAL CHEMICALS

ARIZONA
 Route 1, Box 33
 Somerton, Arizona 85350
 Telephone: 627-2479

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. 10163-41

PROKIL Cryolite 96

ACTIVE INGREDIENTS	% by Wt.
Sodium Fluoaluminate*	96
INERT INGREDIENTS	4
	<hr/> 100

*Fluorine expressed as elemental 50%

CAUTION

KEEP OUT OF REACH OF CHILDREN
AVOID INHALATION OF DUST

THIS PRODUCT IS HIGHLY TOXIC TO BEES exposed to treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

NET WEIGHT 50 LBS.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS, WARNINGS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS. USES, TIMING AND DOSAGE MAY VARY AS A CONSEQUENCE OF LOCAL WEATHER OR CONDITIONS. WE RECOMMEND REFERENCE TO STATE AGRICULTURAL EXTENSION SERVICE CONCERNING SPECIFIC USES.

KEEP PESTICIDE IN ORIGINAL CONTAINER!

DO NOT REUSE EMPTY CONTAINER. DESTROY IT BY BURYING WITH WASTE or BURNING. STAY AWAY FROM SMOKE or FUMES.

This product is formulated for application by ground or air equipment.

Use at rate of 25 to 50 pounds per acre depending on plant size and insect population. Repeat application as necessary. Remove residues remaining at harvest by washing, brushing or other effective means. Do not apply under conditions involving possible drift to food, forage or other planting that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

7-72—164 PROOF

EPA Reg. No. 10163-41 AA

BEANS—Mexican bean beetle, bean leaf beetle, bean leaf roller, corn earworm, cucumber beetles.

CABBAGE, COLLARDS, MUSTARD, TURNIPS, BROCCOLI, RADISHES, and CAULIFLOWER—Imported cabbage worm, cabbage looper, diamond-back moth caterpillar, yellow striped armyworm, climbing cut worms, corn earworm, flea beetles.

CANTALOUPE, WATERMELON, CUCUMBER and SQUASH—Melon worms, pickle-worm, cucumber beetles.

TOMATO—Hornworm, fruitworm, pinworm, blister beetle.

APPLES, PEACHES, PEARS, SHADE TREES, and ORNAMENTALS—Flea weevil, codling moth, gypsy moth caterpillar, Fuller's rose beetle, plum curculio.

GRAPEFRUIT, LEMONS, ORANGES—Orange Tortrix.

CRANBERRY—Cranberry weevil, gypsy moth caterpillar, false army worm.

LETTUCE—Cabbage looper.

GRAPES—Grape leaf folder.

STRAWBERRIES—Strawberry leaf roller.

PEPPERS—Pepper weevil.

LIMITATIONS—Do not use in combination with lime or compounds containing free lime. Do not apply immediately before harvest. Do not apply to bush and vine fruits after fruit starts to form. Do not apply to cabbage, broccoli or grapes after heads or edible parts start to form. Do not apply to collards, mustard, cauliflower or lettuce after the seedling stage or to turnips after seedling stage if tops are to be used for food.

Remove excess residues on edible portions of fruit and vegetables by washing, brushing or other effective means.

NOTICE ON CONDITIONS OF SALE

Our recommendations for use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice. The buyer must assume all responsibility, including injury or damage, resulting from its misuse as such, or in combination with other materials.

CALIFORNIA (Plant)
P. O. Box 406
340 East Main Street
Calipatria, Calif. 92233
Telephone: 348-5150

The Dune Company

AGRICULTURAL CHEMICALS

ARIZONA
Route 1, Box 33
Somerton, Arizona 85350
Telephone: 627-2479

SUGGESTED USE

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS, WARNINGS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS. TIMING AND DOSAGE MAY VARY AS A CONSEQUENCE OF LOCAL WEATHER OR CONDITIONS; WE RECOMMEND REFERENCE TO AGRICULTURAL EXPERIMENT STATIONS ON SPECIFIC USAGE

In order that pesticidal residues on food and forage crops will not exceed tolerances established by the Federal Food and Drug Administration, use only at recommended rates and intervals, and do not apply closer to harvest than specified. Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals or onto adjoining food, fiber or pasture crops. **The grower is responsible** for residues on his crops as well as for damages caused by drift from his property to that of others. Consult state agricultural extension service or state agricultural experiment stations for additional information, as the number, and rate of applications needed will vary with local conditions.

NOTICE ON CONDITIONS OF SALE

Our recommendations for use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice. The buyer must assume all responsibility, including injury or damage, resulting from its misuse as such, or in combination with other materials.

12-73 1 1/2 M

E P A Reg. No. 10163-43 AA

PROKIL MALATHION 4 PERTHANE* 4

ACTIVE INGREDIENTS	By Wt.
Malathion (O, O diethyl dithiophosphate of diethyl-mecaptosuccinate)	42.62%
Diethyl Diphenyl Dichloroethane	37.31%
Related Compounds	5.09%
Aromatic petroleum derivatives	5.97%
INERT INGREDIENTS	90.1%
Total	100.00%

*Perthane is a trademark Reg. U.S. Patent Office by Rohm & Haas Company.

Contains 4 lbs. Malathion & 4 lbs. Perthane per gallon

CAUTION

KEEP OUT OF REACH OF CHILDREN
NOT FOR USE OR STORAGE IN OR AROUND THE HOME

See side panel for additional precautions.

NET CONTENTS _____ GALLONS

CALIFORNIA (Plant)
P.O. Box 458
340 East Main St.,
Telephone: 348-2262
Calipatria, CA 92233

The Dine Company

AGRICULTURAL CHEMICALS

ARIZONA
P.O. Box 5696
2760 Cty. 15 St.
Telephone: 344-0040
Yuma, Ariz.

ACCEPTED
APR 29 1975
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 10163-43

Harm
or
Avoic
wil
We
Do n
NOTE
use n
2 PA

Do
or fe
ma in
Do
applic
waste
any c
this l
Thi
ment
be ok