

NET WEIGHT

50 LB.



EPA Reg. No. 9859-51

CHLORDANE 10 GRANULAR

A SOIL INSECTICIDE

ACTIVE INGREDIENTS:

Technical Chlordane*	-----	10.0%
INERT INGREDIENTS:	-----	90.0%
TOTAL	-----	100.0%

*Equivalent to 6% Octachloro-4, 7 methano-tetrahydroindane and 4% related compounds.

CAUTION

- ⊙ AVOID BREATHING DUST.
- ⊙ AVOID CONTACT WITH SKIN.
- ⊙ WASH WITH SOAP AND WATER AFTER EXPOSURE.
- ⊙ DO NOT CONTAMINATE FOOD AND FEED.

OBSERVE THE FOLLOWING USE PRECAUTIONS

Observe use limitations given on the label for specific crops.

This product is toxic to fish, birds, and other wildlife. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Do not use this product on feed or forage to be fed to dairy animals or to livestock being finished for slaughter.

Consult State Agricultural Experiment Stations or the State Agricultural Extension Service for information about timing of applications.

Do not reuse container. Destroy when empty.

CAUTION Keep Out of Reach of Children

NOTICE: Seller's guarantee shall be limited to the terms of the label and insert thereon. The buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.

MANUFACTURED BY

LINDIA CHEMICAL CO.


LAKELAND, FLORIDA 33851

RECOMMENDATIONS AND DIRECTIONS FOR USE

CORN (Field and Sweet): Broadcast Application - Larvae of White-fringed Beetles - Apply 25 to 40 lb. per acre. Cutworms - Apply 20 to 40 lb. per acre. Apply Chlordane after field is plowed. Disk in promptly. Use higher rate on heavier soils and lower rate on light sandy soils. Use higher rates for heavy infestation of cutworms.

The use of this product to control white-fringed beetle larvae on corn is restricted to 8 southern states (AL, FL, GA, LA, MS, NC, SC, TN).

The use of this product on corn is continued until August 1, 1976 only.



OBSERVE THE FOLLOWING USE PRECAUTIONS

Observe use limitations given on the label for specific crops.

This product is toxic to fish, birds, and other wildlife. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Do not use this product on feed or forage to be fed to dairy animals or to livestock being prepared for slaughter.

Consult State Agricultural Experiment Stations or the State Agricultural Extension Service for information about timing of applications.

Do not reuse container. Destroy when empty.

CAUTION Keep Out of Reach of Children

NOTICE: Seller's warranty shall be limited to the terms of the label and subject thereto. The buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.

MANUFACTURED BY

LANDIA CHEMICAL CO.

LAKELAND, FLORIDA 33801

The use of this product on corn for control of beetle larvae on corn is restricted to the eastern states (AL, FL, GA, LA).

The use of this product on corn is restricted to August 1, 1976 only.

