


NET WEIGHT  
50 LB.



EPA Reg. No. 9859-53

# CHLORDANE 5 GRANULAR

A SOIL INSECTICIDE

ACTIVE INGREDIENTS:	
Technical Chlordane*	5%
INERT INGREDIENTS:	95%
TOTAL	100%

\*Equivalent to 3% octachloro-4, 7 methano-tetrahydroindane and 2% related compounds.

### CAUTION

- AVOID BREATHING DUST.
- AVOID CONTACT WITH SKIN.
- WASH WITH SOAP AND WATER AFTER EXPOSURE.
- DO NOT CONTAMINATE FOOD AND FEED.

### OBSERVE THE FOLLOWING USE PRECAUTIONS

Observe use limitations given on the label for specific crops.

This product is toxic to fish, birds, and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Do not use this product on feed or forage to be fed to dairy animals or to livestock being finished for slaughter.

Consult State Agricultural Experiment Stations or the State Agricultural Extension Service for information about timing of applications.

Do not reuse container. Destroy when empty.

### CAUTION Keep Out of Reach of Children

NOTE: Seller's liability shall be limited to the terms of the label and actual directions. The buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.

MANUFACTURED BY

**LANDIA CHEMICAL CO.**

LAKELAND, FLORIDA 33801


EPA Est. 0859-FL-1

### RECOMMENDATIONS AND DIRECTIONS FOR USE

CORN (Field and Sweet): Broadcast Application- Larvae of White-fringed Beetles - Apply 50 to 80 lb. per acre. Cutworms - Apply 40 to 50 lb. per acre. Apply chlordane after field is plowed. Disk promptly. Use higher rates for heavy infestation of cutworms.

The use of this product to control white-fringed beetle larvae on corn is restricted to 8 southeastern states (AL, FL, GA, LA, MS, NC, SC, TN).

The use of this product on corn is continued until August 1, 1976 only.



duct is toxic to fish, birds, and other  
birds feeding on treated areas may  
d. Keep out of lakes, streams, or ponds.  
Apply where runoff is likely to occur.  
Apply when weather conditions favor drift  
as treated. Do not contaminate water by  
use of equipment, or disposal of wastes.  
Use this product only as specified on this

August 1, 1975 only.

Use this product on feed or forage to be  
consumed by dairy animals or to livestock being  
prepared for slaughter.

Consult State Agricultural Experiment Stations  
and State Agricultural Extension Service for  
information about timing of applications.

Reuse container. Destroy when empty.

**CAUTION Keep Out of Reach of Children**

NOTICE: Seller's warranty shall be limited to the terms of the label, and subject thereto. The buyer  
assumes any risk to persons or property arising out of use or handling and accepts the product on  
these conditions.

MANUFACTURED BY

**LANDIA CHEMICAL CO.**

LAKELAND, FLORIDA 33801

EPA Est. 0859-FL-1

**ACCEPTED**

MAY 19 1975

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under  
EPA Reg. No. 9859-53