

10 15 1964

Y₃

Net Contents
25 lbs.

F A S C O
POULTRY DUST C-R

Product
No. 1010

For Control of External Parasites
Contains 0.5% CO-RAL*

Active Ingredients	BY WT.
O,O-Diethyl O-3-chloro-4-methyl -2-oxo-2H-1-benzopyran-7-yl phosphorothioate	0.5%
Inert Ingredients	99.5%

FASCO Poultry Dust is an effective dust formulation designed for use against Northern Fowl Mite, Lice, Poultry Red Mite, and Flies which infest poultry. Direct application of the dust may be made to the birds by means of rotary or hand dusters, puff dusters, or by sprinkling from a can as directed below.

DIRECTIONS

Use contrary to these directions may cause injury to poultry, personal injury, excessive residues, unsatisfactory control or other unintended consequences.

DIRECT APPLICATION TO POULTRY (Chickens, Turkeys, Ducks, Geese) - For the control of Northern Fowl Mite and Lice, dust birds thoroughly, especially under wings and around vent. Supplemental control can be obtained by also treating litter and housing. Normally one pound of dust will treat 30-50 birds. Repeat as necessary but not more often than weekly.

DUST BOX SELF TREATMENT - For the control of Northern Fowl Mite and Lice place 2½ pounds of FASCO POULTRY DUST C-R in a shallow box for self treatment of 50 birds. Supplemental control can be obtained by treatment of litter and housing.

LITTER AND HOUSING TREATMENT - For control of Northern Fowl Mite, Poultry Red Mite, Lice, and flies, apply one pound of dust per 70 square feet of surface area to walls, ceiling, floor, roosts, nests, and adjacent areas. Also apply one pound of dust per 20 square feet of litter area. Force dust into all cracks and crevices. Repeat as necessary.

PRECAUTIONS

Do not treat birds within 10 days of vaccination or other stress influence.

Avoid contamination of feed, water, and water utensils. Provide thorough ventilation while dusting.

Do not use on cats.

CAUTION

Avoid prolonged inhalation of dust. Work after handling.

Do not contaminate feed or food. Keep out of reach of children.

ACCEPTED

FEB 19 1964

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTRY

If misuse results in accidental poisoning, obtain prompt medical aid. Prolonged exposure will result in cholinesterase depression. Atropine Sulfate is antidotal.

CONDITIONS OF SALE

Neither the registrant nor the manufacturer, and neither the distributor nor the retailer, shall be held liable for any injury or damage resulting from the use of this product in accordance with the directions on the label.

ACCEPTED
 2-19-64
 2342-590
 UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND RODENTICIDE ACT
 FOR REGISTRATIONS IN REGISTER-
 ED UNDER NO. SUBJECT
 TO ATTACHED COMMENTS.

