

# CARBAIT-5

Contains Carbaryl Insecticide

**AGRICULTURAL BAIT FOR THE CONTROL OF INSECT PESTS**

**ACTIVE INGREDIENTS:**

Carbaryl (1-naphthyl N-methylcarbamate) . . . . . 5.00%  
 Inert Ingredients . . . . . 95.00%  
 TOTAL . . . . . 100.00%

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

**STATEMENT OF PRACTICAL TREATMENT**

If swallowed: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching finger to back of throat. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.

If inhaled: Remove victim to fresh air. Apply respiration if indicated.

If in eyes: Flush eyes for at least 15 minutes with water. Call a physician immediately.

**PRECAUTIONARY STATEMENTS  
HAZARD TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION:** MAY BE HARMFUL IF SWALLOWED. Do not take internally. Avoid contact with eyes, skin and clothing. Avoid breathing vapors, dust, or spray mist. Wash thoroughly after handling. Wash contaminated clothing before reuse. Flush contaminated eyes with plenty of water and get medical attention if irritation persists. Keep children and pets off treated areas until this material is washed into the soil and grass is dry.

**NOTE TO PHYSICIAN:** Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

**ENVIRONMENTAL HAZARD**

This pesticide is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of wastes.

EPA Est. No. 9859-FL-1

EPA Reg. No. 9859-203

**40 POUNDS NET**

Manufactured By:  
**LANDIA CHEMICAL COMPANY**  
 LAKELAND, FLORIDA 33801

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

**STORAGE:** Store in cool, dry place. (In case of spill, remove spillage with an industrial type vacuum cleaner and wash area with household bleach.)

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Do not reuse bag. Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**DIRECTIONS FOR USE**

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl label(s).

Broadcast applications may be made with either ground equipment or aircraft. Treatments may be repeated as necessary. Dosages from 20 to 40 pounds per acre of CARBAIT-5 will control Cutworms, Armyworms, Crickets, Darkling Ground Beetles, Grasshoppers and Sowbugs.

Crop	Dosage limitations to meet Residue Tolerances	Days Between Final Application and Harvest
Cucumbers, squash melons, peanuts	20 pounds per acre	None
Alfalfa, peas	30 pounds per acre	3 Days
Cotton	30 pounds per acre	7 Days
Beans, carrots, corn, eggplant, okra, peppers, potatoes, tomatoes and sweet corn	40 pounds per acre	None
Asparagus, strawberries	40 pounds per acre	1 Day
Broccoli, Brussels sprouts, cabbage, cauliflower, head lettuce, garden beets (roots), horseradish, parsnips, radishes, rutabagas, and turnips (roots)	40 pounds per acre	3 Days
Sugar beets, collards, endive (escarole), garden beets (tops), kale, leaf lettuce, parsley, spinach, Swiss chard and turnips (tops)	40 pounds per acre	14 Days

**ACCEPTED**

**APR 29 1985**

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 9859-203

**IMPORTANT PRE-HARVEST AND GRAZING USE INFORMATION**

Tolerances established under the Federal Food, Drug and Cosmetic Act permit the sale of crops bearing probable Carbaryl residues when Carbaryl is used in accordance with label directions. If Carbaryl insecticide is used in accordance with label directions, these crops including bean vines, carrot tops, pea vines, sugar beet tops, corn forage and cotton forage may be grazed or harvested for use as feed for dairy and meat animals without resulting in residues in milk or meat.

**CONDITIONS OF SALE**

All statements concerning the use of this product apply only when used as directed. The manufacturer makes no warranties, expressed or implied, concerning this product or its use, which extend beyond the description on the label. Read all directions carefully.

9859-203

04/29/1985

1/1